

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 September 2002 (26.09.2002)

PCT

(10) International Publication Number
WO 02/074981 A2

(51) International Patent Classification⁷: **C12Q 1/00**

Molecular Sciences, South Parks Road, Oxford OX1 3QY (GB).

(21) International Application Number: PCT/GB02/01381

(22) International Filing Date: 21 March 2002 (21.03.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0107123.2 21 March 2001 (21.03.2001) GB
0118952.1 2 August 2001 (02.08.2001) GB

(74) Agents: **ROQUES, Sarah, Elizabeth** et al.; J.A. Kemp & Co., 14 South Square, Gray's Inn, London WC1R 5JJ (GB).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): **ISIS INNOVATION LTD.** [GB/GB]; Ewert House, Ewert Place, Summertown, Oxford OX2 7SG (GB).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **MAXWELL, Patrick, Henry** [GB/GB]; The Henry Wellcome Building of Genomic Medicine, Roosevelt Drive, Oxford OX3 7BN (GB). **PUGH, Christopher, William** [GB/GB]; The Henry Wellcome Building of Genomic Medicine, Roosevelt Drive, Oxford OX3 7BN (GB). **RATCLIFFE, Peter, John** [GB/GB]; The Henry Wellcome Building of Genomic Medicine, Roosevelt Drive, Oxford OX3 7BN (GB). **SCHOFIELD, Christopher, Joseph** [GB/GB]; The Dyson Perrins Laboratory and The Oxford Centre for

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ASSAYS, METHODS AND MEANS

(57) Abstract: A novel class of hydroxylases is described having the amino acid sequence of SEQ ID NO: 2, 4, 6 and 8, and variants and fragments thereof having HIF hydroxylation activity. The polypeptides of the invention have in particular prolyl hydroxylase activity. An assay method monitors the interaction of the HIF hydroxylase with a substrate. Modulators of HIF hydroxylase are provided for use in the treatment of a condition associated with increased or decreased HIF levels or activity or for the treatment of a condition where it is desirable to modulate HIF levels or activity.

ASSAYS, METHODS AND MEANS

Field of Invention

5 The present invention relates to hydroxylases which act on hypoxia inducible factor alpha (HIF- α) and which are involved in the regulation of the cellular turnover of HIF. Compounds, methods and means for the modulation of the activity of these enzymes are provided.

Background of Invention

10 The transcription factor HIF (hypoxia inducible factor) system is a key regulator of responses to hypoxia, occupying a central position in oxygen homeostasis in a wide range of organisms (1). A large number of transcriptional targets have been identified, with critical roles in angiogenesis, erythropoiesis, energy metabolism, inflammation, vasomotor function, and apoptotic/proliferative
15 responses (1). The system is essential for normal development (2, 3), and plays a key role in pathophysiological responses to ischaemia/hypoxia (1). HIF is also important in cancer, in which it is commonly upregulated, and has major effects on tumour growth and angiogenesis (1). The HIF DNA binding complex consists of a heterodimer of α and β subunits (4). Regulation by oxygen occurs through
20 hydroxylation of the α -subunits, which are rapidly destroyed by the proteasome in oxygenated cells (5, 6, 7). This involves binding of HIF- α subunits by the von Hippel-Lindau tumour suppressor protein (pVHL) (8), with pVHL acting as the, or part of the, recognition component for a ubiquitin ligase that promotes ubiquitin dependent proteolysis through interaction with a specific sequence or sequences in
25 HIF- α -subunits (11, 12, 13, 14). In hypoxia, this process is suppressed, so stabilizing HIF- α and permitting transcriptional activation via the HIF α - β .

Disclosure of the invention

30 Investigations by the present inventors have revealed that the interaction between HIF- α and VHL is controlled by oxidation of critical proline residues in the HIF- α protein. In the human HIF-1 α protein these are Pro402 and Pro564, though

the equivalent residue(s) exists in other HIF- α forms and are conserved in *C.elegans*, indicating that these are critical components which have been conserved through evolution.

The data herein demonstrates that hydroxylation of proline residues such as Pro564 in HIF-1 α is mediated by a family of specific prolyl-hydroxylases, referred to here as the HIF hydroxylases, which include the *C. elegans* protein EGL-9 and the human proteins PHD1-3. These enzymes recognise a conserved core LXXLAP motif for prolyl hydroxylation. Different members of the family act differentially on hydroxylation sites within HIF- α and the activity of the recombinant enzymes is directly modulated by oxygen tension, iron availability and cobaltous ions.

The activity of HIF hydroxylases represents a novel target for the control of HIF α . By blocking activity, hydroxylation of HIF α will be reduced, leading to the accumulation of HIF- α in cells. This, in turn, will lead to the promotion or modulation of angiogenesis, erythropoiesis, energy metabolism, inflammation, vasomotor function, and will also affect apoptotic/proliferative responses. Thus, mechanisms which either block, inhibit, reduce or decrease the activity of the HIF hydroxylase, and in particular its prolyl-hydroxylase activity, have therapeutic applications in certain target cells.

Conversely, in hypoxic conditions such as those commonly found in tumours, the lack of hydroxylation may lead to the accumulation of HIF α and the concomitant promotion of angiogenesis and other growth promoting events. Thus mechanisms which either rescue, stimulate, enhance or increase the activity of the HIF hydroxylase, and in particular the prolyl-hydroxylase activity of the enzyme, have different therapeutic applications with respect to certain target cells.

One aspect of the present invention therefore provides an assay method for identifying an agent which modulates the interaction of a HIF hydroxylase with a substrate of the hydroxylase, the method including contacting a HIF hydroxylase and a substrate of the hydroxylase in the presence of a test substance; and, determining the interaction or lack of interaction of the HIF hydroxylase and the substrate.

The HIF hydroxylase and the test substance may be contacted under conditions in which the HIF hydroxylase normally interacts with the substrate of the hydroxylase.

Interaction, or lack of interaction, between the HIF hydroxylase and the substrate may be determined in the presence and/or absence of the test substance. A change, i.e. an increase or decrease in interaction in the presence relative to the absence of test substance being indicative of the test substance being a modulator of said interaction.

Interaction may be determined according to any one of a range of conventional techniques and may include determining the prolyl hydroxylation of the substrate as described below. Interaction in such assays may be any functional interrelation.

Accordingly, the present invention provides an assay method for identifying an agent which modulates the interaction of a hypoxia inducible factor (HIF) hydroxylase with a substrate of the hydroxylase, the method comprising:

- contacting a HIF hydroxylase and a test substance in the presence of a substrate of the hydroxylase under conditions in which the hydroxylase interacts with the substrate in the absence of the test substance; and
- determining the interaction, or lack of interaction, of the hydroxylase and the substrate.

The present invention also provides the HIF hydroxylases themselves. Thus in accordance with the present invention, there is provided a polypeptide comprising:

- (a) the amino acid sequence of SEQ ID NO: 2, 4, 6 or 8;
- (b) a variant thereof having at least 60% identity to the amino acid sequence of SEQ ID NO: 2, 4, 6 or 8 and having HIF hydroxylase activity; or
- (c) a fragment of either thereof having HIF hydroxylase activity.

Preferably, a polypeptide of the invention has prolyl hydroxylase activity.

The present invention also relates to polynucleotides which encode a polypeptide of the invention. Thus, in accordance with another aspect of the invention, a polynucleotide comprises:

- (i) SEQ ID NO: 1, 3, 5 or 7 or a complementary sequence thereto;

(ii) a sequence which hybridises under stringent conditions to the sequence defined in (i);

(iii) a sequence which is degenerate as a result of the genetic code to a sequence as defined in (i) or (ii);

5 (iv) a sequence having at least 60% identity to a sequence as defined in (i);
or

(v) a fragment of any of the sequences (i), (ii), (iii) or (iv), and which encodes a polypeptide having hydroxylase activity or capable of generating antibodies specific for a HIF hydroxylase.

10 The invention also relates to expression vectors comprising a polynucleotide of the invention and antibodies capable of specifically binding a polypeptide of the invention.

The invention also relates to the use of the substances identified in accordance with the assays of the present invention and to the use of inhibitors of the activity of the peptides of the invention in the treatment of a condition or disease
15 associated with altered HIF levels with respect to healthy (or normal) levels, or a condition in which it is desirable to alter HIF activity.

Detailed Description of the Invention

SEQ ID NO: 1 comprises the nucleotide and amino acid sequence for PHD1.

20 SEQ ID NO: 2 comprises the amino acid sequence for PHD1.

SEQ ID NO: 3 comprises the nucleotide and amino acid sequence for PHD2.

SEQ ID NO: 4 comprises the amino acid sequence alone for PHD2.

SEQ ID NO: 5 comprises the nucleotide and amino acid sequence for PHD3.

SEQ ID NO: 6 comprises the amino acid sequence alone for PHD3.

25 SEQ ID NO: 7 comprises the nucleotide and amino acid sequence for EGL-9.

SEQ ID NO: 8 comprises the amino acid sequence alone for EGL-9.

SEQ ID NOs: 9 to 16 represent a number of polypeptides which antagonise the interaction of a HIF α subunit with VHL.

30 SEQ ID NOs: 17 to 22 represent a number of HIF hydroxylase sequence motifs.

SEQ ID NO: 23 provides the amino acid sequence of pVHL minimal binding

domain of HIF-1 α .

SEQ ID NO: 24 provides the amino acid sequence of pVHL minimal binding domain of HIF-2 α .

5 SEQ ID NO: 25 provides the amino acid sequence of pVHL minimal binding domain of HIF- α from *X. laevis*.

SEQ ID NO: 26 provides the amino acid sequence of pVHL minimal binding domain of HIF- α from *D. melanogaster*.

SEQ ID NO: 27 provides the amino acid sequence of pVHL minimal binding domain of HIF- α from *C. elegans*.

10 SEQ ID NOs: 28 to 34 represent the amino acid sequence of a number of synthetic peptides assessed for their ability to block HIF-1 α /pVHL interaction.

SEQ ID NO: 35 comprises the amino acid sequence alone for human HIF- α .

15 SEQ ID NO: 36 comprises the amino acid sequence alone for *C. elegans* HIF- α .

SEQ ID NO: 37 comprises the amino acid sequence of a portion of HIF-1 α which is involved in VHL dependent ubiquitylation and contains an LxxLAP motif.

20 SEQ ID NO: 38 comprises the amino acid sequence of a portion of HIF-2 α which is involved in VHL dependent ubiquitylation and contains an LxxLAP motif.

SEQ ID NO: 39 comprises the amino acid sequence of a second portion of HIF-1 α which is involved in VHL dependent ubiquitylation and contains an LxxLAP motif.

25 SEQ ID NO: 40 comprises the amino acid sequence of the predicted jelly roll core of the *C. elegans* HIF hydroxylase EGL-9.

SEQ ID NO: 41 comprises the amino acid sequence of the predicted jelly roll core of PHD 1.

30 SEQ ID NO: 42 comprises the amino acid sequence of the predicted jelly roll core of PHD 2.

SEQ ID NO: 43 comprises the amino acid sequence of the predicted jelly roll

core of PHD 3.

SEQ ID NO: 44 comprises the amino acid sequence of the predicted jelly roll core of rat SM20.

SEQ ID NO: 45 comprises the amino acid sequence of the prolyl-3-hydroxylase from *Streptomyces*.

SEQ ID NOS: 46 and 47 provide the nucleotide sequences of two primers used to generate a mutagenised ceHIF.

SEQ ID No: 48 provides the amino acid sequence of a possible HIF hydroxylase motif.

HIF hydroxylases

The present invention relates to a family of novel hydroxylases, referred to herein as HIF hydroxylases, functional variants thereof and functional fragments of HIF hydroxylases or of variants thereof. Sequence information for three human HIF hydroxylases termed PHD polypeptides (PHD 1, 2 and 3) are provided in SEQ ID NOS: 1, 3 and 5 (nucleotide and amino acid) and in SEQ ID NOS: 2, 4 and 6 comprising the corresponding amino acid sequence. Sequence information for a *C. elegans* HIF hydroxylase, EGL-9, is provided in SEQ ID NO: 7 (nucleotide and amino acid) and in SEQ ID NO: 8 comprising the corresponding amino acid sequence. A polypeptide of the invention thus consists essentially of the amino acid sequence of SEQ ID NO: 2, 4, 6 or 8 or a variant of any one of these sequences or of a fragment of any one of these sequences or variants of the fragments.

PHD1, 2 and 3 are 2-oxoglutarate dependent non-haem iron-dependent dioxygenases. These dioxygenases have hydroxylase activity, and in particular they mediate hydroxylation HIF 1 α . They are related by sequence to non-haem oxygenases for which crystal structures are known such as proline-3-hydroxylase (Clifton *et al.*, Eur. J. Biochem., 2001, 268, 6625-6636). PHD 1, 2 and 3 are related to EGL9 of *C. elegans* and may also be referred to herein as EGLN 2, 1 and 3 respectively. The PHD 1, 2 and 3 and EGL9 hydroxylases are all considered to be HIF hydroxylases of the invention. The HIF hydroxylases of the invention, and in

particular the human HIF hydroxylases, may also be referred to as EGLN polypeptides.

In a preferred embodiment the HIF hydroxylase of the invention is a prolyl-hydroxylase. Typically the HIF hydroxylase is a human HIF hydroxylase and in particular it is PHD 1, 2 or 3. In a preferred embodiment the various assays, methods, medicaments and other embodiments of the invention employ, or are concerned, with a human HIF hydroxylase and in particular PHD 1, 2 or 3.

A polypeptide of the invention may be in isolated and/or purified form, free or substantially free of material with which it is naturally associated, such as other polypeptides or such as human polypeptides other than that for which the amino acid sequence is encoded by the gene encoding the HIF hydroxylase and in particular the PHD-1, -2 or -3 gene or (for example if produced by expression in a prokaryotic cell) lacking in native glycosylation, e.g. unglycosylated.

It will be understood that the polypeptide may be mixed with carriers or diluents which will not interfere with the intended purpose of the polypeptide and still be regarded as substantially isolated. A polypeptide in substantially purified form will generally comprise the polypeptide in a preparation in which more than 50% e.g. more than 80%, 90%, 95% or 99%, by weight of the polypeptide in the preparation is a polypeptide of the invention. Routine methods can be employed to purify and/or synthesize the proteins according to the invention. Such methods are well understood by persons skilled in the art and, include techniques such as those disclosed in Sambrook *et al*, Molecular Cloning: A Laboratory Manual, Second Edition, CSH Laboratory Press, 1989, the disclosure of which is included herein in its entirety by way of reference.

The term "variant" refers to a polypeptide which shares at least one property or function with the HIF hydroxylases of SEQ ID NOS: 2, 4, 6 or 8 and in particular those of SEQ ID NOS: 2, 4 or 6. A "fragment" of the invention also possesses at least one function or property of the HIF hydroxylase of SEQ ID NO: 2, 4, 6 or 8 and in particular of SEQ ID NOS: 2, 4 or 6. The HIF hydroxylases of the invention are hydroxylases, that is they have the ability to hydroxylate an amino acid residue in a peptide. Preferably, a polypeptide of the invention is capable of hydroxylating one or

more prolyl residues of a peptide substrate. In preferred aspects of the invention, a HIF hydroxylase, variant or fragment in accordance with the invention has the ability to hydroxylate one or more residues of HIF-1 α , preferably a prolyl residue of HIF and in particular Pro 564 and/or Pro 402 of HIF-1 α or a peptide analogue of HIF-1 α or fragment thereof incorporating such a prolyl. Preferably, a variant of a HIF hydroxylase in accordance with the present invention has at least 60% sequence identity with the amino acid sequence of SEQ ID NO: 2, 4, 6 or 8 and in particular with that of SEQ ID NO: 2, 4 or 6.

The present invention also includes active portions, fragments, derivatives and functional mimetics of the polypeptides of the invention. An "active portion" of a polypeptide means a peptide which is less than said full length polypeptide, but which retains hydroxylase activity and in particular maintains HIF hydroxylase activity, preferably HIF prolyl hydroxylase activity. Such an active fragment may be included as part of a fusion protein, e.g. including a binding portion for a different i.e. heterologous ligand.

A "fragment" of a polypeptide generally means a stretch of amino acid residues of at least about five contiguous amino acids, often at least about seven contiguous amino acids, typically at least about nine contiguous amino acids, more preferably at least about 13 contiguous amino acids, and, more preferably, at least about 20 to 30 or more contiguous amino acids. Fragments of the HIF hydroxylase sequence may include antigenic determinants or epitopes useful for raising antibodies to a portion of the amino acid sequence. Alanine scans are commonly used to find and refine peptide motifs within polypeptides, this involving the systematic replacement of each residue in turn with the amino acid alanine, followed by an assessment of biological activity. Such scans may therefore be used in the identification of preferred fragments of the invention.

The polypeptides of the present invention generally have hydroxylase activity, preferably prolyl hydroxylase activity. Thus, the invention also relates to such polypeptides, in particular, for use in assays of hydroxylase activity on substrates such as HIF. The polypeptides may also be used for hydroxylation of

suitable substrates and in particular prolyl hydroxylation of such substrates. A variant or an active fragment of a HIF hydroxylase of the invention may typically be identified by monitoring for hydroxylase activity as described in more detail below. In preferred embodiments the HIF hydroxylase has prolyl hydroxylase activity such as prolyl-4-hydroxylase activity.

Such HIF hydroxylases may be a eukaryotic polypeptide, preferably a mammalian polypeptide, more preferably a human polypeptide.

A HIF hydroxylase preferably has HIF prolyl hydroxylase (HPH) activity and preferably recognises and/or has specificity for the substrate amino acid sequence motif LXXLXP, in particular LXXLAP, or LXXLRP where X is any amino acid i.e. hydroxylates the proline residue of the LXXLXP or LXXLAP motif of a polypeptide which comprises this sequence.

A HIF hydroxylase preferably contains a β -barrel jelly roll structure consisting of a minimum of eight strands. Typically, the jelly roll structure may have eight strands. Figure 9 shows an alignment of various HIF hydroxylases with the locations of the eight β -barrel strands of the jelly roll motif indicated. A diagram of the jelly roll structure is shown in Figure 10.

Preferred HIF hydroxylases comprise the sequence;



where X is any amino acid and n is between 1 and 200, 20 and 150 or 30 and 100 amino acids, for example 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 amino acids.

In especially preferred embodiments, the HXD portion of the motif is located on the second strand of the jelly roll motif of the HIF hydroxylase and the remaining H is on or close to the second strand of the motif.

In some enzymes related to the PHD 1, 2 and 3 enzymes isolated, such as clavaminic acid synthase, the HXD motif is replaced by a HXE motif. Thus the invention also encompasses HIF hydroxylases which have in place of a HXD motif a HXE motif. This may be because the HIF hydroxylase normally has such a motif, or alternatively, because the HXD motif originally present has been replaced by a HXE motif. Thus for any of the HXD motifs described herein, the invention also

encompasses enzymes with a motif where the Aspartic acid residue has been replaced with a Glutamic acid residue.


A suitable HXD[X]_nH motif may comprise the residues His487, Asp489 and His 548 with reference of the Egl-9 sequence. A suitable HIF hydroxylase may thus
5 comprise or include the sequence;



Amino acid residues described herein are numbered according to the EGL-9 sequence (GI5923812), unless otherwise stated. Sequences of the catalytic regions of EGL-9 and other HIF hydroxylases are shown in Figure 9. It will be appreciated that
10 because of variations in sequence, the equivalent or corresponding residues in other HIF hydroxylase sequences may have different numbers. Reference herein to a residue numbered according to the EGL-9 sequence is understood to include the equivalent residue in other HIF hydroxylases.

Preferred HIF hydroxylases may comprise one or more of the following
15 residues; Met473, Asp494, Tyr502, Leu517, Pro532, Asp543, Val550, Arg557.

Such a preferred polypeptide may comprise the following amino acid sequence;



where X is any amino acid residue.

20 Especially preferred HIF hydroxylases may additionally comprise one or more of the following residues;

Arg469, Tyr477, Pro478, Gly479, Asn480, Gly481, Tyr584, Val585, Val488, Asn490, Pro491, Gly495, Arg496, Cys497, Thr499, Ile501, Tyr503, Asn505, Trp508, Asp509, Gly514, Gly515, Phe520, Pro521, Glu522, Asp535, Arg536,
25 Leu537, Phe539, Trp541, Ser542, Arg544, Arg545, Asn546, Pro547, Glu549, Pro552, Ala559, Thr561, Val562, Trp563, Tyr564, Asp566, Glu569, Arg570, Ala573, Ala575, Lys576, Lys578.

Such an especially preferred polypeptide may, for example, comprise the following amino acid sequence;

RXXXMXXXYP GNGXXYVXHV DNPXXDGRGX TXIYYXNXXW
 D(X)₄GGXLX XFPE(X)₉PX XDRLXFXWSD RRNPHEVXP(X)₄RXAXTVWYXD
 XXERXXAXAK XK

where X is any amino acid residue.

- 5 In other preferred embodiments one or more of the following variations of the above sequence may be present. Residue 478 may be Asn, residue 485 may be Ile, residue 496 may be Lys, residue 497 may be Val, residue 515 may be Ser, residue 520 may be Tyr, residue 530 may be Ile, Val or Met, residue 536 may be Lys, residue 537 may be Ile, residue 539 may be Ile, residue 546 may be Thr, residue 559
 10 may be Ser, residue 560 may be Ile, Met or Leu, residue 561 may be Cys and/or residue 564 may be Phe.

- A suitable HIF hydroxylase may comprise a polypeptide sequence selected from the group consisting of SM20 (NCBI Acc No: NP071334), EGL-9 (GI5923812), CG1114 (AAF52050), C1orf12 (NP071334), EGLN1/PHD2
 15 (gi1457146), EGLN2/PHD1 (gi1457148), EGLN3/PHD3 (gi14547150), FALKOR (gi13649965), and FLJ21620 (BAB15101) as shown in Table 1 .

- A polypeptide of the invention may further comprise an amino acid sequence which shares greater than about 60% sequence identity with one of the above amino acid sequences, preferably greater than about 70%, more preferably greater than about
 20 80%, more preferably greater than about 90%, most preferably greater than about 95%. Suitable sequences have prolyl hydroxylase and in particular HIF prolyl hydroxylase activity.

In one embodiment the invention provides a polypeptide having a least 60% sequence identity with the amino acid sequence encoded by the PHD2 (EGLN1) gene.

- 25 Further aspects of the present invention relate to methods for identifying HIF hydroxylases. Such a method may comprise;
 screening a database for an open reading frame encoding a polypeptide comprising the sequence;

M(X)₁₃HXD(X)₄D(X)₇Y(X)₁₄L(X)₁₄P(X)₁₀D(X)₄HXV(X)₆R

expressing said open reading frame to produce said polypeptide; and,
determining the ability of said polypeptide to hydroxylate a prolyl or other residue of
HIF polypeptide as described herein. Crystallographic information may also be used
to identify other HIF hydroxylases, for use in subsequent assays of HIF hydroxylase
5 activity.

In an alternative aspect of the present invention, a HIF hydroxylase of the
invention may be a variant which does not show the same activity as the HIF
hydroxylase, but is one which inhibits a function of the wild type polypeptide. For
example, a modified or variant HIF hydroxylase may be one which competes for HIF
10 hydroxylase substrates but which does not lead to prolyl hydroxylation of such
substrate. Such variants may be used in the various embodiments of the invention.

Amino Acid Sequence Identity

A polypeptide may comprise an amino acid sequence which shares greater
than about 60% sequence identity with a polypeptide sequence described or
15 referenced herein, greater than about 70%, greater than about 80% greater than about
90%, greater than about 95%, or greater than about 98%.

For amino acid "homology", this may be understood to be identity e.g. as
determined using the algorithm GAP (as described below).

Amino acid identity is generally defined with reference to the algorithm GAP
20 (Genetics Computer Group, Madison, WI). GAP uses the Needleman and Wunsch
algorithm to align two complete sequences that maximizes the number of matches
and minimizes the number of gaps. Generally, the default parameters are used, with
a gap creation penalty = 12 and gap extension penalty = 4. Use of GAP may be
preferred but other algorithms may be used, e.g. BLAST, (which uses the method of
25 Altschul *et al.* (1990) *J. Mol. Biol.* 215: 405-410), gapped BLAST, PSI-BLAST,
(Altschul S. (1997) *Nucleic Acid Res.* 17 3389-33402), FASTA (which uses the
method of Pearson and Lipman (1988) *PNAS USA* 85: 2444-2448), or the Smith-
Waterman algorithm (Smith and Waterman (1981) *J. Mol Biol.* 147: 195-197).
Generally, the default parameters are used, with a gap creation penalty = 12 and gap
30 extension penalty = 4.

Sequence comparison may be made over the full-length of the relevant sequence shown herein, or may more preferably be over a contiguous sequence of about or greater than about 20, 25, 30, 33, 40, 50, 67, 133, 167, 200, 233, 267, 300, 333, 400 or more amino acids, compared with the relevant amino acid sequence.

5 Where default parameters or other features of these programs are subject to revision, it is to be understood that reference to the programs and their parameters are as of the priority date of the instant application.

Substitutions made to polypeptides of the invention may include conserved substitutions, for example according to the following table, where amino acids on the
10 same block in the second column and preferably in the same line in the third column may be substituted for each other:

15	ALIPHATIC	Non-polar	G A P
			I L V
		Polar - uncharged	C S T M
			N Q
		Polar - charged	D E
			K R H
20	AROMATIC		F W Y

Alternatively, any amino acid may be replaced by a small aliphatic amino acid,
25 preferably glycine or alanine.

In addition, deletions and insertions (e.g. from 1 to 5 subject to a maximum of 40% of the amino acids) may also be made. Insertions are preferably insertions of small aliphatic amino acids, such as glycine or alanine, although other insertions are not excluded.

Variant polypeptides may also be modified in any of the ways described herein for polypeptides of the invention. This includes for example "reverse" C-terminal to N-terminal sequences, synthetic amino acids, glycosylated peptides, phosphorylated peptides, addition of metal ions such as ions of calcium, zinc, iron or manganese modified side chains and labelling. Polypeptides may be provided in the form of molecules which contain multiple copies of the polypeptide (or mixtures of polypeptides). For example, the amino group of the side chain of lysine may be used as an attachment point for the carboxy terminus of an amino acid. Thus two amino acids may be joined to lysine via carbonyl linkages, leading to a branched structure which may in turn be branched one or more times. By way of example, four copies of a polypeptide of the invention may be joined to such a multiple antigen peptide (MAP), such as a MAP of the structure $\text{Pep}_4\text{-Lys}_2\text{-Lys-X}$, where Pep is a polypeptide from the HIF hydroxylase or variant thereof (optionally in the form of a heterologous fusion), Lys is lysine and X is a terminal group such as β -alanine which provides for joining of the MAP core to a solid support such as a resin for synthesis of the $\text{Pep}_4\text{-MAP}$ peptide and which may be removed from the support once synthesis is complete.

Other multiple polypeptide structures may be obtained using the MAP cores described in: Lu *et al.*, 1991, *Mol Immunol*, 28, 623-30; Briand *et al.*, 1992, *J Immunol Methods*, 156, 255-65; Ahlborg, 1995, *J Immunol Methods*, 179, 269-75.

Where multimers of the invention are provided, they may comprise different polypeptides of the invention or be multimers of the same polypeptide.

Except where specified to the contrary, the polypeptide sequences described herein are shown in the conventional 1-letter code and in the N-terminal to C-terminal orientation. The amino acid sequence of polypeptides of the invention may also be further modified to include non-naturally-occurring amino acids or to increase the stability of the compound *in vivo*. When the compounds are produced by synthetic means, such amino acids may be introduced during production. The compound may also be modified following either synthetic or recombinant production.

Polypeptides of the invention may also be made synthetically using D-amino acids. In such cases, the amino acids may be linked in a reverse sequence in the C to N orientation. β -amino acids (or higher homologues) may also be used.

5 A number of side-chain modifications for amino acids are known in the art and may be made to the side chains of polypeptides of the present invention. Such modifications include for example, modifications of amino groups by reductive alkylation by reaction with an aldehyde followed by reduction with NaBH_4 , amidination with methylacetimidate or acylation with acetic anhydride.

10 The guanidino groups of arginine residues may be modified by the formation of heterocyclic condensation products with reagents such as 2,3-butanedione or glyoxal. Sulphydryl groups may be modified by methods such as carboxymethylation, tryptophan residues may be modified by oxidation or alkylation of the indole ring and the imidazole ring of histidine residues may be modified by alkylation.

15 The carboxy terminus and any other carboxy side chains may be blocked in the form of an ester group, e.g. a C_{1-6} alkyl ester.

The above examples of modifications to amino acids are not exhaustive. Those of skill in the art may modify amino acid side chains where desired using chemistry known per se in the art.

20 Polypeptides may be made synthetically or recombinantly, using techniques which are widely available in the art. Synthetic production generally involves step-wise addition of individual amino acid residues to a reaction vessel in which a polypeptide of a desired sequence is being made.

Polynucleotides

25 The invention also includes nucleotide sequences that encode for a HIF hydroxylase or a variant or fragment thereof as well as nucleotide sequences which are complementary thereto. In particular, the invention provides nucleotide sequences which encode a human HIF hydroxylase or a fragment or variant of a human HIF hydroxylase as well as nucleotide sequences complementary to any of
30 these sequences: The nucleotide sequence may be RNA or DNA including genomic

DNA, synthetic DNA or cDNA. Preferably the nucleotide sequence is a DNA sequence and most preferably, a cDNA sequence. The invention also encompasses PNA (protein nucleic acid) molecules comprising the sequences of the invention.

Nucleotide sequence information for human PHD 1, 2 and 3 is provided in SEQ ID

5 NOs: 1, 3 and 5 respectively. Nucleotide sequence information is provided in SEQ ID NO: 7, for the EGL-9 polypeptide of *C. elegans*. Such nucleotides can be isolated from cells or synthesised according to methods well known in the art, as described by way of example in Sambrook *et al*, 1989.

Typically a polynucleotide of the invention comprises a contiguous sequence
10 of nucleotides which is capable of hybridizing under selective conditions to the coding sequence or the complement of the coding sequence of SEQ ID NO: 1, 3, 5 or 7 and in particular to the coding sequence or the complement of SEQ ID NO: 1, 3 or 5.

A polynucleotide of the invention can hybridize to the coding sequence or the
15 complement of the coding sequence of SEQ ID NO: 1, 3, 5 or 7, and in particular to the coding sequence or the complement of SEQ ID NO: 1, 3 or 5, at a level significantly above background. Background hybridization may occur, for example, because of other cDNAs present in a cDNA library. The signal level generated by the interaction between a polynucleotide of the invention and the coding sequence or
20 complement of the coding sequence of SEQ ID NO: 1, 3, 5 or 7 is typically at least 10 fold, preferably at least 100 fold, as intense as interactions between other polynucleotides and the coding sequence of SEQ ID NO: 1, 3, 5 or 7. The intensity of interaction may be measured, for example, by radiolabelling the probe, e.g. with ³²P. Selective hybridisation may typically be achieved using conditions of medium
25 to high stringency. However, such hybridisation may be carried out under any suitable conditions known in the art (see Sambrook *et al*, 1989. For example, if high stringency is required suitable conditions include from 0.1 to 0.2 x SSC at 60°C up to 65°C. If lower stringency is required suitable conditions include 2 x SSC at 60°C.

The coding sequence of SEQ ID NO: 1, 3, 5 or 7 may be modified by
30 nucleotide substitutions, for example from 1, 2 or 3 to 10, 25, 50 or 100 substitutions. The polynucleotide of SEQ ID NO: 1, 3, 5 or 7 may alternatively or additionally be

modified by one or more insertions and/or deletions and/or by an extension at either or both ends. A polynucleotide may include one or more introns, for example may comprise genomic DNA. The modified polynucleotide generally encodes a polypeptide which has HIF hydroxylase activity, typically which has hydroxylase activity and in particular prolyl hydroxylase activity. Alternatively, a polynucleotide encodes a ligand-binding portion of a polypeptide or a polypeptide which modulates HIF hydroxylase activity. Degenerate substitutions may be made and/or substitutions may be made which would result in a conservative amino acid substitution when the modified sequence is translated, for example as shown in the Table above.

A nucleotide sequence which is capable of selectively hybridizing to the complement of the DNA coding sequence of SEQ ID NO: 1, 3, 5 or 7 will generally have at least 60%, at least 70%, at least 80%, at least 88%, at least 90%, at least 95%, at least 98% or at least 99% sequence identity to the coding sequence of SEQ ID NO: 1, 3, 5 or 7 over a region of at least 20, preferably at least 30, for instance at least 40, at least 60, more preferably at least 100 contiguous nucleotides or most preferably over the full length of SEQ ID NO: 1, 3, 5 or 7. Preferably the nucleotide sequence encodes a polypeptide which has the same domain structure as a HIF hydroxylase as described in more detail above.

For example the UWGCG Package provides the BESTFIT program which can be used to calculate homology (for example used on its default settings) (Devereux *et al* (1984) *Nucleic Acids Research* 12, p387-395). The PILEUP and BLAST algorithms can be used to calculate homology or line up sequences (typically on their default settings), for example as described in Altschul (1993) *J. Mol. Evol.* 36:290-300; Altschul *et al* (1990) *J. Mol. Biol.* 215:403-10.

Software for performing BLAST analyses is publicly available through the National Centre for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pair (HSPs) by identifying short words of length W in the query sequence that either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighbourhood word score threshold (Altschul *et al*, 1990). These initial neighbourhood word hits act as seeds for

initiating searches to find HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Extensions for the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLAST program uses as defaults a word length (W) of 11, the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1992) *Proc. Natl. Acad. Sci. USA* 89: 10915-10919) alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.

The BLAST algorithm performs a statistical analysis of the similarity between two sequences; see e.g., Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90: 5873-5787 and Altschul and Gish (1996) *Methods Enzymol.* 266: 460-480. One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a sequence is considered similar to another sequence if the smallest sum probability in comparison of the first sequence to the second sequence is less than about 1, preferably less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

Any combination of the above mentioned degrees of sequence identity and minimum sizes may be used to define polynucleotides of the invention, with the more stringent combinations (i.e. higher sequence identity over longer lengths) being preferred. Thus, for example a polynucleotide which has at least 90% sequence identity over 25, preferably over 30 nucleotides forms one aspect of the invention, as does a polynucleotide which has at least 95% sequence identity over 40 nucleotides.

The nucleotides of the invention may comprise a label for example, they may be radiolabelled or fluorescently labelled. The label may be such that it is only visualised on hybridisation to a complementary nucleic acid. For example, the label

may be quenched until hybridisation. The nucleotides of the invention may be immobilised to a support such as a membrane or as a microarray.

The nucleotides according to the invention have utility in production of the proteins according to the invention, which may take place *in vitro*, *in vivo* or *ex vivo*.

5 Accordingly, the invention provides a polypeptide encoded by a polynucleotide of the invention and in particular encoded by SEQ ID NO: 1, 3, 5 or 7. The invention includes a PHD polypeptide encoded by PHD gene, in particular by the PHD 1, 2 or 3 genes. The invention also provides fragments of such polypeptides which have, HIF hydroxylase activity and in particular prolyl hydroxylase activity.

10 The nucleotides may be involved in recombinant protein synthesis or indeed as therapeutic agents in their own right, utilised in gene therapy techniques. Nucleotides complementary to those encoding HIF hydroxylase, or antisense sequences, or interfering RNA may also be used in gene therapy.

The present invention also includes expression vectors that comprise
15 nucleotide sequences encoding the proteins of the invention. Such expression vectors are routinely constructed in the art of molecular biology and may for example involve the use of plasmid DNA and appropriate initiators, promoters, enhancers and other elements, such as for example polyadenylation signals which may be necessary, and which are positioned in the correct orientation, in order to allow for protein
20 expression. Other suitable vectors would be apparent to persons skilled in the art. By way of further example in this regard we refer to Sambrook *et al.* 1989.

Polynucleotides according to the invention may also be inserted into the vectors described above in an antisense orientation in order to provide for the production of antisense RNA. Antisense RNA or other antisense polynucleotides
25 may also be produced by synthetic means. Such antisense polynucleotides may be used as test compounds in the assays of the invention or may be useful in a method of treatment of the human or animal body by therapy.

Polynucleotides of the invention may also be used to design double stranded RNAs for use in RNA interference. Such RNA comprises short stretches of double
30 stranded RNA having the same sequence as a target mRNA. Such sequences can be used to inhibit translation of the mRNA. Alternatively, small fragments of the gene

encoding a HIF hydroxylase may be provided, cloned back to back in a plasmid. Expression leads to production of the desired double stranded RNA. Such short interfering RNA (siRNA) may be used for example to reduce or inhibit expression of a HIF hydroxylase of the invention, in assays or in a method of therapy. The invention also relates to such siRNAs. Such siRNAs may be designed to inhibit groups of HIF hydroxylases of the invention by basing their sequences on regions of conserved sequence in the encoding genes of the hydroxylases. Alternatively, the siRNAs may be made specific to a particular HIF hydroxylase by choosing a sequence unique to the encoding gene of the particular hydroxylase gene to be inhibited.

Preferably, a polynucleotide of the invention in a vector is operably linked to a control sequence which is capable of providing for the expression of the coding sequence by the host cell, i.e. the vector is an expression vector. The term "operably linked" refers to a juxtaposition wherein the components described are in a relationship permitting them to function in their intended manner. A regulatory sequence, such as a promoter, "operably linked" to a coding sequence is positioned in such a way that expression of the coding sequence is achieved under conditions compatible with the regulatory sequence.

The vectors may be for example, plasmid, virus or phage vectors provided with a origin of replication, optionally a promoter for the expression of the said polynucleotide and optionally a regulator of the promoter. The vector may be an artificial chromosome. The vectors may contain one or more selectable marker genes, for example an ampicillin resistance gene in the case of a bacterial plasmid or a resistance gene for a fungal vector. Vectors may be used *in vitro*, for example for the production of DNA or RNA or used to transfect or transform a host cell, for example, a mammalian host cell. The vectors may also be adapted to be used *in vivo*, for example in a method of gene therapy.

Promoters and other expression regulation signals may be selected to be compatible with the host cell for which expression is designed. For example, yeast promoters include *S. cerevisiae* GAL4 and ADH promoters, *S. pombe* *nmt1* and *adh* promoter. Mammalian promoters include the metallothionein promoter which can be

induced in response to heavy metals such as cadmium. Viral promoters such as the SV40 large T antigen promoter or adenovirus promoters may also be used. An IRES promoter may also be used. All these promoters are readily available in the art.

Mammalian promoters, such as β -actin promoters, may be used. Tissue-specific promoters are especially preferred. Inducible promoters are also preferred. Promoters inducible by hypoxic conditions may, for example, be employed. Viral promoters may also be used, for example the Moloney murine leukaemia virus long terminal repeat (MMLV LTR), the rous sarcoma virus (RSV) LTR promoter, the, SV40 promoter, the human cytomegalovirus (CMV) IE promoter, adenovirus, HSV promoters (such as the HSV IE promoters), or HPV promoters, particularly the HPV upstream regulatory region (URR). Viral promoters are readily available in the art.

The vector may further include sequences flanking the polynucleotide giving rise to polynucleotides which comprise sequences homologous to eukaryotic genomic sequences, preferably mammalian genomic sequences, or viral genomic sequences. This will allow the introduction of the polynucleotides of the invention into the genome of eukaryotic cells or viruses by homologous recombination. In particular, a plasmid vector comprising the expression cassette flanked by viral sequences can be used to prepare a viral vector suitable for delivering the polynucleotides of the invention to a mammalian cell. Homologous recombination may also be used to disrupt or mutate endogenous sequences in cells encoding HIF hydroxylases. Other examples of suitable viral vectors include herpes simplex viral vectors and retroviruses, including lentiviruses, adenoviruses, adeno-associated viruses and HPV viruses. Gene transfer techniques using these viruses are known to those skilled in the art. Retrovirus vectors for example may be used to stably integrate the polynucleotide giving rise to the polynucleotide into the host genome. Replication-defective adenovirus vectors by contrast remain episomal and therefore allow transient expression.

The invention also includes cells that have been modified to express a HIF hydroxylase of the invention. Such cells include transient, or preferably stable higher eukaryotic cell lines, such as mammalian cells or insect cells, using for example a

baculovirus expression system, lower eukaryotic cells, such as yeast or prokaryotic cells such as bacterial cells. Particular examples of cells which may be modified by insertion of vectors encoding for a polypeptide according to the invention include mammalian thymic epithelial cells, fibroblasts, HEK293T, U20S, CHO, HeLa, BHK, 3T3 and COS cells.

A polypeptide of the invention may be expressed in cells of a transgenic non-human animal, typically a mammal, preferably a rodent, more preferably a mouse. The animal may be a larger animal such as a pig or sheep. A transgenic non-human animal expressing a polypeptide of the invention is included within the scope of the invention. Also included are transgenic animals expressing an antisense RNA, siRNA or ribozyme designed to inhibit expressions of a polypeptide of the invention. The transgenic animals of the invention may have a gene encoding an endogenous HIF hydroxylase disrupted or mutated. For example, the endogenous HIF hydroxylase may be rendered inactive and lack hydroxylase activity.

Antibodies

According to another aspect, the present invention also relates to antibodies, specific for a polypeptide of the invention. Such antibodies are for example useful in purification, isolation or screening methods involving immunoprecipitation techniques or, indeed, as therapeutic agents in their own right. Antibodies may be raised against specific epitopes of the polypeptides according to the invention.

Antibodies may be used to impair HIF hydroxylase function. An antibody, or other compound, "specifically binds" to a protein when it binds with preferential or high affinity to the protein for which it is specific but does not substantially bind or binds with only low affinity to other proteins. A variety of protocols for competitive binding or immunoradiometric assays to determine the specific binding capability of an antibody are well known in the art (see for example Maddox *et al*, J. Exp. Med. 158, 1211-1226, 1993). Such immunoassays typically involve the formation of complexes between the specific protein and its antibody and the measurement of complex formation.

Antibodies of the invention may be antibodies to human polypeptides or fragments thereof. For the purposes of this invention, the term "antibody", unless specified to the contrary, includes fragments which bind a polypeptide of the invention. Such fragments include Fv, F(ab') and F(ab')₂ fragments, as well as single chain antibodies. Furthermore, the antibodies and fragment thereof may be chimeric antibodies, CDR-grafted antibodies or humanised antibodies.

Antibodies may be used in a method for detecting polypeptides of the invention in a biological sample, which method comprises:

- I providing an antibody of the invention;
- 10 II incubating a biological sample with said antibody under conditions which allow for the formation of an antibody-antigen complex; and
- III determining whether antibody-antigen complex comprising said antibody is formed.

A sample may be for example a tissue extract, blood, serum and saliva.

15 Antibodies of the invention may be bound to a solid support and/or packaged into kits in a suitable container along with suitable reagents, controls, instructions, etc. Antibodies may be linked to a revealing label and thus may be suitable for use in methods of *in vivo* HIF hydroxylase imaging.

Antibodies of the invention can be produced by any suitable method. Means for preparing and characterising antibodies are well known in the art, see for example Harlow and Lane (1988) "Antibodies: A Laboratory Manual", Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. For example, an antibody may be produced by raising antibody in a host animal against the whole polypeptide or a fragment thereof, for example an antigenic epitope thereof, herein after the

25 "immunogen".

A method for producing a polyclonal antibody comprises immunising a suitable host animal, for example an experimental animal, with the immunogen and isolating immunoglobulins from the animal's serum. The animal may therefore be inoculated with the immunogen, blood subsequently removed from the animal and

30 the IgG fraction purified.

A method for producing a monoclonal antibody comprises immortalising cells which produce the desired antibody. Hybridoma cells may be produced by fusing spleen cells from an inoculated experimental animal with tumour cells (Kohler and Milstein (1975) *Nature* 256, 495-497).

- 5 An immortalized cell producing the desired antibody may be selected by a conventional procedure. The hybridomas may be grown in culture or injected intraperitoneally for formation of ascites fluid or into the blood stream of an allogenic host or immunocompromised host. Human antibody may be prepared by *in vitro* immunisation of human lymphocytes, followed by transformation of the
- 10 lymphocytes with Epstein-Barr virus and in transgenic mice enabling production of human antibodies.

For the production of both monoclonal and polyclonal antibodies, the experimental animal is suitably a goat, rabbit, rat or mouse. If desired, the immunogen may be administered as a conjugate in which the immunogen is coupled,

15 for example via a side chain of one of the amino acid residues, to a suitable carrier. The carrier molecule is typically a physiologically acceptable carrier. The immunogen may, for example, be administered with an adjuvant. The antibody obtained may be isolated and, if desired, purified.

Assays

- 20 Our data shows that hydroxylation of HIF- α is mediated by a hydroxylase enzyme which has specificity or selectivity for HIF- α . The enzyme responsible are referred to as HIF hydroxylases and include EGL-9 and PHD1, 2 and 3. The action of HIF hydroxylases, and in particular human HIF hydroxylases, represent a novel target for the control of HIF α . By blocking HIF hydroxylase activity, this will
- 25 reduce hydroxylation of HIF- α and thus lead to the accumulation of HIF- α in cells. This in turn will lead to the activation of systemic local defences against hypoxia or ischaemia that may include the promotion of angiogenesis, erythropoiesis, energy metabolism, inflammation, vasomotor function, and will also affect apoptotic/proliferative responses.

- 30 We describe below in more detail a number of different assays that may be carried out to identify modulators of HIF hydroxylase activity and in particular of

prolyl hydroxylase activity, or which affect regulation of HIF- α levels in a cell and hence which affect HIF mediated activity. Some of these assays utilise HIF polypeptides and VHL polypeptides, and in particular HIF hydroxylases in accordance with the present invention. Typically, the assays may utilise a human
5 HIF hydroxylase such as PHD 1, 2 or 3 or a fragment or variant of a human HIF hydroxylase. In a preferred embodiment an enzyme with HIF prolyl-hydroxylase activity may be used. These components are described in more detail below. Each of these components, where required may be provided either in purified or unpurified form, for example, as cellular extracts or by purification of the relevant component
10 from such extracts. Alternatively, the relevant component can be expressed using recombinant expression techniques and purified for use in the assay. Alternatively, the components may be expressed recombinantly in a cell for use in cell based assays.

Typically, a polynucleotide encoding the relevant component is provided
15 within an expression vector. Such expression vectors are routinely constructed in the art and may for example involve the use of plasmid DNA and appropriate initiators, promoters, enhancers and other elements, such as for example polyadenylation signals which may be necessary and which are positioned in the correct orientation in order to allow full protein expression. Suitable vectors would be very readily
20 apparent to those of skill in the art, such as those described in more detail herein with reference to the HIF hydroxylases. Promoter sequences may be inducible or constitutive promoters depending on the selected assay format. The promoter may be tissue specific. Examples of promoters and other flanking sequences for use in the expression vectors are described in more detail herein with reference to the HIF
25 hydroxylases of the invention and in particular to the human HIF hydroxylases of the invention.

HIF Polypeptides and Peptide Analogues

The assays of the present invention may use a substrate of a HIF hydroxylase
30 and in particular a prolyl containing substrate of the enzyme. In particular, such

substrates may be used in assays to monitor for the activity of a modulator of HIF hydroxylase activity. The substrate may be a HIF polypeptides or peptide analogue thereof. Typically, a HIF polypeptide will be used as the substrate.

Any suitable substrate in which a residue, preferably a proline residue, is hydroxylated by a HIF hydroxylase of SEQ ID NO: 2, 4, 6 or 8 may be used and in particular one which is hydroxylated by the HIF hydroxylase of SEQ ID NO: 2, 4 or 6. In preferred embodiments of the invention, such a substrate is a HIF polypeptide such as a HIF-1 α or HIF-2 α subunit protein or fragment of either or peptide analogue of the subunit or fragment. Preferably, the HIF- α peptide conveys an oxygen regulated response. More preferably, the HIF- α peptide is capable of oxygen regulated binding to pVHL. Preferably, such HIF polypeptides, fragments or peptide analogues incorporate a proline residue equivalent to Pro 564 and/or Pro 402 as defined with reference to HIF-1 α . The proline equivalent to Pro 564 and/or Pro 402 of HIF-1 α may be determined by aligning the HIF variant, fragment or analogue to the sequence of HIF-1 α to obtain the best sequence alignment and identifying thereby the proline equivalent to Pro 564 and/or Pro 402 of HIF-1 α . In the assays of the invention the hydroxylation of one or both of these prolines may be determined.

A HIF polypeptide may be of eukaryotic origin, in particular a human or other mammalian, HIF- α subunit protein or fragment thereof. Alternatively, the polypeptide may be of *C. elegans* origin. In those assays which monitor for hydroxylation of HIF- α through its interaction with and subsequent destruction by VHL, the HIF polypeptide has the ability to bind to a wild type full length pVHL protein, such that the binding is able, in a normoxic cellular environment, to target the HIF- α subunit for destruction i.e. the polypeptide comprises a pVHL binding domain.

A number of HIF α subunit proteins have been cloned. These include HIF-1 α , the sequence of which is available as Genbank accession number U22431, HIF-2 α , available as Genbank accession number U81984 and HIF-3 α , available as Genbank accession numbers AC007193 and AC079154. These are all human HIF α subunit proteins and all may be used in the invention. HIF- α subunit proteins from other

species, including murine HIF-1 α (accession numbers AF003695, U59496 and X95580), rat HIF-1 α (accession number Y09507), murine HIF-2 α (accession numbers U81983 and D89787) and murine HIF-3 α (accession number AF060194) may also be used in the invention. Other mammalian, vertebrate, invertebrate or
5 other homologues may be obtained by techniques similar to those described above for obtaining pVHL homologues.

One HIF- α protein of particular interest is the *C.elegans* HIF- α subunit protein. The HIF- α /VHL system of regulation is conserved in *C.elegans*, so that the *C.elegans* system may be used in assays of the present invention.

10 There are a number of common structural features found in the two HIF- α subunit proteins identified to date. Some of these features are identified in O'Rourke *et al* (1999, J. Biol. Chem., 274; 2060-2071) and may be involved in the trans-activation functions of the HIF- α subunit proteins. One or more of these common structural features are preferred features of the HIF polypeptides.

15 Fragments of HIF-1 α or peptide analogues preferably include proline residue 402 and/or 564 (U22431), which are hydroxylated by HIF prolyl hydroxylases. Suitable fragments may include or consist of residues 344-698, particularly residues 364-678, more particularly residues 364-638 or 384-638 and still more particularly residues 364 - 598 or 394-598. Other suitable fragments may include or consist of
20 residues 549-652 and even more particularly the N-terminal region thereof which interacts with the VHL protein. C-terminal fragments may include residues 549 to 582 and in particular residues 556-574. Other suitable fragments comprise or consist of residues 344-417, more preferably 380-417. Such a region, or its equivalent in other HIF- α subunit proteins, is desirably present in the HIF α polypeptides described
25 herein. The substrates used in the assays of the invention may typically comprise residues 549 to 582 of the human HIF-1 α sequence.

Variants of the above HIF- α subunits may be used, such as synthetic variants which have at least 45% amino acid identity to a naturally occurring HIF- α subunit (particularly to a human HIF- α subunit such as, for example HIF-1 α), preferably at
30 least 50%, 60%, 70%, 80%, 90%, 95% or 98% identity. Such variants may include substitutions or modifications as described above with respect to HIF hydroxylases.

Amino acid activity may also be calculated as described above with reference to HIF hydroxylases.

HIF fragments may also include non-peptidyl functionalities and may be optimised for assay purposes such that the level of identity is lowered. Such
5 functionalities may be covalently bound such as sugars or non-covalently bound such as metal ions.

HIF α polypeptides as described herein may be fragments of the HIF- α subunit protein or variants as described above, provided that said fragments retain the ability to interact with a wild-type pVHL, preferably wild-type human pVHL. When using
10 proteinogenic amino acid residues, such fragments are desirably at least 20, preferably at least 40, 50, 75, 100, 200, 250 or 400 amino acids in size. Desirably, such fragments include proline residue 402 and/or 564. Some preferred fragments include the region 556-574 found in human HIF-1 α or its equivalent regions in other HIF- α subunit proteins, e.g. 517-542 of HIF-2 α . Optionally, the fragments also
15 include one or more domains of the protein responsible for trans-activation. Reference herein to a HIF- α polypeptide or HIF- α subunit protein includes the above mentioned mutants and fragments or other HIF- α fragments which are functionally able to bind VHL protein unless the context is explicitly to the contrary.

Cell based assays of the present invention may involve upregulation of an
20 endogenous HIF- α or expression of a HIF- α by recombinant techniques and in particular of HIF-1 α .

VHL

Some assays in accordance with the present invention utilise VHL and in particular monitor the interaction between hydroxylated HIF and VHL and the
25 subsequent destruction of HIF- α . The VHL may be any suitable mammalian VHL, particularly human VHL. It may be a *C. elegans* VHL. Human VHL has been cloned and sources of the gene can be readily identified by those of skill in the art. Its sequence is available as Genbank accession numbers AF010238 and L15409. Other mammalian VHLs are also available, such as murine VHL (accession number
30 U12570) and rat (accession numbers U14746 and S80345). Non-mammalian

homologues include the VHL-like protein of *C.elegans*, accession number F08G12.4. VHL gene sequences may also be obtained by routine cloning techniques, for example by using all or part of the human VHL gene sequence as a probe to recover and to determine the sequence of the VHL gene in other species. A wide variety of techniques are available for this, for example PCR amplification and cloning of the gene using a suitable source of mRNA (e.g. from an embryo or a liver cell), obtaining a cDNA library from a mammalian, vertebrate, invertebrate or other source, e.g a cDNA library from one of the above-mentioned sources, probing said library with a polynucleotide of the invention under stringent conditions, and recovering a cDNA encoding all or part of the VHL protein of that mammal. Suitable stringent conditions include hybridization on a solid support (filter) overnight incubation at 42°C in a solution containing 50% formamide, 5xSSC (750 mM NaCl, 75 mM sodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulphate and 20 µg/ml salmon sperm DNA, followed by washing in 0.2x SSC at from about 50°C to about 60°C). Where a partial cDNA is obtained, the full length coding sequence may be determined by primer extension techniques.

A further approach is to use the above-identified sequences as query sequences to search databases for homologous gene sequences or partial gene sequences (particularly ESTs). Matches identified may be examined and where an actual or putative VHL sequence is found, the gene recovered by physical cloning using, for example PCR and RACE-PCR based on the sequence of the match.

Although wild-type VHL is preferred, mutants and variants of VHL which still retain the ability to interact directly with the HIF- α subunit may also be used. Examples of VHL mutants are well known in the art and include mutants described by Stebbins *et al* (Science, 1999, 284; 55-61) which have changes to the Elongin C interacting interface.

Mutants and other variants will generally be based on wild-type mammalian VHLs and have a degree of amino acid identity which is desirably at least 70%, preferably at least 80%, 90%, 95% or even 98% homologous to a wild type mammalian VHL, preferably to human VHL.

It is not necessary to use the entire VHL proteins (including their mutants and other variants) for assays of the invention. Fragments of the VHL may be used provided such fragments retain the ability to interact with the target domain of the HIF α subunit. Optionally, the fragment may include the Elongin C interacting interface domain. Fragments of VHL may be generated in any suitable way known to those of skill in the art. Suitable ways include, but are not limited to, recombinant expression of a fragment of the DNA encoding the VHL. Such fragments may be generated by taking DNA encoding the VHL, identifying suitable restriction enzyme recognition sites either side of the portion to be expressed, and cutting out said portion from the DNA. The portion may then be operably linked to a suitable promoter in a standard commercially available expression system. Another recombinant approach is to amplify the relevant portion of the DNA with suitable PCR primers. Small fragments of the VHL (up to about 20 or 30 amino acids) may also be generated using peptide synthesis methods which are well known in the art. Generally fragments will be at least 40, preferably at least 50, 60, 70, 80 or 100 amino acids in size.

Particularly preferred fragments include those which are based upon the beta domain located within the fragment 63-156 of the 213 amino acid human VHL protein, or the equivalent domain in other variants. In a preferred embodiment, such domains will have at least 70%, preferably 80%, 90%, 95% or even 98% degree of sequence identity to the 64-156 fragment of human VHL. Fragments of this region and its variants may be used. These fragments may be 15-80 amino acids in length, for example from 20 to 80, such as 30-60 amino acids in length. Fragments may include the regions 71-90 or 90-109 of human VHL or their equivalents in the above described variants. Desirably, the wild-type sequence of the beta domain is retained.

One fragment which may be used is that in which up to 53 of the N-terminal residues, e.g. from 1 to n wherein n is an integer of from 2 to 53, have been deleted, the rest of the protein being wild-type.

The ability of suitable fragments to bind to the HIF α subunit (or fragment thereof) may be tested using routine procedures such as those described in the accompanying Examples relating to intact VHL. Reference herein to a VHL protein

includes the above mentioned mutants and fragments which are functionally able to bind the HIF α subunit unless the context is explicitly to the contrary.

Hydroxylases

In a number of the assays in accordance with the present invention,
5 hydroxylase enzyme is provided. In preferred embodiments, the hydroxylase enzyme is a HIF hydroxylase in accordance with the present invention. The enzyme is preferably a prolyl hydroxylase. In a particularly preferred embodiment of the invention the HIF hydroxylase used comprises:

- (a) the amino acid sequence of SEQ ID NO: 2, 4, 6 or 8;
- 10 (b) a variant thereof having at least 60% identity to the amino acid sequence of SEQ ID NO: 2, 4, 6 or 8 and having HIF hydroxylase activity; or
- (c) a fragment of either thereof having HIF hydroxylase activity.

Such hydroxylase enzymes, and in particular prolyl-hydroxylases such as for example 4-prolyl hydroxylase, are obtainable from extracts of mammalian cells,
15 including immortalised mammalian cells in culture such as HeLa, RCC or CHO-K1 cells, primary cells, tissues or primary cell lysates (e.g. rabbit reticulocyte or human placental lysates). Cell extracts may be prepared in accordance with standard techniques available in the art by reference to the accompanying examples. Assays may alternatively be carried out as cell based assays in which hydroxylase enzyme is
20 expressed endogenously.

In a preferred embodiment of any one of the assays in accordance with the present invention, the assay utilises a HIF hydroxylase, typically a human HIF hydroxylase, and in particular a PHD hydroxylase of the present invention. Such hydroxylases may be upregulated before or during the course of the assay.
25 Alternatively, the enzyme may be expressed recombinantly, and the HIF hydroxylase of the invention isolated from such recombinant expression systems in purified or unpurified form for use in the assays. Alternatively, cells may be provided which have been transformed or transfected with expression vectors expressing a HIF hydroxylase in accordance with the present invention. Such methods provide assays
30 for substances that inhibit, promote or otherwise modulate the individual activities of HIF hydroxylases in either a specific or a general manner. The methods may also be

used to identify substances that inhibit, promote or otherwise modulate the activities of a group of HIF hydroxylases such as, for example, all of PHD 1, 2 and 3 or any two of the three enzymes.

The assays of the invention may use an EGLN polypeptide such as EGLN1
5 (gil457146), EGLN2 (gil457148), EGLN3 (gil4547150), FLJ21620 (BAB15101) or Clorf12 (NP071334).

In general, the HIF hydroxylases of the invention are iron dependent, that is they typically require ferrous (FeII) ions for activity. Accordingly, the assays of the invention will typically include ferrous compounds, unless the purpose of the assay is
10 to determine the effect of the absence of ferrous ions or it is desired to carry out control reactions where no ferrous ions are present.

Assay methods

The present invention provides an assay method for identifying an agent which modulates the interaction of a hypoxia inducible factor (HIF) hydroxylase with
15 a substrate of the hydroxylase, the method comprising:

- contacting a HIF hydroxylase and a test substance in the presence of a substrate of the hydroxylase under conditions in which the hydroxylase interacts with the substrate in the absence of the test substance; and
- determining the interaction, or lack of interaction, of the hydroxylase
20 and the substrate. The interaction of the hydroxylase with the substrate may be determined by measuring the hydroxylase activity of the hydroxylase.

The interaction between hydroxylase and substrate refers to physical interaction or to functional interaction. The interaction may therefore be measured by any suitable method, including binding of the hydroxylase to a substrate, the activity
25 of the hydroxylase on the substrate, or any activity related to the action of the hydroxylase on the substrate, such as the levels of co-factors or by-products used or produced in the hydroxylation reaction, or downstream effects mediated through hydroxylation of the substrate.

In another aspect of the present invention, there is provided an assay for an
30 inhibitor of HIF- α destruction or HIF- α transcription inactivation comprising

providing HIF- α or a peptide analogue thereof, incubating HIF- α or the peptide analogue with a test substance under conditions which allow for hydroxylation of HIF- α in the absence of the test substance, and monitoring for hydroxylation of HIF- α . Preferably, HIF- α or the peptide analogue thereof includes a prolyl residue such as Pro 564 and/or Pro 402 of HIF- α or an equivalent prolyl in a peptide analogue, and said assay is carried out under conditions which allow for hydroxylation of Pro 564 and/or Pro 402 in the absence of the test substance, and monitoring for hydroxylation of Pro 564 and/or Pro 402.

In a further aspect, there is provided an assay for an inhibitor of VHL-mediated HIF- α destruction, which comprises

- providing a HIF- α , or fragment thereof which includes a VHL-binding portion, together with its cognate prolyl-hydroxylase under conditions suitable for the hydroxylation of a proline residue in the HIF- α VHL-binding domain;
- providing a putative modulator of hydroxylation; and
- determining whether the amount of hydroxylation of said proline residue has been modulated by said putative modulator. In one embodiment of the invention the cognate prolyl-hydroxylase is a prolyl-4-hydroxylase.

Conversely, in hypoxic conditions such as those commonly found in tumours, the lack of hydroxylation of HIF, and in particular of proline hydroxylation, may lead to the accumulation of HIF α and the concomitant promotion of angiogenesis and other growth promoting events. Alternatively, in ischaemic/hypoxic conditions in which normal levels of HIF activity are present, it may also be desirable to increase existing HIF activity. Thus mechanisms which either upregulate a HIF hydroxylase, increase the activity of a HIF hydroxylase, rescue or bypass the hydroxylase are a target in such cells.

The invention also provides an assay for a promoter of hydroxylation of HIF- α , for example prolyl hydroxylation at Pro 564 and/or Pro 402 which comprises providing HIF- α or a peptide analogue; incubating HIF- α or the peptide analogue under hypoxic conditions or conditions under which hydroxylation of HIF- α does not

occur in the absence of the test substance, and monitoring for hydroxylation of HIF- α or the peptide analogue thereof, such as at Pro 564 and/or Pro 402.

Accordingly, there is provided an assay for a promoter of hydroxylation of a proline residue in HIF- α , which comprises;

- 5 - providing HIF- α , or fragment thereof which includes a VHL-binding portion, under hypoxic conditions, said HIF- α or fragment thereof containing a proline residue in the VHL-binding domain;
- providing a putative hydroxylation promoting agent; and
- determining whether said agent provides for hydroxylation of said proline.

10 In the experiments described herein, HIF hydroxylases, and in particular PHD polypeptides, have been found to hydroxylate HIF- α at one or more proline residues within the pVHL binding domain. This hydroxylation mediates pVHL binding. Accordingly, the present invention provides an assay for a modulator of HIF hydroxylase activity comprising contacting a HIF hydroxylase and a substrate of the hydroxylase, preferably a prolyl-containing substrate, in the presence of a test
15 substance; and, determining the hydroxylase activity of the HIF hydroxylase, and in particular the prolyl hydroxylase thereof.

Such an assay may be used to identify inhibitors of HIF hydroxylase activity and are thus preferably carried out under conditions under which hydroxylation, and
20 in particular prolyl hydroxylation, takes place in the absence of the test substance. As an alternative, the assays may be used to look for promoters of hydroxylase activity, for example, by looking for increased hydroxylation of the proline substrate compared to an assay carried out in the absence of a test substance. Alternatively, the assays may be carried out under conditions in which hydroxylation is reduced or
25 absent, such as under hypoxic conditions and monitoring for the presence of or increased hydroxylation under such conditions.

Such an assay method may by virtue of using a specific HIF hydroxylase polypeptide be specific for inhibitors or promoters of the activity of that polypeptide and may by way of comparison be used to define inhibitors or activators that are
30 specific for that HIF hydroxylase and not active or less active on a different HIF

hydroxylase. In particular, such assays may be used to identify inhibitors or activators specific for a particular human HIF hydroxylase such as PHD 1, 2 or 3.

An assay method for obtaining an agent which modulates the activity of a HIF hydroxylase may include:

- 5 - contacting an HIF hydroxylase polypeptide and a substrate thereof, such as an HIF α polypeptide in the presence of a test substance; and,
- determining the hydroxylase activity of said HIF hydroxylase, preferably the prolyl hydroxylase activity, or HIF- α hydroxylase activity thereof.

10 In one embodiment the assay method may be for obtaining an agent which modulates the activity of an EGLN polypeptide and comprise:

- contacting an EGLN polypeptide and a test compound in the presence of an HIF polypeptide under conditions in which the EGLN polypeptide normally catalyses prolyl hydroxylation of said HIF polypeptide; and
- 15 - determining the HIF prolyl hydroxylase activity of said EGLN polypeptide.

Such assays may be performed under conditions in which the HIF hydroxylase/EGLN polypeptide normally catalyses hydroxylation and, in particular prolyl hydroxylation of a HIF α polypeptide. Suitable conditions may include pH 6.6
20 to 8.5 in an appropriate buffer (for example, Tris.HCl or MOPS) in the presence of 2-oxoglutarate, dioxygen and preferably ascorbate and ferrous iron.

Reducing agents such as dithiothreitol or tris(carboxyethyl)phosphine may also be present to optimise activity. Other enzymes such as catalase and protein disulphide isomerase may be used for the optimisation of activity. The enzymes,
25 such as protein disulphide isomerase, may be added in purified or unpurified form. Further components capable of promoting or facilitating the activity of protein disulphide isomerase may also be added.

In an alternative embodiment, the invention provides an assay method for identifying an agent which modulates the interaction of a EGLN polypeptide and a
30 HIF polypeptide comprising:

- contacting an EGLN polypeptide and a test compound in the presence of an HIF polypeptide, under conditions in which the EGLN polypeptide normally interacts with the HIF polypeptide; and

- determining the interaction of said HIF polypeptide and said EGLN polypeptide.

The present invention also provides an assay method for the identification of a HIF hydroxylase and in particular for the identification of a HIF prolyl hydroxylase. The method typically comprising:

- (a) providing a test polypeptide;
- 10 (b) bringing into contact a HIF polypeptide and the test polypeptide under conditions in which the HIF polypeptide is hydroxylated by a HIF hydroxylase; and
- (c) determining whether or not the HIF polypeptide is hydroxylated.

In one embodiment the assay method may, in step (b), bring the HIF polypeptide and test polypeptide into contact under conditions in which the HIF polypeptide is hydroxylated by a PHD (EGLN) polypeptide.

Typically, libraries of test polypeptides may be screened to identify a HIF hydroxylase, for example an expression library from a particular species or tissue may be screened or one produced under a particular set of conditions. Alternatively, candidate HIF hydroxylases identified on the basis of criteria such as sequence

20 homology or a particular protein structure may be assessed and their hydroxylase activity confirmed.

The HIF polypeptide used in the screening may be any of those described herein. It may be a human HIF polypeptide or a homolog from another species such as, for example, ceHIF. It may be from the species the library of test polypeptides is

25 derived from. The hydroxylation of the HIF polypeptide, and in particular the hydroxylation of proline, may be identified by any of the methods discussed herein. For example hydroxylation may be confirmed by using a functional assay based on the effect of the hydroxylation on the HIF polypeptide, such as its decreased stability.

Once a HIF hydroxylase has been identified it may be further characterised

30 by, for example, assessing whether or not it is inhibited by compounds such as dimethyloxalolylglycine (DMOG) a precursor or pro-drug for oxalolylglycine. The

effect of the HIF hydroxylase on HIF stability in the organism or tissue the hydroxylase is identified from may be assessed. The identified hydroxylase may be used in the same way as the other hydroxylases of the invention and in particular in the assays and therapeutic applications of the invention.

- 5 The present invention also provides an assay method for identifying alternative substrates of a HIF hydroxylase of the invention. Thus polypeptides or polypeptide analogues which can be hydroxylated and in particular have proline residues hydroxylated may be identified. The assay method typically comprises:
- 10 (b) contacting a test polypeptide with a HIF hydroxylase of the invention under conditions which HIF would normally be hydroxylated by the hydroxylase;
- (c) determining whether the polypeptide is hydroxylated.

Typically, hydroxylation and in particular prolyl hydroxylation of the test substance may be confirmed using any of the methods discussed herein.

- 15 The present invention also provides an assay method for identifying a polypeptide or polypeptide analogue capable of specifically interacting with a HIF hydroxylase of the invention and in particular which is capable of specifically binding to the active site of the HIF hydroxylase in a manner which mimics or resembles the binding of the normal substrate of the enzyme. The method typically
- 20 comprises:
- (a) contacting a test polypeptide with a HIF hydroxylase of the invention under conditions which HIF would bind to the hydroxylase;
- (b) determining whether the test polypeptide or analogue binds the hydroxylase.

- 25 The binding of the test polypeptide to the hydroxylase may be confirmed by any of the techniques discussed herein. In one embodiment binding of the polypeptide to a HIF hydroxylase may be identified by looking for co-immunoprecipitation of the test polypeptide with the hydroxylase. The ability of the test polypeptide to inhibit binding of HIF to the hydroxylase may also be used.

- 30 The alternative polypeptide substrates and the polypeptides identified as being capable of specifically binding the hydroxylases of the invention may be used

in the assay methods of the invention, for example, to identify modulators. Those polypeptides capable of preventing the normal interaction of HIF with a hydroxylase of the invention may also be used therapeutically.

The assays of the invention may also comprise modifying the agent identified. The assays may also comprise formulating the identified agent into a pharmaceutical composition. Typically, such pharmaceutical compositions may be for the treatment of a condition associated with increased or decreased HIF levels or activity.

Methods for monitoring modulation

The precise format of any of the screening or assay methods of the present invention may be varied by those of skill in the art using routine skill and knowledge. The skilled person is well aware of the need to additionally employ appropriate controlled experiments. The assays of the present invention may involve monitoring for hydroxylation of a suitable substrate (in particular monitoring for prolyl hydroxylation), monitoring for the utilisation of substrates and co-substrates, monitoring for the production of the expected products between the enzyme and its substrate. Assay methods of the present invention may also involve screening for the direct interaction between components in the system. Alternatively, assays may be carried out which monitor for downstream effects such as binding and subsequent destruction of HIF by VHL, alterations to the levels of HIF in the system and downstream effects mediated by HIF such as HIF mediated transcription using suitable reporter constructs or by monitoring for the upregulation of genes or alterations in the expression patterns of genes known to be regulated directly or indirectly by HIF.

Various methods for determining hydroxylation are known in the art and are described and exemplified herein. Any suitable method may be used for determining activity of the HIF hydroxylase such as by substrate or co-substrate utilization, product appearance such as peptide hydroxylation or down-stream effects mediated by hydroxylated or non-hydroxylated products.

Our finding that the Pro564 residue of HIF-1 α is hydroxylated by a prolyl-hydroxylase provides the basis for assay methods designed to screen for inhibitors or

promoters of this process. Any suitable method may be used to monitor for hydroxylation of HIF-1 α or a HIF polypeptide or analogue thereof. Assays may be carried out to monitor directly for hydroxylation of the relevant proline residue or another position. Alternatively, assays may be carried out to monitor for depletion of co-factors and co-substrates. Alternatively, such assays may monitor the downstream effects of hydroxylation of HIF or indeed inhibition of hydroxylation of HIF, for example, by monitoring the interaction between HIF and VHL levels of HIF protein or HIF mediated transcription. Alternatively, reporter gene constructs driven by HIF regulated promoters may be used. Assays are also provided for the identification of enhancers of the activity of the HIF hydroxylase and in particular of the HIF prolyl hydroxylase activity of these enzymes. The assay may be used to identify an enhancer a human HIF hydroxylases and, in particular, of PHD 1,2 or 3. Such enhancers may be used to reduce HIF α activity.

In one embodiment, to perform an assay for an inhibitor of VHL-mediated HIF- α destruction a suitable substrate of the HIF hydroxylase is provided. This may be HIF- α or a fragment thereof which includes a VHL binding portion and which included the Pro564 and/or Pro 402 residue is provided. The substrate may not be hydroxylated at the Pro564 and/or Pro 402 position. This may be achieved by providing synthetic polypeptide substrates, or by producing HIF- α polypeptides in bacterial cells, insect cells or mammalian cells or in *in vitro* transcription and translation systems. Alternatively, assays may be carried out over a selected time course such that the substrate is produced during the course of the assay, initially in un-hydroxylated form.

The substrate, enzyme and potential inhibitor compound may be incubated together under conditions which, in the absence of inhibitor provide for hydroxylation of Pro564 and/or Pro 402, and the effect of the inhibitor may be determined by determining hydroxylation of the substrate. This may be accomplished by any suitable means. Small polypeptide substrates may be recovered and subject to physical analysis, such as mass spectrometry or chromatography, or to functional analysis, such as the ability to bind to VHL (or displace a reporter

molecule from VHL) and be targeted for destruction. Such methods are known as such in the art and may be practiced using routine skill and knowledge.

Determination may be quantitative or qualitative. In both cases, but particularly in the latter, qualitative determination may be carried out in comparison to a suitable control, e.g. a substrate incubated without the potential inhibitor.

Inhibitor compounds which are identified in this manner may be recovered and formulated as described above for polypeptides of the invention.

Another assay of the invention is for a promoter of hydroxylation of HIF- α subunits. Typically, a HIF- α subunit or portion thereof is prepared as described above, and incubated under hypoxic conditions. By "hypoxic", it is meant less than 5%, preferably less than 3%, more preferably less than 1%, and preferably less than 0.5%, such as less than 0.1% O₂. The HIF- α subunit is incubated with a cell extract which includes the HIF hydroxylase as described above, optionally further in the presence of a source of ferrous (FeII) ions and/or ascorbate. A suitable concentration of ferrous ions is in the range of from 1 to 500 μ M, such as from 25 to 250 μ M and in particular from 50-200 μ M. Ferrous ions may be supplied in the form of ferrous chloride, ferrous sulphate, and the like. Ascorbate may be provided in the form of a salt, such as sodium ascorbate, and in a concentration range of from 0.1 to 10mM, such as from 1 to 5 mM. Another cofactor is α -ketoglutarate, which may also be supplied in the form of a salt at a range of from 0.1 to 5mM, such as from 1 to 5 mM.

In this embodiment of the invention, the substrate will be incubated in the presence of a potential hydroxylation promoting agent, and the effect of the agent determined, by determining the hydroxylation of the Pro564 and/or Pro 402. As with the assay of the other aspect of the invention described above, determination may be quantitative or qualitative, and in either case determined relative to a suitable control.

The interaction between the polypeptides may be studied *in vitro* by labelling one with a detectable label and bringing it into contact with the other which has been immobilised on a solid support. Suitable detectable labels include ³⁵S, which may be incorporated into recombinantly produced peptides and polypeptides.

Recombinantly produced peptides and polypeptides may also be expressed as a fusion protein containing an epitope which can be labelled with an antibody.

Fusion proteins may, for example, incorporate six histidine residues at either the N-terminus or C-terminus of the recombinant protein. Such a histidine tag may be used for purification of the protein by using commercially available columns which contain a metal ion, either nickel or cobalt (Clontech, Palo Alto, CA, USA).

5 These tags also serve for detecting the protein using commercially available monoclonal antibodies directed against the six histidine residues (Clontech, Palo Alto, CA, USA).


The protein which is immobilized on a solid support may be immobilized using an antibody against that protein bound to a solid support or via other
10 technologies which are known *per se*. A preferred *in vitro* interaction may utilise a fusion protein including glutathione-S-transferase (GST). This may be immobilized on glutathione agarose beads. In an *in vitro* assay format of the type described above, a test compound can be assayed by determining its ability to diminish the amount of labelled peptide or polypeptide which binds to the immobilized GST-
15 fusion polypeptide. This may be determined by fractionating the glutathione-agarose beads by SDS-polyacrylamide gel electrophoresis. Alternatively, the beads may be rinsed to remove unbound protein and the amount of protein which has bound can be determined for example, by counting the amount of label present in a suitable scintillation counter.

20 Thus, assays in accordance with the present invention may involve monitoring for the interaction between VHL and HIF. The interaction between HIF and VHL is mediated by hydroxylation of HIF. The VHL-HIF interaction leads to ubiquitylation of HIF. Assays to monitor for test substances which interfere with the interaction between HIF and VHL, and in particular which interfere with
25 hydroxylation of HIF may be monitored by any suitable method using HIF associated regulation. For example, in assay systems making use of recombinant HIF hydroxylase in accordance with the present invention, or in which HIF hydroxylase expression is upregulated within a cell, the effect of test substances can be monitored through monitoring the levels of HIF in the cell. Alternatively, transcription and
30 expression of genes known to be upregulated or down regulated by the presence of HIF could be monitored. In particular, upregulation of HIF regulated genes would

demonstrate inhibition of prolyl hydroxylation whereas down regulation would suggest enhancement or promotion of prolyl hydroxylation.

In alternative embodiments, reporter constructs may be provided in which promoters mediated by HIF are provided operably linked to a reporter gene. Any
5 suitable reporter gene could be used, such as for example enzymes which may then be used in colorimetric, fluorometric, fluorescence resonance or spectrometric assays.

HIF hydroxylase is a 2OG dependent oxygenase which catalyses the following reaction, in which R is HIF α and ROH singly is hydroxylated HIF α ;



10

Typically, the hydroxylation is prolyl hydroxylation. The hydroxylase may catalyse more than one hydroxylation of HIF- α .

In the assay methods described herein, typically the HIF hydroxylase and the substrate of the hydroxylase are contacted in the presence of a co-substrate, such as
15 2-oxoglutarate (2OG). The hydroxylase activity of the HIF hydroxylase may be determined by determining the turnover of the co-substrate. This may be achieved by determining the presence and/or amount of reaction products, such as hydroxylated substrate or succinic acid. The amount of product may be determined relative to the amount of substrate. Typically, in such embodiments the substrate may be an HIF- α
20 polypeptide and, for example, the product measured may be hydroxylated HIF- α polypeptide.

HIF α prolyl hydroxylase activity may be determined by determining the turnover of said 2OG to succinate and CO₂, as described in Myllyharju J. *et al* EMBO J. 16 (6): 1173-1180 (1991) or as in Cunliffe C.J. *et al* Biochem. J. 240 617-

619 (1986), or other suitable assays for CO₂, bicarbonate or succinate production. These methods may be used in the assays of the invention and in particular to assess the HIF- α prolyl hydroxylase activity of the HIF hydroxylase of the invention, including that of human HIF hydroxylases and in particular of PHD or EGLN
5 polypeptides of the invention. Such assays can be modified to high throughput format and the invention encompasses such high throughput assays for hydroxylase activity.

Alternatively, the end-point determination may be based on conversion of HIF α or peptide fragments (including synthetic and recombinant peptides) derived
10 from HIF α into detectable products. Peptides may be modified to facilitate the assays so that they can be rapidly carried out and may be suitable for high throughput screening.

For example, reverse phase HPLC (C-18 octadecylsilane column), as exemplified herein, may be used to separate starting synthetic peptide substrates for
15 HIF hydroxylase from the hydroxylated products, as the latter have a shorter retention time in the column. Modifications of this assay or alternative assays for HIF hydroxylase activity may employ, for example, mass spectrometric, spectroscopic, and/or fluorescence techniques as are well known in the art (Masimirembwa C. *et al* Combinatorial Chemistry & High Throughput Screening (2001) 4 (3) 245-263,
20 Owicki J. (2000) J. Biomol. Screen. 5 (5) 297-305, Gershkovich A *et al* (1996) J. Biochem. & Biophys. Meths. 33 (3) 135-162, Kraaft G. *et al* (1994) Meths. Enzymol. 241 70-86). Fluorescent techniques may employ versions of the substrate modified
21 in such a way as to carry out or optimise spectroscopic or fluorescence assays.

For example, HIF α polypeptide may be immobilised e.g. on a bead or plate,
25 and hydroxylation of the appropriate residue detected using an antibody or other binding molecule which binds the pVHL binding domain of HIF α with a different affinity when a proline residue such as proline 402 or proline 564 is hydroxylated from when the residue is not hydroxylated. Such antibodies may be obtained by means of standard techniques which are well known in the art, e.g. using a
30 hydroxylated HIF α peptide.

Binding of a molecule which discriminates between the hydroxylated and non-hydroxylated form of a HIF α polypeptide may be assessed using any technique available to those skilled in the art, which may involve determination of the presence of a suitable label.

5 Assays may be used to screen for inhibitors of HIF hydroxylase and in particular for inhibitors of HIF prolyl hydroxylase (HPH) activity in a similar way to that described for the human prolyl hydroxylase involved in collagen biosynthesis (CPH) (Cunliffe *et al.* Biochemical J. 240 611-619 (1986); Cunliffe CJ *et al.* Biochem. J. 239 311-315 (1986), Franklin TJ and Hitchen M Biochem. J. 261: 127-10 130 (1989) Franklin TJ. *et al.* Biochemical Society Transactions 19 (4): 812-815 (1991)).

HIF prolyl-hydroxylase activity of a HIF hydroxylase polypeptide may be determined by determining the hydroxylation of one or more proline residues of the substrate of the HIF hydroxylase used, which will typically be a HIF α 15 polypeptide. Preferably, the hydroxylation of one or more proline residues within the pVHL binding domain of the HIF- α polypeptide, for example, proline 402 and/or proline 564. For convenience, these proline residues are referred to herein as Pro402 and Pro564 or position 402 and position 564. It will be understood that this terminology is also applied to polypeptides which contain far fewer than 564 20 residues, and to other HIF- α isoforms where the equivalent proline residue may occur at a slightly different position.

Assay methods of the present invention may also take the form of an *in vivo* assay. The *in vivo* assay may be performed in a cell line such as a yeast strain in which the relevant polypeptides or peptides are expressed from one or more vectors 25 introduced into the cell.

C.elegans assay systems

Our finding that the HIF-VHL interaction is conserved in *C.elegans* provides a system to study the interaction in an *in vivo* environment, and its consequences.

Thus in a further aspect, the invention provides an assay for a modulator of 30 HIF-VHL interaction, said method comprising:

- providing a *C.elegans* which has wild-type HIF and VHL genes in normoxic or hypoxic conditions (wherein hypoxic conditions are as defined above);

- exposing said *C.elegans* to a potential modulator of the HIF-VHL interaction; and

5 - determining the extent to which the modulator promotes or decreases the interaction between HIF and VHL in said *C.elegans*.

The determining may comprise immunoprecipitating one or other of the HIF and VHL components and then determining, e.g. by antibody detection, the amount of the other of the HIF and VHL component which is associated with the

10 immunoprecipitated protein. Radiolabelling of one of the two proteins may allow determination of the amount of VHL or HIF captured. Alternatively, a reporter gene may be linked to a HIF-responsive promoter, and the amount of HIF in the subject *C.elegans* determined by measuring the activity of a reporter gene product, such as green fluorescent protein, luciferase, chloramphenicol acetyl transferase, beta-
15 galactosidase, and the like.

In alternative aspects of the invention, the source of 2-oxoglutarate dependent dioxygenase and in particular the prolyl-hydroxylase of the invention is a HIF hydroxylase according to the invention. Such polypeptides may be introduced using recombinant systems so as to allow for the assay of inhibitory, augmenting, blocking or
20 other modulating activities that show differential effects among the said HIF hydroxylase.

In vivo assays

The assays may be carried out using cell based, organ based or whole animal assays conducted *in vivo*. Such assays may utilize the endogenous expression of the HIF
25 hydroxylase nucleotides and/or polypeptides. In other forms of the invention, upregulation of specific endogenous HIF hydroxylases may be achieved by stimulators of the expression thereof. Such stimulators may be growth factor such as platelet derived growth factor or angiotensin II, or chemicals such as phorbol esters that are known to upregulate specific HIF hydroxylases. In another form of the invention, nucleotide
30 constructs may be introduced into cells or transgenic animals to increase production of one or more specific HIF hydroxylases. Alternatively nucleotide constructs may be

introduced into cells so as to reduce or abrogate expression of one or more specific HIF hydroxylases. Appropriate methods that include but are not limited to homologous recombination, antisense expression, ribozyme expression and RNA interference are outlined herein and known by those skilled in the art.

5 Tissue culture cells, organs, animals and other biological systems, obtained by the aforementioned forms of the invention, may be used to provide a further source of a HIF hydroxylase, or may be used for the assay, or especially comparative assay, of the activity of test substances may inhibit, augment, block or otherwise modulate the activity of specific HIF hydroxylases.

10 The activity of the HIF hydroxylases may be assayed by any of the aforementioned methods or by cell, tissue, or other assays conducted *in vivo* that measure the effects of altered activity of the HIF hydroxylases. A preferred form of these assays measures the level of a HIF- α polypeptide or the level of activity of a HIF- α polypeptide that is a substrate for the PHD polypeptide.

15 The level of a HIF- α peptide may be measured by such methods as immunoblotting, immunoprecipitation, or other immunological methods using specific antibodies and methods that are known to those skilled in the art. The amounts and the activity of HIF- α peptides can be related to each other but are not necessarily related to each other. Therefore in a further form of the invention, the activity of a HIF- α peptide that is a
20 target for a HIF hydroxylase is assayed by measurement of transcriptional activity or another property of the said HIF- α polypeptide.

HIF- α polypeptides are known to form complexes with other molecules that include other HIF subunits and co-activator molecules such as p300. In this form HIF complexes activate hypoxia response elements that are found in the promoters and/or
25 enhancers of endogenous genes that are regulated by the said HIF complexes. Such hypoxia response elements may also be isolated and operationally linked to reporter genes so as to assay the activity of the HIF complex through detection and/or quantitation of the reporter gene or its product. Therefore in a further form of the invention the activity of a HIF- α polypeptide that is regulated by its cognate HIF
30 hydroxylase will be assayed by measuring the effects of the HIF complex on the

expression of an endogenous gene or reporter gene that is functionally linked to a HIF binding hypoxia response element. Examples of endogenous genes that are regulated in this way are to be found in the role of the aryl hydrocarbon nuclear translocator (ARNT) in hypoxic induction of gene expression, see for example, Studies in ARNT-deficient
5 cells. S.M. Wood, J.M. Gleadle, C.W. Pugh, O. Hankinson, P.J. Ratcliffe. *Journal of Biological Chemistry* 271 (1996) 15117-15123, and Hypoxia inducible expression of tumor-associated carbonic anhydrases, C.C. Wykoff, N.J.P. Beasley, K.J. Turner, J. Pastorek, A. Sibtain. G.D. Wilson, H. Turley, K. Talks, P.H. Maxwell, C.W. Pugh, P.J. Ratcliffe, A.L. Harris. *Cancer Research* 60 (2000) 7075-7083. Examples include but
10 are not limited to glucose transporter isoform 1, phosphoglycerate kinase-1, carbon anhydrase isoform 9, vascular endothelial growth factor. Each of said genes contains one or hypoxia response elements that may be isolated and operationally linked as single or multiple copies to a reporter gene for the measurement of activity of a HIF- α polypeptide that varies in accordance with the activity of a HIF hydroxylase.

15 The activity of genes or gene products that are regulated by a HIF- α polypeptide in accordance with the activity of a HIF hydroxylase affects cellular, organ, and animal physiology in a manner that provide further aspects of the invention. Thus a further embodiment of the invention provides for assays that utilise a specific functional response that is regulated in accordance with the activity of a HIF- α polypeptide in
20 accordance with the activity of a HIF hydroxylase. Such responses include the uptake rate of glucose or glucose analogues that are not metabolized, the ingrowth of blood vessels by angiogenesis, the activity of a carbonic anhydrase enzyme. It is recognised that many other responses that operate at a cellular or systemic level are controlled by the activity of a HIF- α polypeptide in accordance with the activity of a HIF hydroxylase
25 and may be utilized as assays of the said HIF hydroxylase activity in further aspects of the invention.

A HIF- α polypeptide that is a substrate for a HIF hydroxylase may be fused to a further polypeptide so as to cause the activity of the said HIF hydroxylase to regulate the activity of the fusion peptide. Accordingly a further form of the invention provides
30 for the assay of the activity of a fusion polypeptide. In the preferred form such a fusion

polypeptide may contain the whole or part of a HIF- α polypeptide, for example human HIF-1 α residues 344-698, 344-417, 554-698, 652-826, or 775-826 linked to a heterologous transcription factor and expressed together with its cognate DNA response element. The Gal4 DNA binding domain including the amino acids 1-143 together with
5 the Gal binding upstream activating sequence (UAS) is an example of such a transcription factor and cognate DNA response element whose operation can be assayed by those skilled in the art.

In a preferred embodiment, the assays discussed herein relate to, or utilise, human HIF hydroxylases and in particular PHD 1, 2 or 3. These may also be referred
10 to as EGLN polypeptides.

Test Compounds

Compounds which may be screened using the assay methods described herein may be natural or synthetic chemical compounds used in drug screening programmes. Extracts of plants, microbes or other organisms, which contain several
15 characterised or uncharacterised components may also be used.

Combinatorial library technology (including solid phase synthesis and parallel synthesis methodologies) provides an efficient way of testing a potentially vast number of different substances for ability to modulate an interaction. Such libraries and their use are known in the art, for all manner of natural products, small
20 molecules and peptides, among others. The use of peptide libraries may be preferred in certain circumstances.

Potential inhibitor compounds may be polypeptides, small molecules such as molecules from commercially available combinatorial libraries, or the like. Small molecule compounds which may be used include 2-oxoglutarate analogues, or HIF- α
25 analogues, or those that incorporate features of both 2-oxoglutarate and affect HIF- α , which inhibit the action of the enzyme. We have found in particular that compounds dimethyl-oxalylglycine, N-oxalylglycine and N-oxalyl-2S-alanine and certain thiols all act as inhibitors of the HIF- α hydroxylase. N-oxalyl-2R-alanine, an enantiomer of N-oxalyl-2S-alanine, is also an inhibitor of HIF hydroxylase and may be used in the
30 invention. Thus the invention provides the use of a compound which acts as a

hydroxylase inhibitor, and in particular prolyl hydroxylase inhibitor for the manufacture of a medicament for the treatment of a condition in a patient which requires the promotion of cell growth, such as angiogenesis. The invention also provides a method of treatment of a patient suffering from a condition which is

5 treatable by promoting cell growth, which comprises administering to said patient an effective amount of a HIF hydroxylase inhibitor. Such inhibitors include dimethyl-oxalylglycine, N-oxalylglycine and N-oxalyl-2S-alanine, and salts thereof. More generally, such inhibitors include other N-oxalyl-amino acid compounds and salts thereof, wherein the amino acids are either naturally occurring amino acids or

10 synthetic amino acids with a side chain which provides for the compound to act as an inhibitor. Such side chains include hydrocarbyl side chains containing a carbon chain which may be straight or branched, optionally including one or two heteroatoms such as N, O or S, and optionally substituted by a group such as halogen (particularly fluoro, chloro, bromo or iodo), thiol, hydroxy, methoxy, amino, mono-

15 or di- C1-3 alkyl amino or nitro. Salts include pharmaceutically acceptable salts such as sodium, potassium, magnesium and the like.

Potential promoting agents may be screened from a wide variety of sources, particularly from libraries of small compounds which are commercially available. Oxygen-containing compounds may be included in candidate compounds to be

20 screened, for example 2-oxoglutarate analogues.

CPH Inhibitors

Inhibitors of the 2-OG dependent enzyme collagen prolyl hydroxylase (CPH) are well known in the art and have been previously proposed for use in the treatment of lung fibrosis, skin fibrosis (scleroderma), atherosclerosis and other conditions

25 associated with collagen biosynthesis. Inhibitors of parahydroxyphenylpyruvate oxygenase (a non-haem oxygenase employing ferrous iron as a co-factor) such as triketones are used as herbicides (Lee D. *et al* (1998) Pestic. Sci. 54(4) 377-384).

The present inventors have now found that certain of these CPH inhibitors also inhibit the biological activity of HIF hydroxylases and in particular the ability

30 of the HIF hydroxylase to catalyse prolyl hydroxylation of HIF (HPH activity).

Another aspect of the present invention therefore provides the use of a CPH inhibitor or modified CPH inhibitor which inhibits the biological activity of a HIF hydroxylase, and in particular its HPH activity, in the manufacture of a medicament for use in the treatment of a condition associated with reduced or suboptimal HIF levels or activity, for example ischaemia, wound healing, auto-, allo-, and xeno-transplantation, systemic high blood pressure, cancer, and inflammatory disorders. In one embodiment a CPH inhibitor which inhibits the biological activity of a human HIF hydroxylase such as a PHD polypeptide (EGLN polypeptide) may be used.

CPH inhibitors which inhibit HIF hydroxylases, and in particular the prolyl hydroxylase activity (HPH activity) of a HIF hydroxylase, may be modified to generate selective inhibitors of HIF hydroxylases and in particular of HPH activity. Further, the discovery allows for the development of collagen prolyl hydroxylase inhibitors that do not inhibit HIF hydroxylases, and in particular HPH, by the use of comparative screening assays.

Another aspect of the present invention therefore provides the use of a modulator of a HIF hydroxylase in the manufacture of a medicament for the treatment of a condition associated with reduced HIF levels or activity as described above and below.

Such an modulator may be a selective inhibitor. A selective inhibitor is an inhibitor which shows a greater level of inhibition of a HIF hydroxylase than on other enzymes including collagen proly hydroxylase. In particular, a selective inhibitor is one which inhibits HPH activity relative to CPH activity as described above.

HPH Modulators

Various methods and uses of modulators which inhibit, potentiate, increase or stimulate hydroxylation of HIF- α by HIF hydroxylase are provided as further aspects of the present invention.

The purpose of disruption, interference with or modulation of the hydroxylation of HIF-1 α by a HIF hydroxylase may be to modulate cellular functions such as angiogenesis, erythropoiesis, energy metabolism, inflammation, matrix metabolism, vasomotor function, and apoptotic/proliferative responses and

pathophysiological responses to ischaemia/hypoxia, all of which are mediated by HIF α , as discussed above and further below.

A test compound which increases, potentiates, stimulates, disrupts, reduces, interferes with or wholly or partially abolishes hydroxylation of HIF- α polypeptide and which may thereby modulate HIF hydroxylation activity, may be identified
5 and/or obtained using the assay methods described herein.

Agents which increase or potentiate hydroxylation, and in particular prolyl hydroxylation of HIF, may be identified and/or obtained under conditions which, in the absence of a positively-testing agent, limit or prevent hydroxylation. Such agents
10 may be used to potentiate, increase, enhance or stimulate the function of a HIF hydroxylase, and may have an effect on cells under hypoxic conditions such as those found in tumours, in which the lack of hydroxylation leads to the accumulation of HIF α and the concomitant promotion of angiogenesis and other growth promoting events.

15 The term 'agent' includes a compound having one of the formulae I to XXVIII as described herein in particular a compound shown in Table 3.

Methods of determining the presence of, and optionally quantifying the amount of HIF hydroxylase in a test sample may have a diagnostic or prognostic purpose, e.g. in the diagnosis or prognosis of any medical condition discussed herein
20 (e.g. a proliferative disorder such as cancer) or in the evaluation of a therapy to treat such a condition.

In various aspects, the present invention provides an agent or compound identified by a screening method of the invention to be a modulator of HIF α hydroxylation e.g. a substance which inhibits or reduces, increases or potentiates the
25 hydroxylase activity of a HIF hydroxylase.

Following identification of a modulator, the substance may be purified and/or investigated further (e.g. modified) and/or manufactured. A modulator may be used to obtain peptidyl or non-peptidyl mimetics, e.g. by methods well known to those skilled in the art and discussed herein. A modulator may be modified, for example to
30 increase selectively, as described herein. It may be used in a therapeutic context as discussed below.

Agents according to the present invention, which are useful in modulating the hydroxylation of HIF- α and therefore the modulation of HIF's intracellular levels and hence one or more of its cellular functions, may modulate the hydroxylase activity of the HIF hydroxylase. Such agents may specifically inhibit the ability of the HIF
5 hydroxylase, and in particular of a PHD polypeptide, to hydroxylate the appropriate residue of HIF- α . Assays and screens for such agents are provided as described above in accordance with the present invention, along with the agents themselves and their use in modulating the hydroxylation and thereby the function of HIF- α .

An agent able to inhibit hydroxylation of HIF- α by a HIF hydroxylase may
10 include a substance able to affect the catalytic properties of the enzymatically active site of the hydroxylase. An inhibitor of hydroxylation may interact with the HIF hydroxylase within the active prolyl hydroxylase domain, for example within the HXD[X]_nH or jelly roll motifs described herein or in the HXE[X]_nH motifs. Residues within this domain are involved in interaction with HIF- α and catalysis of
15 the hydroxylation. An inhibitor may, for example, interact with His358 or ARG 557 of the EGLN2/PHD1 sequence using the EGL9 numbering system or the equivalent residue of other HIF hydroxylases, or may interact with residues in the region between residues 369 and 389 or other residues of the jelly roll motif of the PHD1 sequence or the equivalent residues of other HIF hydroxylases. Residues outside of
20 the domain may also be involved in interacting with HIF- α and agents which interfere with such interaction may also affect the hydroxylation as discussed elsewhere herein. Alternatively or additionally, the inhibitor may bind in such a way as to inhibit dioxygen binding. For example, it is appreciated that dioxygen may approach the iron in the HIF hydroxylase from a different direction to the HIF- α
25 substrate and in particular through a tunnel through the centre of the jelly roll motif. Inhibitors may bind to residues within or at the entrance to this tunnel.


Further aspects of the present invention relate to methods for modulating the amount of HIF polypeptide in a cell. Such a method may comprise contacting the cell with a substance which inhibits the hydroxylase activity of a HIF hydroxylase and in
30 particular its prolyl hydroxylase activity.

A suitable substance is an agent as described herein. A substance which is a selective HIF hydroxylase inhibitor may be used in such a method. A selective HIF hydroxylase inhibitor is an inhibitor which inhibits the biological activity of a HIF hydroxylase but does not inhibit biological activity of a collagen prolyl hydroxylase, as described herein and in particular one which inhibits HPH activity of a HIF hydroxylase but not CPH activity of a collagen prolyl hydroxylase.

Examples of HPH Modulators

Compounds which modulate 2OG oxygenases, in particular collagen 4-prolyl hydroxylase, may be useful as modulators of HIF prolyl hydroxylase, or may be used as 'lead' compounds which may be modified and/or optimised to develop modulators of HIF prolyl hydroxylase, in particular selective modulators.


Some of these compounds generally possess the formula:




where the group R^1 is capable of forming an electrostatic interaction with the side chain of the arginine which together with other residues binds the 5-carboxylate of 2-oxoglutarate during catalysis, $A*B$ is a chain of two atoms which are, independently, carbon, oxygen, nitrogen or sulphur, which chain can be functionalised, y is 0 or 1 and $C*D$ is a chain of two atoms which are, independently, carbon, oxygen, nitrogen, or sulphur, which chain can be functionalised, A , B , C and D being linked to one another by single and/or double and/or triple bonds, such that when y is 0 or 1 at least one of the atoms of which is capable of chelating with a metal group and when y is 1 said chain is attached to R^2 which is capable of chelating with a metal group. Generally at least one of A , B , C and D is not carbon. Typical chains include $C-N-C$, $C-C-C=O$ and $C-O-C-C$. The chain atoms can form part of a ring, such as pyrrolidine and tetrahydro-pyran, and unsaturated derivatives thereof including pyridine and pyran or partially hydrogenated pyrans. The rings can be fused. When $y = 0$ C and/or D is attached to, for example, $=S$, $=O$, $-SH$ or $-OH$. Typically R^1 is an acid group such as carboxylate, $-SO_3H$, $-B(OH)_2$ or $-PO_3H_2$. Typical values for R^2 include $-SH$, $-OH$, $-CO_2H$, $-SO_3H$, $-B(OH)_2$ or $-PO_3H_2$, $-NHOH$, $-CONHR^3$, $-CONHOR^3$, $-CONR^3OH$ and $-CONR^3OR^3$ where R^3 is a branched or straight chain alkyl group of 1 to 6 carbon atoms which can be functionalised. Preferred

compounds will have typical values for more than one group, for example for all groups.

One class of compounds which have been found to modulate the activity of HIF α hydroxylases and in particular their prolyl hydroxylase activity are oxalo-amino acid derivatives, including oxalo derivatives having one of the following
 5 general formulae (I to IV), for example compounds having the formula V, VI or VII;
 I: N-oxalo inhibitors for in vitro use




II:




10

III: N-oxalo inhibitors for in vivo use




IV:


55



V: (N-oxalyl glycine)




5 VI: (N-oxalyl alanine)



such as oxalyl-L-alanine (IS70) as well as oxalyl-D-alanine (IS71)

VII: (N-oxalyl valine)



- 10 where R¹ and R may independently be H, a branched or straight C₁ to C₆ alkyl chain, especially methyl, which can be functionalised, e.g. as -C₂H₄CO₂C₂H₅, any natural amino acid side chain such as alanine, valine and glutamic acid, a 4 to 7 membered

heterocyclic ring optionally containing 1 or 2 N, S, O or P atoms or a 5 or 6 membered aromatic ring, optionally containing 1 or 2 N, O or S atoms, such as phenyl or naphthyl which may be fused to another ring or a said alkyl chain substituted by a said aromatic ring;

- 5 R^2 is C1 to C6 alkyl chain which may be functionalised such that R^2 is $(CR^1R^1)_n$ where $n = 1$ to 6 and where the R^1 groups may be the same or different and are as defined above or R^2 is absent;

- X is NH, NR'' , where R'' is OH, a branched or straight C_1 to C_6 alkyl chain which can be functionalised, or O i.e. XR is O-alkyl having a branched or straight C_1 to C_6 alkyl chain, especially MeO, which can be functionalised; and,
10 Y is O or S.

The said alkyl groups and chains are typically functionalised by fluorine, alcohol, thiol, a carboxylic acid, phosphonic or phosphinic acid, sulphonic acid or other chelating group, in the case of the chains typically via an alkyl group.

- 15 Formula I is further exemplified by compound IS3 and oxalylglycine (IS2), Formula II is by oxaloylamino-L-alanine, and Formula III by compounds IS1 and dimethyloxaloylglycine (MMOG) in Table 3 as well as the methyl esters of methyloxalyl-L-alanine (IS80) and methyloxalyl-D-alanine (IS81), methyloxalyl-L-alanine (IS68) and methoxalyl-D-alanine (IS69), diethyl N-methoxyoxalyl-L-glutamate (IS12), -oxalyl-L-glutamate (IS13), oxalyl S-alanine (IS70) and oxalyl R-alanine (IS71).
20

These compounds may be obtained by synthesis as described in Cunliffe *et al* (1992) J. Med. Chem. 35 2652-2658.

- The present inventors have found that N-oxalo derivatives of amino acids, which are known to inhibit collagen prolyl-hydroxylase (CPH), inhibit the
25 modification of HIF- α by HIF hydroxylase and in particular by inhibiting their prolyl hydroxylase activity (HPH). This leads to reduced pVHL binding and increased cellular HIF levels.


- N-oxaloglycine has been found to be an inhibitor of both CPH and HPH.
30 However, (RS)- N-oxaloalanine is a poor inhibitor of CPH compared to N-oxaloglycine and (S) oxaloalanine is a preferred inhibitor of HIF hydroxylase, and in

particular of HPH activity, as described below. (S)-oxalovaline has little or no activity as an HPH inhibitor.


Selectivity for particular HIF hydroxylase may be increased or enhanced by modification of the oxaloglycine backbone. N-Oxalo amino acid derivatives may be converted into methyl or ethyl ester form for use as a 'pro drug' as described in E. Baeder *et al.* (1994) Biochem. J. 300.525-530 and exemplified in Table 2.

Another class of compounds of interest for use in the modulation of HIF hydroxylases, and in particular of HPH activity, are hydroxamic acid derivatives of the general formulae (VIII-XII):

VIII: (Alahopcin)




IX:




X:


58




XI:




5 XII:



XIII:



XIIIA:



where R^I to R^{VI} may independently be H, OH, a branched or straight C_1 to C_6 alkyl chain, optionally with 1, 2, 3, 4 or 5 halo substitutions, which can be functionalised, a 4 to 7 membered heterocyclic ring optionally containing 1 or 2 N, S, O or P atoms, or a 5 or 6 membered aromatic ring, optionally containing 1 or 2 N, O or S atoms which may be fused to another ring or a said alkyl chain substituted by a said aromatic ring such that R^{III} can also be NH_2 or a salt thereof such as HCl, or NHR^{IX} where R^{IX} is acyl, such as unsubstituted or substituted alkanoyl such as acetyl or phenoxyacetyl; and,

XR is OH, NH_2 or NHR^X , where R^X is OH, a branched or straight C_1 to C_6 alkyl chain, or O-alkyl having a branched or straight C_1 to C_6 alkyl chain.

Preferred compounds are those where R is methyl or benzyl and/or R^{II} is hydrogen and/or R^{III} is NH_2 or $NHCOCH_2O_6H_5$.

Hydroxamic acids may be obtained as described in Walter M. W. *et al* (1999) Bioorg. Chem. 27 (1): 35-40.

Hydroxamic acids, including cyclic and natural products such as alahopcin (Higashide E, *et al* (1995) J. Antibiotics 38: 285-295), are known to be inhibitors of CPH and are therefore of interest for the inhibition of HIF hydroxylases and in particular of HPH. Modulators based on these compounds, which are specific for HPH, may be developed using the methods described within.


Compounds of formula X are exemplified by benzo-hydroxamic acid and compounds of formula XI, which are a sub-class of formula X, are exemplified by compounds NK45, NK46, NK47 and NK84 in Table 3.

Another class of compounds of interest for use in the modulation of HIF hydroxylases, and in particular of HPH activity, are hydroxylated aromatic

60


compounds including catechols, phenanthrolines and hydroxanthroquinones,
including hydroxyanthroquinones of the formulae;

XIV:




5

XV:



XVA




where R¹ to R¹¹ may independently be H, a branched or straight C₁ to C₆ alkyl chain,
OH, O-alkyl having a branched or straight C₁ to C₆ alkyl chain optionally containing
10 1 or 2 N, O or S atoms, COOH, a branched or straight C₁ to C₆ alkyl ester

61

(alkoxycarbonyl), a 4 to 7 membered heterocyclic ring optionally containing 1 or 2 N, S, O or P atoms or

a 5 or 6 membered aromatic ring, optionally containing 1 or more N, O or S atoms, which can be fused to another ring, or a said alkyl chain substituted by a said

5 aromatic ring, and



Dihydroxybenzoate (EDB), as shown in Table 3, and 3,4-dihydroxybenzoic acid (protocatechuic acid) are examples of a compound of formula XV.

These compounds of formula XIV are known CPH inhibitors (Franklin *et al* (2001) Biochem. J. 353: 333-338, Cunliffe *et al* Biochem J. (1986) 239(2) 311-315, Franklin *et al* (1989) Biochem. J. 261 (1) 127-130). Modulators based on these compounds which are specific for HPH may be developed using the methods described within.

Specific compounds of formula XVA include:


15

20


62

R1


5




R2



R3



10 R4




Thus typically R^1, R^3, R^7, R^8 and R^{11} are hydrogen and R^2, R^4, R^9 and R^{10} are OH.

Another class of compounds of interest for use in the modulation of HIF


- 5 hydroxylases, and in particular of HPH activity are N-containing heterocyclic compounds which have one of the following general formulae:

XVI: 3-hydroxyquinolone-2-carboximide derivatives




- 10 where R^1 to R^5 may be H, a branched or straight C_1 to C_6 alkyl chain such as Me, a 4 to 7 membered heterocyclic ring optionally containing 1 or more N, S, O or P atoms, or a 5 or 6 membered aromatic ring, optionally containing 1 or more N, O or S atoms, which can be fused to another ring, or a said alkyl chain substituted by a said aromatic group, A = substituted alkylene, B=CO₂H, NHSO₂CF₃, tetrazolyl,
- 15 imidazolyl or 3-hydroxyisoxazolyl, and m is 0 or 1.

XVII: pyridine and pyridine N-oxide derivatives




XVIII:



where R^I to R^{IV} may independently be H, a branched or straight chain alkyl of from 1 to 6 C atoms, a halogen group (i.e. fluoro-, chloro-, bromo- or iodo-), a carboxylate group, a 4 to 7 membered heterocyclic ring optionally containing 1 or more N, S, O or P atoms, a 5 or 6 membered aromatic ring, optionally containing 1 or more N, O or S atoms which can be fused to another ring or a said alkyl chain substituted by a said aromatic ring, or a $C(=O)XR$ group as defined below,

X is O, NH, NR, where R is H, OH, a branched or straight chain alkyl of from 1 to 6 C atoms which can be functionalised, alkoxy containing a branched or straight chain alkyl of from 1 to 6 C atoms which can be functionalised, a 4 to 7 membered heterocyclic ring optionally containing 1 or 2 N, S, O or P atoms, a 5 or 6 membered aromatic ring, optionally containing 1 or 2 N, O or S atoms which can be fused to another ring, such that RX is typically straight or branched C_1 to C_6 alkoxy, and m is 0 or 1.

XIX:



where R^I is as defined in EP0114031: i.e. C_1 to C_4 alkyl chain, such as C_2H_5 , which may be substituted with an alkoxy group with a C_1 to C_4 alkyl chain such as methyl


65

and the CONHR₁ groups are typically in the 2 and 4 positions. Thus R' is typically methoxyethyl.

Compounds of formula XVII and XVIII are exemplified by 2,5-(C8), 2,4-(C9), 2,3-(C10) and 3,4-(C11)-pyridine dicarboxylic acids, and compounds of the
5 formula XIX are exemplified by the compounds IS4, IS5, IS6, IS7, IS8 and IS9 in Table 3.


Another suitable N containing heterocycle may have the formula;

XX:



10


XXI: Pyridylcarbonyl glycines and derivatives



15

where X = O, Y = N or CR₃, m = 0 or 1, A = substituted alkylene, B=CO₂H, NHSO₂CF₃, tetrazolyl, imidazolyl or 3-hydroxyisoxazolyl, R₁, R₂ and R₃ may independently be H, OH, halo, cyano, CF₃, NO₂, CO₂H, alkyl, cycloalkyl, cycloalkoxy, aryl, aralkynyl, alkynylcarbonyl, alkylcarbonyloxy, carbamoyl, alkynyloxyalkyl, alkenyloxy, alkoxyalkoxy, alkynyl, retinyloxycarbonyl, alkenyloxycarbonyloxy, where R¹ and R² or R² and R³ = (CH₂)_o in which 1-2 CH₂ groups of the saturated or C:C unsaturated chain may be replaced by O, S, SO, SO₂ or imino, o= 3-5, R₄ = H.

XXII: 3-alkoxypyridine-2-carboxamide derivatives



where A, B and R⁴ as defined in WO97/41103: A=(substituted alkylene), B=(modified) carboxy, tetrazolyl, imidazolyl, 3-hydroxyisoxazolyl, R₄ = H, OH, halo, cyano, CF₃, NO₂, CO₂H, alkyl (e.g. branched or straight chain C₁ -C₆ alkyl), cycloalkyl, cycloalkylalkyl, cycloalkylalkoxy, cycloalkoxyalkyl, aryl, aralkyl, aralkoxy, hydroxyalkyl, alkenyl, alkynyl, alkynyloxyalkyl, alkoxycarbonyl, alkylcarbonyloxy, arylcarbonyloxy, cinnamoyl, alkenylcarbonyl, arylcarbamoyl or aralkoxycarbonyloxy.


Various - containing heterocycles and derivatives are known to be inhibitors of CPH. Their mechanism of action is believed to be via bidentate chelation of the aromatic nitrogen (or its N-oxide) and a carbonyl group located at the 2-position on the heterocyclic ring. Suitable compounds are described in Bickel *et al* Hepatology 28(2) 404-411, DE-A-1974628, EP-A-0846685, WO-A-9741103, EP07865871, DE-A-19504226, EP-A-0673932, EP-A-0673931 and EP-A-0673930.

Analogues of these compounds, appropriately derivatised for pharmaceutical use, may be made selective for HIF hydroxylases, and in particular for HPH activity, using the methods described herein.

2,4 -diethylpyridine dicarboxylate is a known CPH inhibitor (Friedman L. et al (2000) PNAS 97 (9) 4736-4741) which was found not to inhibit HPH. This provides indication that selective HIF hydroxylase inhibitors, and in particular selective HPH inhibitors, are possible.


5 Another class of compounds of interest for use in the modulation of HIF hydroxylases, and in particular of HPH activity, have the general formulae (XXV-XXVIII, XXVIII A and XXVIII B):

XXV:




15

XXVI:



XXVII (M6635):

20




where R, R^I to R^{VI} may independently be H, a branched or straight C₁ to C₆ alkyl

chain, a 4 to 7 membered heterocyclic ring optionally containing 1 or 2 N, S, O or P atoms, a 5 or 6 membered aromatic ring, optionally containing 1 or 2 N, O or S atoms, which can be fused to another ring, or a said alkyl chain substituted by a said aromatic ring, preferably H or methyl, R²⁰ is hydrogen or acyl typically aromatic acyl such as benzoyl.

X is NH, NR'', where R'' is OH, Me, alkyl, OMe, Oalkyl with a C₁ to C₆ alkyl chain, and


Y is O or S.

10 XXVIII:



such as

XXVIII A:



15

where A*B*C*D* are as defined for formula (A), R²¹ is hydrogen or acyl, typically aromatic acyl such as benzoyl, R is as defined for formula III, n is from 1 to 5, x is from 1 to 5 and y is from 1 to 5, such that the resulting methylene chains can be functionalised by one or more groups as defined for R or by NH₂, such as glutathione (gamma-glutamyl-cysteinyl-glycine) (C16) and cysteinyl-glycine (CO166).

20


These compounds may be obtained from commercial sources (for example, Sigma Chemical Co.) or prepared using standard methodology.

N-(Mercaptopropionyl)glycine and glutathione have been found to be inhibitors of HPH. This provides indication that appropriately functionalised thiols may be inhibitors of 2OG dependent oxygenases. N-(Mercaptopropionyl)glycine was not an inhibitor of clavamate synthase from *Streptomyces clavuligerus* under standard assay conditions, demonstrating that this family of compounds can be selective for different 2OG oxygenases. Modification of N-(mercaptpropionyl)glycine may improve selectivity for HPH. These compounds may be made into useful pharmaceutical agents by conversion into their ester 'pro drug' forms. These are commonly methyl or ethyl esters although others are possible.


N-(Mercaptopropionyl)glycine and glutathione are structurally related to L- δ (α -aminoadipoyl)-L-cysteinyl-D-valine (ACV) which is the substrate of isopenicillin-N-synthase (IPNS). IPNS is an oxidase which is closely related to the 2OG dependent oxygenases by sequence and structure, although it does not use a 2OG co-substrate. Glutathione is important in maintaining the correct redox potential inside cells and the observation that it is an inhibitor of HPH may have physiological relevance.

Compounds of Formula XXVIII include N-(3-mercaptopropanoyl)-L-alanine (IS37) and the corresponding D isomer (IS38) as well as N-(3-benzoylthiopropionoyl)-L-alanine (IS20) and the corresponding D isomer (IS21). Other compounds include:

XXIXA:



XXIX:



where R' is H, a branched or straight C₁ to C₆ alkyl chain which can be functionalised, any natural amino acid side chain for example of glutamic acid, a 4 to 7 membered heterocyclic ring optionally containing 1 or 2 N, S, O or P atoms or a 5 or 6 membered aromatic ring, optionally containing 1 or 2 N, O or S atoms which
 5 may be fused to another ring or a said alkyl chain substituted by a said aromatic ring and each of R² to R⁶, which may be the same or different, is as defined for R' or is NH₂ or OR⁷ where R⁷ is as defined for R' and E represents a monocyclic ring system such as thiophene or pyran and E' is absent or forms with E a bicyclic ring system such as naphthalene or indole, E' typically being benzene. The ring or rings can be
 10 functionalised at any of their carbon atoms with a group as defined for R'. Typical compounds include 2-hydroxy-hippuric acid, N(2-hydroxy-benzoyl)-glycine (C14) and N-benzoyl-glutamic acid (C15).

Where appropriate the acids can be in the form of salts, eg. sodium salts.

Peptide fragments derived from the sequence of a HIF hydroxylase or an
 15 HIF-α polypeptide form another class of compounds which may have modulating activity. Nucleic acid encoding such peptides, vectors and host cells containing such nucleic acid, and methods of expressing nucleic acid encoding such peptides are further aspects of the present invention. In a preferred embodiment such fragments are fragments of human HIF hydroxylases and in particular of any of PHD 1, 2 or 3.

20 A suitable modulator may be an analogue of HIFα, the prime substrate for HIF prolyl hydroxylation by HIF hydroxylase, and may act, for example, as a competitive inhibitor of HIFα, or through another mechanism, such as by irreversibly

modifying the HIF hydroxylase. Uncoupled oxidation of the co-substrate 2OG may still occur in the event of competitive inhibition of HIF- α .


A suitable fragment of an HIF α polypeptide may comprise a proline residue which is hydroxylated by a HIF hydroxylase in the full-length polypeptide, and which is, itself, capable of being hydroxylated by a HIF hydroxylase, for example a fragment containing a proline residue which corresponds to proline residue 402 or 564 of the human HIF-1 α sequence. Smaller fragments, and analogues and variants of these fragments may similarly be employed, e.g. as identified using techniques such as deletion analysis or alanine scanning.

Knowledge of the HIF α sequence, in particular the identity of the residues which are hydroxylated, therefore allows an inhibitor to be designed with a proline analogue in the position of hydroxylation to inhibit the hydroxylation reaction.


In particular, a modulator, such as an inhibitor, may include a peptide fragment of HIF α or analogue thereof in which the proline residue which undergoes hydroxylation, e.g. proline 564, is replaced by a proline analogue which is not a HIF hydroxylase substrate, such as 5-oxaproline, 3, 4-dehydroproline and 4-thiaproline (Wu *et al* (1999) J. Am. Chem. Soc. 121(3)587-588, DE-A-3818850). The proline inhibitor analogues may be modified such that they also bind to the 2-oxoglutarate binding residues of the hydroxylase.

Examples of proline analogues are;

XXIII:



XXIV:



where one of Y or X = O (oxaprolines) and the other is -CR'R''- or Y=S (4-thiaprolines) where one of X and Y is C=O or -SO₂ or -P(=O)O(H) and the other is -CR'R''-,

and R is a peptide or peptide analogue and R, R^I, R^{II} are H, a branched or straight C₁ to C₆ alkyl chain, a 4 to 7 membered heterocyclic ring optionally containing 1 or more F, N, S, O or P atoms, a 5 or 6 membered aromatic ring, optionally containing 1 or more N, O or S atoms, which can be fused to another ring.

5 Peptides or peptide analogues suitable for use in accordance with the present invention tend to be short, and may be about 40 amino acids in length or less, preferably about 35 amino acids in length or less, more preferably about 30 amino acids in length, or less, more preferably about 25 amino acids or less, more preferably about 20 amino acids or less, more preferably about 15 amino acids or
10 less, more preferably about 10 amino acids or less, or 9, 8, 7, 6, 5 or less in length. Peptides according to the present invention may be about 10-40 amino acids in length, about 5-10, about 10-15, about 10-20, about 10-30, about 20-30, or about 30-40 amino acids in length. Peptides which are HIF α fragments generally include one or more of the relevant proline residues.

15 A peptide modulator which is a derivative of a peptide for which the specific sequence is disclosed herein may be in certain embodiments the same length or shorter than the specific peptide. In other embodiments the peptide sequence or a variant thereof may be included in a larger peptide, as discussed above, which may or may not include an additional portion of HIF hydroxylase or HIF-1 polypeptide. 1,
20 2, 3, 4 or 5 or more additional amino acids, adjacent to the relevant specific peptide fragment of the HIF hydroxylase or HIF-1 polypeptide, or heterologous thereto may be included at one end or both ends of the peptide.

 Peptides may be modified, for example, for use as pharmaceuticals, for example by replacing amide bonds and/or using D rather than L α -amino acids or β -
25 amino acids or by using conformational restraints.

 Examples of potential HIF hydroxylase inhibitors, and in particular of HPH activity, of the classes described above are shown in Table 3.

 In the formulae described herein, a branched or straight C₁ to C₆ alkyl chain may be a methyl, ethyl, propyl, butyl, iso-butyl, *sec*-butyl, *tert*-butyl, pentyl,
30 neopentyl *tert*-pentyl or a primary, secondary or tertiary hexyl group. Preferably the alkyl group is methyl or ethyl while the preferred heterocyclic rings are pyrrolidine or

tetrahydropyran and the aromatic rings are benzene, naphthalene or pyridine.

Without being limited to any particular mechanism, analysis of the structure of IPNS complexed to Fe(II) and ACV, together with the structural relationship between N-(mercaptopropionyl) glycine, 2OG, and N-oxaloglycine, provides
5 indication that the former inhibits HIF hydroxylase and in particular HPH activity via a complex in its thiol binds to the Fe(II) and the carboxylate to the same residues as the S-carboxylate of 2OG.

HPH Selectivity

A number of distinct HIF hydroxylases with HIF prolyl hydroxylase activity
10 exist in humans and it may be also be advantageous to modulate these selectively; as single targets, or in selected groups as well as an entire family. Agents which modulate HIF hydroxylase activity and in particular HIF prolyl hydroxylase activity are therefore preferably specific i.e. they have an increased or enhanced effect on a HIF hydroxylase relative to other 2OG dependent oxygenases as defined below, in
15 particular collagen prolyl hydroxylases (CPH). Such agents may be specific for a particular human HIF hydroxylases and in particular PHD 1, 2 or 3.

Assay methods as described herein may therefore further comprise contacting the test compound with one or more 2OG dependent oxygenases under conditions in which said 2OG dependent oxygenases are normally active and determining activity
20 of said oxygenases.

A difference in activity in the presence relative to the absence of test compound is indicative of the test compound modulating the activity of the one or more 2OG dependent oxygenases.

A test compound which provides increased or enhanced modulation of a HIF
25 hydroxylase, relative to the one or more 2OG dependent oxygenases shows selectivity or specificity for the HIF hydroxylase.

2OG dependent oxygenases may include for example, clavamint synthase, deacetoxycephalosporin C synthase, collagen-prolyl-4-hydroxylase, collagen prolyl-3-hydroxylase, lysyl hydroxylase, aspartyl hydroxylase, phytanoyl coenzyme A
30 hydroxylase or gamma-butyrobetaine hydroxylase. 2OG dependent oxygenases may be mammalian, preferably human polypeptides.

The structures of various 2-OG oxygenase enzymes have been reported; oxygenase clavaminic acid synthase (Zhang Z. *et al* (2000) Nature Structural Biol. 7 127-133), deacetoxycephalosporin C synthase (Lloyd *et al* (1999) J. Mol. Biol. 287 943-960), cephalosporin synthase (Valégard K. *et al* (1998) Nature 394 805-809),
5 isopenicillin N-synthase (Roach P. (1995) Nature 375 700-704) and 2OG dependent oxygenases (Schofield, C. & Zhang, Z. (1999) Curr. Opin. Struct. Biol. 9 722-731).

The assays of the invention may also be used to identify agents which are selective for HIF hydroxylases, and in particular HPH activity, but not for other 2OG dependent oxygenases; agents which are selective for HIF hydroxylase, and in
10 particular HIF prolyl hydroxylase activity, compared to other hydroxylases and prolyl hydroxylases; and agents which are specific for a particular HIF hydroxylase. The assays may be used to identify agents selective for a particular human HIF hydroxylase, such as for example specific for an enzyme with the amino acid sequence of SEQ ID NO: 2, or SEQ ID NO: 4 or SEQ ID NO: 6.

15 The invention provides for the use of such selective inhibitors of HIF hydroxylases in the manufacture of a medicament for the treatment of a condition associated with reduced HIF levels of activity.

In alternative aspects of the present invention, the assays can be used to establish whether agents which have been identified as inhibitors or activators of
20 other 2OG dependent oxygenases are specific for such oxygenases, or at least do not affect HIF hydroxylase and in particular HIF prolyl hydroxylase activity of the polypeptides of the present invention. In particular, the assays may be used to establish that such agents do not affect human HIF hydroxylases. Thus, the assays may be carried out using agents which have been identified as inhibitors of a 2OG
25 dependent oxygenase, such as collagen prolyl hydroxylase to identify whether such an agent is specific for collagen prolyl hydroxylase and is not active or shows reduced activity against HIF hydroxylases and in particular their prolyl hydroxylase activity.

Assay Formats

A screening or assay method may include purifying and/or isolating a test compound, agent, or substance of interest from a mixture or extract, i.e. reducing the content of at least one component of the mixture or extract, e.g. a component with
5 which the test substance is naturally associated. The screening or assay method may include determining the ability of one or more fractions of a test mixture or extract to modulate the hydroxylase and in particular the prolyl activity of the HIF hydroxylase, and typically these activities in relation to HIF- α .

The purification and/or isolation may employ any method known to those
10 skilled in the art. An agent or compound obtained and/or identified using an assay method described herein may be modified, for example to increase selectivity for a HIF hydroxylase relative to other 2OG dependent oxygenases such as CPH or to increase selectivity for a particular HIF hydroxylase relative to other HIF hydroxylases.

15 The approach of modifying a class of compounds containing specific functional groups to be selective for particular enzymes is well-known, for example the inhibition of specific serine proteases by differently modified trifluoroketones, chloromethylketones, beta lactams, or other generic serine protease inhibitors (Walker B. & Lynas J. (2001) Cell. Mol. Life Sci. 58(4) 596-624, Rai R. (2001) *et al*
20 Curr. Med. Chem. 8 (2) 101-119, Marquis R. (2000) Ann. Rep. Med. Chem. 35 309-320, Lebon, F. & Ledecq, M. (2000) Curr. Med. Chem. 7 (4) 455-477).

Selectivity for a particular HIF hydroxylase can be achieved by performing assays as described herein with a plurality of different HIF hydroxylases. The preferential or selective modulation of the HIF hydroxylase activity, and in particular
25 the prolyl hydroxylase activity, of one or more HIF hydroxylases relative to the other HIF hydroxylase by a test compound may thereby be determined. Similarly, selectivity for the PHD family of HIF hydroxylases can be achieved by performing assays as described herein with a plurality of different 2-oxoglutarate dependent oxygenases i.e. a panel of related enzymes. The preferential or selective modulation
30 of the HIF prolyl hydroxylase activity of one or more PHD polypeptides relative to

the activity of other 2-oxoglutarate dependent oxygenases by a test compound may thereby be determined.

Structural information, including primary sequence data and 3D information such as crystallographic data, may also be used to identify structural differences between HIF hydroxylases and other 2-oxoglutarate dependent oxygenases. These differences may be used to design compounds which selectively or preferentially modulate HIF hydroxylases as described herein relative to other 2-oxoglutarate dependent oxygenases.

Structural Analysis and Rational Drug Design

Secondary structural analysis predicts that the HIF hydroxylases fold to produce a common jelly roll structure that positions a non-haem iron co-ordinating $\text{HXD}[\text{X}]_n\text{H}$ motif at the catalytic site.

Kinetic and time resolved crystallographic studies of the catalytic mechanism among members of this class of oxygenase have indicated ordered binding of Iron (II), 2-oxoglutarate, and prime substrate (Zhang *et al.*, (2000) *supra*; Zhou *et al* (1998) J. Am. Chem. Soc. 120 13539-13540).

Binding of the latter 'primes' the enzyme for reversible binding of dioxygen, probably by displacing a water molecule from the iron. Weak binding at the iron centre is associated with a cofactor requirement for iron (II). The activity of the recombinant enzyme requires iron, and is directly inhibited by cobaltous ions by substitution at the catalytic center. A mutation (EGLN2/PHD1; H358A) that is predicted to abrogate iron binding, but not otherwise alter the 3-dimensional structure was observed to completely ablate enzyme activity,

Structural analysis and sequence studies show that SM20 comprises an iron atom at its active site in complex with residues His313 and Asp315 (Acc No: af229245 or gi11320937). Related enzymes clavamate synthase (CAS) and deacetoxycephalosporin C synthase (DAOCS) also show iron coordination at the active site as shown in Figure 8.

An inhibitor may form a mono-, di-, or tri-dentate complex with the coordinate iron atom to inhibit the activity of the enzyme. Examples of such inhibitors include the hydroxamates and hydroxyanthroquinones described herein.

The coordination of an inhibitor to the iron atom may be determined by spectroscopic analysis or crystallography.

As noted above, an agent may be peptidyl, e.g. a peptide which includes a sequence as recited above, or may be a functional analogue of such a peptide.

5 As used herein, the expression "functional analogue" relates to peptide variants or organic compounds having the same functional activity as the peptide in question, which may interfere with the hydroxylation and in particular the prolyl hydroxylation of HIF α by a HIF hydroxylase. Examples of such analogues include chemical compounds which are modelled to resemble the three dimensional structure
10 of the substrate of the HIF hydroxylase (HIF α) in the contact area or in the pVHL binding domain, and in particular the arrangement of the key amino acid residues, including proline 564.

In a further aspect, the present invention provides the use of a HIF α polypeptide, in particular a peptide fragment which undergoes hydroxylation by a
15 HIF hydroxylase, in a method of designing a peptide or non-peptidyl mimetic, which mimetic is able to interact with the HIF hydroxylase active site and modulate the hydroxylation of a proline residue i.e. proline 402 or 564 of HIF α , by the HIF hydroxylase.

Accordingly, the present invention provides a method of designing a mimetic,
20 for example of a HIF α polypeptide, which modulates the hydroxylation of a proline residue by a HIF hydroxylase, said method comprising:

- (i) analysing a substance to determine the amino acid residues essential and important for biological activity to define a pharmacophore; and,
- (ii) modelling the pharmacophore to design and/or screen candidate
25 mimetics which modulate the hydroxylation as described.

Suitable modelling techniques are known in the art. This includes the study of the bonding between a HIF hydroxylase and HIF- α and the design of compounds which contain corresponding functional groups arranged in such a manner that they could reproduce that bonding.

30 The iron atom at the HIF hydroxylase active site is normally hexacoordinate (but can be pentacoordinate during catalysis). The amino acid sequence provides

three ligands so there are three positions vacant for binding to inhibitors. Inhibitor molecules may therefore be designed to coordinate with the iron atom at these three positions.

The designing of mimetics to a known pharmaceutically active compound is a known approach to the development of pharmaceuticals based on a "lead" compound, for example a compound as described herein. This might be desirable where the active compound is difficult or expensive to synthesise or where it is unsuitable for a particular method of administration, for instance, HIF- α or HIF hydroxylase and in particular peptides derived from them may not be well suited as active agents for oral compositions as they tend to be quickly degraded by proteases in the alimentary canal.

There are several steps commonly taken in the design of a mimetic from a compound having a given target property. Firstly, the particular parts of the compound that are critical and/or important in determining the target property are determined. In the case of a peptide, this can be done by systematically varying the amino acid residues in the peptide, e.g. by substituting each residue in turn. These parts or residues constituting the active region of the compound are known as its "pharmacophore".

Once the pharmacophore has been found, its structure is modelled according to its physical properties, e.g. stereochemistry, bonding, size and/or charge, using data from a range of sources, e.g. spectroscopic techniques, X-ray diffraction data and NMR. Computational analysis, similarity mapping (which models the charge and/or volume of a pharmacophore, rather than the bonding between atoms) and other techniques can be used in this modelling process.

In a variant of the above approach, the three-dimensional structure of a ligand and its binding partner are modelled. This can be especially useful where the ligand and/or binding partner change conformation on binding, allowing the model to take account of this in the design of the mimetic.

Polypeptide inhibitors of the invention.

The invention also provides a polypeptide of SEQ ID NO: 9
DLDLEMLAP*YIPMDDDFQL wherein P* is 4-hydroxy proline. Such peptides

may be provided in an isolated form. The invention also provides variants of SEQ ID NO:9, as defined above. Variants will retain the ability to antagonise the interaction of a HIF- α subunit with VHL. This means that the presence of the variant in a cell will lead to an increase in HIF- α subunit protein compared to the amount of

5 HIF- α subunit protein present in the absence of the polypeptide.

Particular examples of such substitutions include the following:

DLDLEMLAP*YISMDDDFQL SEQ ID NO:10;
DLDLEMLLP*YIPMDDDFQL; SEQ ID NO:11
DLDLEMLVP*YIPMDDDFQL; SEQ ID NO:12
10 DLDLEMIAP*YIPMDDDFQL; SEQ ID NO:13
DLDLEMIAP*YIPMEDDFQL; SEQ ID NO:14, and
DLDLEMLVP*YISMDDDFQL SEQ ID NO:15.

The invention also provides an isolated polypeptide which is a variant of SEQ ID NO: 9, wherein such variants comprise from 1 to 4 amino acid substitutions of

15 any amino acid apart from P*, said variant retaining the ability to antagonise the interaction of a HIF- α subunit with VHL.

A particular polypeptide of the latter type is:

PFSTQDLDLEMLAPYIPMDDDFQLRSFDQLSP (SEQ ID NO:16);
or variants thereof as defined for SEQ ID NO:9 above; and

20 polypeptides consisting of from 35 to 50 amino acids which contain SEQ ID NO:16.

Fragments of these polypeptides which retain the P* residue, and which are at least 6, preferably at least 10, such as at least 12 or at least 15 amino acids in length are also a further aspect of the invention, and are also referred to herein as

25 polypeptides of the invention. Preferably these fragments retain the motif LXP*, e.g. LAP*, and more preferably the fragments retain the motif LXXLXP*, e.g. LXXLAP*. "X" means any amino acid. In particular, such a polypeptide has or includes the sequence LAP*YIP. Thus the invention also relates to a peptide comprising or having the sequence LAP*YIP and having the ability to antagonise the

30 interaction between HIF and VHL. Such peptides may be formulated or used as described for peptides of SEQ ID NO:9.

We have also found that a second proline residue in HIF-1 α is subject to hydroxylation. Our findings indicate that residue 402 which shares a common motif of "LXXLAP" with the site of hydroxylation at position 564. Our findings indicate that this site. Polypeptides of at least 8, e.g. at least 10, at least 12, at least 15 or at
5 least 18 amino acids, up to no more than 50, such as no more than 35 or no more than 20 amino acids, based upon the sequences of this region also form a further aspect of the invention. Such polypeptides include those in which proline at 402 is hydroxylated. Substitutions, modifications, purification, isolation and/or synthesis may be carried out as described for HIF hydroxylases described above.

10 The hydroxylated peptides in accordance with the present invention can be used in assays to monitor for agents which inhibit the interaction between VHL and HIF. In accordance with this aspect of the invention, a hydroxylated polypeptide as described above is incubated with VHL or a HIF binding region thereof in the presence of a test substance and the interaction between the hydroxylated polypeptide
15 and the VHL polypeptide is monitored. The polypeptides may also be used as displacement probes in high throughput assays for inhibitors.

These polypeptides of the present invention may be prepared as a pharmaceutical preparation. Such preparations will comprise the polypeptide together with suitable carriers, diluents and excipients. Typically, they will comprise
20 the polypeptide together with a pharmaceutically acceptable polypeptide. Such formulations form a further aspect of the present invention.

Formulations may be prepared suitable for any desired route of administration, including oral, buccal, topical, intramuscular, intravenous, subcutaneous and the like.

25 Formulations for topical administration to the skin may include ingredients which enhance the permeability of the skin to the polypeptides. Such formulations may be in the form of ointments, creams, transdermal patches and the like.

Formulations for administration by injection (i.m., i.v., subcutaneous and the like) will include sterile carriers such as physiological saline, optionally together with
30 agents which preserve or stabilise the polypeptide. Albumin may be a suitable agent.

Formulations of polypeptides in particular may be used in methods of treatment ischaemic conditions, such as organ ischaemia, such as is manifest in coronary, cerebrovascular and peripheral vascular insufficiency. Any ischaemia is a therapeutic target. The therapy may be applied in two ways; following declared
5 tissue damage, e.g. myocardial infarction (in order to limit tissue damage), or prophylactically to prevent ischaemia, e.g. promotion of coronary collaterals in the treatment of angina. Additionally, vasomotor control is subject to regulation by HIF. Activation of HIF might affect systemic vascular resistance and hence systemic blood pressure.

10 Polypeptides may also be used in combination with promoters of angiogenesis. These include vascular endothelial growth factor and other angiogenic growth factors such as basic fibroblast growth factors and thymidine phosphorylase and pro-angiogenic and might be used in combination therapy. Other compounds which might conceivably be used in combination are 2-deoxy ribose and
15 prostaglandin E.

In administering polypeptides of the invention to a subject, the doses will be determined at the discretion of the physician, taking into account the needs of the patient and condition to be treated. Generally, doses will be provided to achieve concentrations at a desired site of action that are from 0.1 μ M to 1mM, for example in
20 the 1-10 μ M range.

Therapeutic Applications

A compound, substance or agent which is found to have the ability to affect the hydroxylase activity of a HIF hydroxylase, and in particular its prolyl hydroxylase activity, has therapeutic and other potential in a number of contexts, as
25 discussed. For therapeutic treatment, such a compound may be used in combination with any other active substance, e.g. for anti-tumour therapy another anti-tumour compound or therapy, such as radiotherapy or chemotherapy.

An agent identified using one or more primary screens (e.g. in a cell-free system) as having ability to modulate the HIF α hydroxylation activity of a HIF
30 hydroxylase may be assessed further using one or more secondary screens. A secondary screen may involve testing for an increase or decrease in the amount of

HIF- α or HIF activity, for example as manifest by the level of a HIF target gene or process present in a cell in the presence of the agent relative to the absence of the agent.

A HIF hydroxylase or a HIF polypeptide may be used in therapies which
5 include treatment with full length polypeptides or fragments thereof, or otherwise modified polypeptides (e.g. to enhance stability or ensure targeting, including in conjunction with other active agents such as antibodies.

Generally, an agent, compound or substance which is a modulator according to the present invention is provided in an isolated and/or purified form, i.e.
10 substantially pure. This may include being in a composition where it represents at least about 90% active ingredient, more preferably at least about 95%, more preferably at least about 98%. Any such composition may, however, include inert carrier materials or other pharmaceutically and physiologically acceptable excipients, such as those required for correct delivery, release and/or stabilisation of the active
15 agent. As noted below, a composition according to the present invention may include in addition to an modulator compound as disclosed, one or more other molecules of therapeutic use, such as an anti-tumour agent.

Products obtained by assays of the invention

The invention further provides compounds obtained by assay methods of the
20 present invention, and compositions comprising said compounds, such as pharmaceutical compositions wherein the compound is in a mixture with a pharmaceutically acceptable carrier or diluent. The carrier may be liquid, e.g. saline, ethanol, glycerol and mixtures thereof, or solid, e.g. in the form of a tablet, or in a semi-solid form such as a gel formulated as a depot formulation or in a transdermally
25 administerable vehicle, such as a transdermal patch.

The invention further provides a method of treatment which includes administering to a patient an agent which interferes with the hydroxylation of the target residue of an HIF α polypeptide by a HIF hydroxylase. Such agents may include inhibitors of hydroxylase activity, typically of prolyl hydroxylase activity and
30 in particular these activities in relation to HIF. Examples of inhibitors of HIF prolyl hydroxylase activity include, for example compounds of structures I to XXVIII as

described herein. Exemplary purposes of such treatment are discussed elsewhere herein.

The invention further provides various therapeutic methods and uses of one or more substances selected from (i) a HIF hydroxylase which is able to bind to HIF-1; 5 (ii) a modulator identified by a screening method of the present invention; (iii) a mimetic of any of the above substances which can bind to HIF-1 or a HIF hydroxylase, or the polypeptide inhibitors of the invention.

The therapeutic/prophylactic purpose of such a method or use may be the modulation of the level of HIF α in a cell by modulation, e.g. disruption or 10 interference, of the hydroxylation of HIF α , which may occur for example at proline 402, 564 or other proline residue. Hydroxylation of HIF α promotes pVHL binding which leads to ubiquitin dependent proteolysis of HIF α as described above.

The therapeutic/prophylactic purpose may be related to the treatment of a condition associated with reduced or suboptimal or increased HIF levels or activity, 15 or conditions in which have normal HIF levels, but where an modulation in HIF levels such as an increase or decrease in HIF levels is desirable such as:

- (i) ischaemic conditions, for example organ ischaemia, including coronary, cerebrovascular and peripheral vascular insufficiency. The therapy may be applied in two ways; following declared tissue damage, e.g. myocardial infarction (in order to 20 limit tissue damage), or prophylactically to prevent ischaemia, e.g. promotion of coronary collaterals in the treatment of angina.
- (ii) wound healing and organ regeneration
- (iii) auto-, allo-, and xeno- transplantation.
- (iv) systemic blood pressure
- 25 (v) cancer; HIF α is commonly up-regulated in tumour cells and has major effects on tumour growth and angiogenesis.
- (vi) inflammatory disorders.
- (vii) pulmonary arterial blood pressure, neurodegenerative disease.

Pharmaceutical Compositions

In various further aspects, the present invention thus provides a pharmaceutical composition, medicament, drug or other composition for such a purpose, the composition comprising one or more agents, compounds or substances as described herein, including HIF hydroxylase inhibitors and in particular inhibitors of their HIF prolyl hydroxylase (HPH) activity such as compounds of formulae I to XXVIII, the use of such an composition in a method of medical treatment, a method comprising administration of such a composition to a patient, e.g. for treatment(which may include preventative treatment) of a medical condition as described above, use of such an agent compound or substance in the manufacture of a composition, medicament or drug for administration for any such purpose, e.g. for treatment of a condition as described herein, and a method of making a pharmaceutical composition comprising admixing such an agent, compound or substance with a pharmaceutically acceptable excipient, vehicle or carrier, and optionally other ingredients.

In one embodiment the method for providing a pharmaceutical composition may typically comprise:

- (a) identifying an agent by an assay method of the invention; and
- (b) formulating the agent thus identified with a pharmaceutically acceptable excipient.

The pharmaceutical compositions of the invention may comprise an agent, polypeptide, polynucleotide, vector or antibody according to the invention and a pharmaceutically acceptable excipient.

The agent may be used as sole active agent or in combination with one another or with any other active substance, e.g. for anti-tumour therapy another anti-tumour compound or therapy, such as radiotherapy or chemotherapy.

Whatever the agent used in a method of medical treatment of the present invention, administration is preferably in a "prophylactically effective amount" or a "therapeutically effective amount" (as the case may be, although prophylaxis may be considered therapy), this being sufficient to show benefit to the individual. The actual amount administered, and rate and time-course of administration, will depend

Nucleic acid encoding a HIF hydroxylase as described herein may also be used in the anti-sense regulation of the HIF hydroxylase activity and in particular, of the HIF prolyl hydroxylase activity within a cell.

Down-regulation of expression of a gene encoding a HIF hydroxylase may be achieved using anti-sense technology, or RNA interference.

In using anti-sense genes or partial gene sequences to down-regulate gene expression, a nucleotide sequence is placed under the control of a promoter in a "reverse orientation" such that transcription yields RNA which is complementary to normal mRNA transcribed from the "sense" strand of the target gene. See, for example, Smith *et al.*, (1988) *Nature* 334, 724-726;. Antisense technology is also reviewed in Flavell, (1994) *PNAS USA* 91, 3490-3496.

The complete sequence corresponding to the reverse orientation of the coding sequence need not be used. For example, fragments of sufficient length may be used. It is a routine matter for the person skilled in the art to screen fragments of various sizes and from various parts of the coding sequence to optimise the level of anti-sense inhibition. It may be advantageous to include the initiating methionine ATG codon, and perhaps one or more nucleotides upstream of the initiating codon. A further possibility is to target a conserved sequence of a gene, e.g. a sequence that is characteristic of one or more genes, such as a regulatory sequence.

The sequence employed may be 500 nucleotides or less, possibly about 400 nucleotides, about 300 nucleotides, about 200 nucleotides, or about 100 nucleotides. It may be possible to use oligonucleotides of much shorter lengths, 14-23 nucleotides, although longer fragments, and generally even longer than 500 nucleotides are preferable where possible.

Anti-sense oligonucleotides may be designed to hybridise to the complementary sequence of nucleic acid, pre-mRNA or mature mRNA, interfering with the production of a HIF hydroxylase encoded by a given DNA sequence (e.g. either native polypeptide or a mutant form thereof), so that its expression is reduced or prevented altogether. Anti-sense techniques may be used to target a coding sequence, a control sequence of a gene, e.g. in the 5' flanking sequence, whereby the anti-sense oligonucleotides can interfere with control sequences. Anti-sense

oligonucleotides may be DNA or RNA and may be of around 14-23 nucleotides, particularly around 15-18 nucleotides, in length. The construction of antisense sequences and their use is described in Peyman and Ulman, Chemical Reviews, 90:543-584, (1990), and Crooke, Ann. Rev. Pharmacol. Toxicol., 32:329-376, 5 (1992).

It may be preferable that there is complete sequence identity in the sequence used for down-regulation of expression of a target sequence, and the target sequence, though total complementarity or similarity of sequence is not essential. One or more nucleotides may differ in the sequence used from the target gene. Thus, a sequence 10 employed in a down-regulation of gene expression in accordance with the present invention may be a wild-type sequence (e.g. gene) selected from those available, or a mutant, derivative, variant or allele, by way of insertion, addition, deletion or substitution of one or more nucleotides, of such a sequence.

The sequence need not include an open reading frame or specify an RNA that 15 would be translatable. It may be preferred for there to be sufficient homology for the respective sense RNA molecules to hybridise. There may be down regulation of gene expression even where there is about 5%, 10%, 15% or 20% or more mismatch between the sequence used and the target gene.

Other approaches to specific down-regulation of genes which may be used to 20 modulate HIF hydroxylase expression are well known, including the use of ribozymes designed to cleave specific nucleic acid sequences. Ribozymes are nucleic acid molecules, actually RNA, which specifically cleave single-stranded RNA, such as mRNA, at defined sequences, and their specificity can be engineered. Hammerhead ribozymes may be preferred because they recognise base sequences of 25 about 11-18 bases in length, and so have greater specificity than ribozymes of the Tetrahymena type which recognise sequences of about 4 bases in length, though the latter type of ribozymes are useful in certain circumstances. References on the use of ribozymes include Marschall, *et al.* Cellular and Molecular Neurobiology, 1994. 14(5): 523; Hasselhoff, Nature 334: 585 (1988) and Cech, J. Amer. Med. Assn., 260: 30 3030 (1988).

Vectors such as viral vectors have been used in the prior art to introduce nucleic acid into a wide variety of different target cells. Typically the vectors are exposed to the target cells so that transfection can take place in a sufficient proportion of the cells to provide a useful therapeutic or prophylactic effect from the expression of the desired peptide. The transfected nucleic acid may be permanently
5 incorporated into the genome of each of the targeted cells, providing long lasting effect, or alternatively the treatment may have to be repeated periodically.

A variety of vectors, both viral vectors and plasmid vectors, are known in the art, see US Patent No:5,252,479 and WO93/07282. In particular, a number of
10 viruses have been used as gene transfer vectors, including papovaviruses, such as SV40, vaccinia virus, herpesviruses, including HSV and EBV, and retroviruses. Many gene therapy protocols in the prior art have used disabled murine retroviruses.

As an alternative to the use of viral vectors in gene therapy other known methods of introducing nucleic acid into cells includes mechanical techniques such
15 as microinjection, transfer mediated by liposomes and receptor-mediated DNA transfer.

Receptor-mediated gene transfer, in which the nucleic acid is linked to a protein ligand via polylysine, with the ligand being specific for a receptor present on the surface of the target cells, is an example of a technique for specifically targeting
20 nucleic acid to particular cells.

In various further aspects, the present invention thus provides a pharmaceutical composition, medicament, drug or other composition for use in a method of treating a medical condition described above, the composition comprising an isolated nucleic acid molecule as described herein, the use of such an composition
25 in a method of medical treatment, a method comprising administration of such a composition to a patient, e.g. for treatment (which may include preventative treatment) of a medical condition as described above, use of such an agent compound or substance in the manufacture of a composition, medicament or drug for administration for any such purpose, e.g. for treatment of a condition as described
30 herein, and a method of making a pharmaceutical composition comprising admixing such an agent, compound or substance with a pharmaceutically acceptable excipient,

vehicle or carrier, and optionally other ingredients.

A peptide or other substance having an ability to modulate or interfere with the prolyl hydroxylation of the residue of HIF- α by a polypeptide, or a nucleic acid molecule which encodes a peptide having that ability, may be provided in a kit, e.g. sealed in a suitable container which protects its contents from the external environment. Such a kit may include instructions for use.

Use of polypeptides

Another aspect of the present invention provides the use of a HIF hydroxylase as described herein or a fragment thereof for the hydroxylation, and in particular the prolyl hydroxylation, of an HIF polypeptide, or a proline-containing substrate of HIF hydroxylation.

Another aspect of the present invention provides a method of producing a HIF hydroxylase comprising:

- (a) causing expression from nucleic acid which encodes a HIF hydroxylase in a suitable expression system to produce the polypeptide recombinantly;
- (b) determining the prolyl hydroxylation of an HIF α polypeptide by said recombinantly produced polypeptide. The polypeptide expressed by the method may be a PHD (EGLN) polypeptide.

Suitable expression systems are well-known in the art. HIF hydroxylases may be expressed in a prokaryote, such as *E. coli*, lower eukaryote such as *S. cerevisiae* or a higher eukaryotic cell, such as a mammalian cell e.g. a CHO or COS cell.

Prolyl hydroxylation of an HIF α polypeptide, in particular within the pVHL binding domain, such as residue 402 or 564, may be determined as described herein.

Another aspect of the present invention provides an assay method for identifying/obtaining a HIF hydroxylase, and in particular a HIF α prolyl hydroxylase, comprising,

- (a) providing a test polypeptide,
- (b) bringing into contact an HIF α polypeptide and the test polypeptide under conditions in which the HIF α polypeptide is hydroxylated by a HIF hydroxylase; and
- (c) determining hydroxylation and in particular the prolyl hydroxylation of the HIF α polypeptide.

A HIF hydroxylase polypeptide according to the present invention can also be used to identify additional substrates of HIF hydroxylases. For example, peptides which have either previously been demonstrated to be hydroxylated by other hydroxylases, or other peptides may be brought into contact with a HIF hydroxylase according to the present invention and monitoring for hydroxylation of such peptides. Any suitable conditions may be selected including the provision of agents and co-factors known to enhance hydroxylation by the hydroxylases of the present invention. In a preferred aspect, prolyl containing substrates are contacted with a HIF hydroxylase of the present invention, and hydroxylation of the prolyl residue is monitored. Hydroxylation of the substrate may be monitored by any suitable method including monitoring levels of co-factors or by products of hydroxylation.

The invention also provides a method of modulating the amount of HIF polypeptide in a cell comprising contacting the cell with a substance which inhibits the 4-prolyl hydroxylase activity of a HIF hydroxylase such as, for example, an EGLN polypeptide. The substance may, for example, be an agent of the invention. In a preferred embodiment, the substance inhibits the biological activity of HIF hydroxylase, such as, for example an EGLN polypeptide, but does not inhibit biological activity of a collagen prolyl hydroxylase.

Various further aspects and embodiments of the present invention will be apparent to those skilled in the art in view of the present disclosure.

Certain aspects and embodiments of the invention will now be illustrated by way of example and with reference to the figures described below.

Figures

Figure 1. Analysis of the minimal pVHL binding domain of HIF-1 α .
A. Left panel; sequence alignment of the minimal pVHL binding domain from HIF-1 α and HIF-2 α , with HIF- α genes from other organisms. B. Summary of the ability of synthetic polypeptides to block the HIF-1 α /pVHL interaction before and after exposure to reticulocyte lysate supplemented with Fe(II). Treated and untreated polypeptides were added to a mixture of HIF-1 α and pVHL.HA IVTTs that was then assayed for interaction by anti-HA immunoprecipitation. Substituted residues are underlined. Y* denotes phosphotyrosine.

Figure 2. Amino acid sequences of human HIF-1 α and *C. elegans* HIF.

Identical amino acids are boxed in black. The PAS domains are indicated. The VHL minimal binding region defined in studies of human HIF-1 α is indicated.

Figure 3A Line up of the indicated amino acids of the indicated HIF alpha chains showing conservation of the LxxLAP motif in all three domains demonstrated to be involved in VHL dependent ubiquitylation.

Figure 3B U2OS cells were transiently co-transfected with plasmids encoding the indicated Gal-Hif-1 alpha-VP16 sequences in combination with pUAS-tk-luc (a Gal 4 upstream activating sequence dependent luciferase reporter gene plasmid). Luciferase activities were determined in extracts made from transfected cells maintained for 48 hours, either entirely in normoxia (white bars) or with hypoxic stimulation for the last 16 hours. (black bars). Low normoxic activity (explicable by rapid destruction of the fusion protein) is seen when the Gal-Hif-VP16 fusion protein contains wild type Hif-1 alpha amino acids 344-417 but not when the sequence bears the P402A mutation or is truncated to amino acid 400.

Figure 4 The CHO mutant cell line Ka13 (deficient in Hif alpha subunit activity) was co-transfected with pcDNA3 expression plasmids encoding wild type full length Hif-1 alpha (HIF-1), full length Hif-1 alpha bearing the P402A mutation (P402A), full length Hif-1 alpha bearing the P564G mutation (P564G) or full length Hif-1 alpha bearing the double mutation (P402A+P564G) in combination with a hypoxia response element dependent luciferase reporter gene construct. Luciferase activities were determined in extracts made from transfected cells maintained for 48 hours, either entirely in normoxia (white bars) or with hypoxic stimulation for the last 16 hours (black bars). The individual mutations showed slightly enhanced normoxic activity compared with the wild type sequence but the combined mutant showed constitutive activity in normoxia with no further induction by hypoxia.

Figure 5 shows a schematic of the on-bead modification assay used to assay for HIF prolyl hydroxylase activity.

Figure 6 shows the effect of dimethyl oxalyl glycine on HIF activity in Hep3B and U2OS cell lines.

Figure 7 shows the effects of forced expression of EGLN2 (PHD 1) or a naturally occurring splice variant lacking amino acids 369-389 (PHD4) on the action of HIF in cells incubated in atmospheres containing 20% or 2% oxygen.

Figure 8 shows views derived from the active sites of clavamate synthase
5 and deacetoxycephalosporin C synthase.

Figure 9 shows sequence alignments of the predicted jelly roll cores of HIF-PHs. Sequences shown are *C. elegans* EGL-9 (462-580), Human EGLN1(288-403), Human EGLN2 (462-580), Human EGLN3 (111-225), rat SM20 (226-341). Also shown is the *Streptomyces* sp. Prolyl-3-hydroxylase (P3OH). The residues of the 2-
10 His-1-Asp motif are indicated by arrows, as is the arginine proposed to bind the 5-carboxylate of 2-oxoglutarate (Mukherji et al (2001) Chem. Comm. 11 972-973).

Figure 10 shows a topographical diagram of the conserved jelly roll core (strands 1 to 8) of 2-oxoglutarate dependent, showing the approximate location of the conserved 2-histidine-1 carboxylate iron binding ligands and 2 oxoglutarate binding
15 basic residue (Arg 557) used to identify candidate HIF-PHs. Numbering refers to the positions in the EGL-9 sequence.

Figure 11 shows an HPLC elution profile of absorbance at 218nm of the hydroxylation of a synthetic peptide by purified PHD1. Synthetic HIF-1 α (B19Pro) peptide was treated with MBP/PHD1 in the presence or absence of 2-oxoglutarate and
20 the products analysed by HPLC as described. Top curve shows results of incubation in the absence of iron(II), upper middle curve shows incubation in the absence of α KG, lower middle shows incubation with all cofactors present. Positions of peaks corresponding to hydroxylated and unhydroxylated B19Pro standards are shown in the bottom panel. Appearance of hydroxylated peptide coincides with the ability to
25 interact with pVHL.

Figure 12 shows an HPLC analysis of the hydroxylation of proline 621 of HIF-1 by EGL-9. *E. coli* were co-transformed with a plasmid expressing either His₆GalHIF-1(590-713) or a mutant (P621G) derivative and plasmids expressing MBP/ Δ .EGL-9 or MBP alone . Retrieved His₆GalHIF-1 was analysed by HPLC for 4-
30 hydroxyproline as shown.

Figure 13 shows NODD and CODD expression plasmids enhance HRE reporter gene activity. (a) Proposed model for peptide effects on HIF α -VHL interaction. Degradation is prevented by the NODD and CODD polypeptides competing for prolyl hydroxylation and/or VHL binding, thereby blocking subsequent ubiquitination. (b) Transfections of the N-terminal or C-terminal ODD (HIF-1 α aa343-417 or aa549-82) led to increased HRE-dependent luciferase activity comparable to hypoxic levels. In contrast, no induction was seen following transfections with corresponding sequences bearing P402A or P564G mutations. *RLU*: Relative Light Units. (c) Use of a CHO cell line, lacking HIF α chains (KA-13), or its stable HIF-1 α transfectant (KH-1) shows dependence of the observed HRE response on HIF-1 α expression. (d) and (e), Amino acids 390-410 for the NODD and amino acids 556-72 for the CODD were the shortest active domains defined. (f), Sequence alignment of human and mouse HIF-1 α NODD and CODD domains.

Figure 14 A and B show dose-response curves for oxalyglycine as well as oxalyl R and S-alanine, Figures 15 to 20 show the relative HIP-PH activity for a variety of inhibitors.

Examples

Example 1. Oxygen regulated modification by enzymatic prolyl hydroxylation targets HIF- α to the von Hippel-Lindau ubiquitylation complex.

In this example it is shown that the interaction between pVHL and a specific domain of the HIF-1 α subunit is regulated by enzymatic hydroxylation of a proline residue (HIF-1 α P564) in a manner that is dependent on oxygen and iron. An absolute requirement of the enzyme for dioxygen as a co-substrate and iron as a co-factor suggests a direct mechanism of cellular oxygen sensing.

In previous studies of the HIF- α /pVHL interaction we found that treatment of cells with cobaltous ions and iron chelators prevented the HIF- α /pVHL association suggesting that the oxygen sensing mechanism might impinge directly on this protein interaction (8). Surprisingly, these studies indicated that the HIF- α /pVHL complex could be retrieved intact from hypoxic cells. Given the rapidity of pVHL dependent proteolysis of HIF- α in oxygenated cells, we considered that re-oxygenation of cell

extracts during cell lysis might have promoted the observed HIF- α /pVHL interaction *in vitro*. To test this we repeated pVHL co-immunoprecipitation experiments in extracts of ^{35}S -methionine/cysteine labelled cells exposed to hypoxia and harvested in a hypoxia work station using deoxygenated buffers, or exposed to hypoxia and
5 harvested conventionally (18).

Experiments were performed on stably transfected renal carcinoma cells expressing haemagglutinin (HA) tagged VHL (RCC4/VHL.HA)(11). RCC4 cells, which lack pVHL, were used as a control. RCC4/VHL.HA cells were labelled with ^{35}S -amino acids in the presence of the proteasomal inhibitor MG132, either in
10 normoxia or hypoxia for 4 hrs. Cells were lysed on ice, either in the hypoxic workstation or on the bench. RCC4 cells were also similarly labelled and lysed and both labelling and lysis were carried out under normoxic conditions. pVHL and associated proteins were captured with anti-HA antibody. As reported previously(8), anti-HA immunoprecipitates captured HIF- α subunits (HIF-1 α and HIF-2 α)
15 efficiently from the proteasomally blocked normoxic cells. However, when hypoxic RCC4/VHL.HA cells were lysed under hypoxic conditions, HIF- α subunits were not co-precipitated with pVHL, despite abundance in the lysate as demonstrated by a HIF-1 α immunoblot. This contrasted with the result using conventional buffers, which had not been deoxygenated, where HIF- α subunits were captured very efficiently.
20 Capture of pVHL and elongins B&C was found to be similar in all RCC4/VHL.HA samples.

Taken together with previously published data, these results indicate that the classical features of regulation by oxygen and iron availability (and interference by cobaltous ions) are reflected in the HIF- α /pVHL interaction *in vivo*, and that
25 promotion of the interaction mediated by oxygen can occur rapidly during the preparation of a cell extract.

1.1: Oxygen sensitivity of the HIF- α /pVHL interaction

^{35}S -Methionine labelled HIF-1 α subunits and pVHL.HA were produced separately by IVTT (in vitro transcription and translation) in reticulocyte lysates.
30 IVTTs were performed under different conditions, mixed, and assayed for interaction

by anti-HA co-immunoprecipitation. These *in vitro* assays allowed analysis of the HIF α /pVHL interaction using the recombinant proteins.

Labelled HIF-1 α and pVHL.HA were generated separately in reticulocyte lysates (IVTT), in the presence or absence of Co (II), desferrioxamine, or Fe(II).

5 Lysates were mixed in various combinations, and interactions assayed by anti-HA immunoprecipitation. We found that supplementary Fe(II) (ferrous chloride, 100 μ M) in the HIF-1 α IVTT greatly enhanced capture by pVHL.HA, whereas addition of Co(II) (cobaltous chloride, 100 μ M) or desferrioxamine, 100 μ M (DFO) to the HIF-1 α IVTT greatly diminished capture. In contrast, pVHL IVTTs performed under these
10 different conditions were all equally effective in supporting the HIF-1 α interaction.

Further experiments were carried out to determine the effect of producing the IVTTs under hypoxic conditions. Labelled HIF-1 α was generated in IVTT reactions in the presence or absence of Fe(II) either under ambient conditions or in a hypoxic workstation. Samples were then diluted in buffer in the hypoxic workstation and
15 purified recombinant GST-VHL-elonginC-elonginB was added. VHL and associated proteins were captured using glutathione-agarose. The results showed that when HIF-1 α IVTTs were prepared under hypoxic conditions, hypoxia reduced their ability to interact with pVHL, irrespective of whether the latter was produced either as a normoxic or hypoxic IVTT, or as a bacterially expressed complex of pVHL and
20 elongins B and C.

Next, we used reticulocyte IVTTs of Gal4/VP16 fusion proteins bearing specific HIF subsequences to show that the regulated interaction with pVHL was supported even by a minimal pVHL binding HIF-1 α subsequence comprising residues 556-574. Fusions bearing amino acid residues 556-574, 549-574, 556-582 or
25 549-582 of HIF-1 α were expressed in reticulocyte lysates with or without added Fe(II). The fusion proteins included the HIF-1 α sequence between a Gal4 DNA binding domain and VP16 transactivation domain. As a control a fusion containing no HIF-1 α sequences was also assessed. Aliquots were assessed for co-immunoprecipitation with pVHL.HA by anti-HA immunoprecipitation. All of the
30 fusions bearing HIF-1 α subsequences displayed iron-dependent recognition by pVHL

including the fusion comprising the shortest region of HIF-1 α subsequence tested comprising residues 556-574. The control fusion, lacking HIF-1 α sequences, did not recognise pVHL either in the presence or absence of iron.

To better understand these findings we surveyed the ability of a series of recombinant pVHL and HIF-1 α products produced in different prokaryotic and eukaryotic expression systems (20) to interact. All pVHL products could interact with HIF-1 α that was derived from mammalian expression systems. In contrast, HIF-1 α could interact only if produced *in vivo* in tissue culture cells, or in reticulocyte IVTT, and not if produced in bacteria, wheat germ lysates, or insect cells. Together, these results indicate that a factor in mammalian cell extracts was necessary to promote the interaction with specific HIF-1 α sequences and that this factor operated in an iron and oxygen dependent manner.

1.2: Modifying activity which promotes interaction of HIF and VHL

To analyse this further we immunopurified a Gal/HIF-1 α /VP16 fusion protein expressing HIF-1 α residues 549-582, from IVTT reactions prepared in the presence of DFO, using anti-Gal antibodies. The unlabelled HIF-1 α substrate was immunopurified on beads, washed, and aliquots incubated under different test conditions in buffer or cell extract. After further washing, the beads were assayed for ability to interact with 35-S labelled pVHL IVTT (21) which was then visualised by fluorography. Increased ability to capture pVHL was seen after exposure of the HIF fusion protein to cell extract in the presence of Fe(II) but not after exposure to Fe(II) without cell extract. The increased ability to capture pVHL after exposure to cell extract and Fe(II) was also found to be oxygen dependent. In analogous experiments it was found that the modifying activity was present in extracts prepared from a variety of mammalian cells, (Hela, RCC, CHO-K1 and rabbit reticulocyte lysate), but that insect cell lysates were essentially inactive on the mammalian HIF fusion protein. The Fe(II) dependent activity of the cell extract was reduced by cooling and was abrogated by pre-heating at 60°C for 10 minutes. The modifying activity did not pass through a 5kDa ultrafilter. Titration of Fe(II) supplementation indicated full activation at 5 μ M. Pre-incubation of the cell extract with hexokinase (50U/ml) and

glucose (50mM) to deplete ATP did not alter activity, though this treatment abrogated the ability of the cell extracts to phosphorylate a control target. Pretreatment of extracts with clotrimazole (10 μ M), methyl-viologen (1mM), or NADase (20mU/ml), did not significantly affect activity.

5 Experiments were also performed using PK epitope tagged HIF-1 α (PK.HIF) expressed in insect cells as a HIF substrate. Both RCC4 cell extract, and reticulocyte lysate, in the presence of Fe(II) promoted the ability of HIF to capture wild type but not mutant (Pro86His) pVHL. Thus human HIF-1 α produced in insect cells required treatment with mammalian extract to promote interaction with wild type but not
10 mutant pVHL (22). Addition of NaCl to the RCC4 cell extract (to 1M final concentration) abrogated the modifying activity, whereas incubation of the PK-HIF in NaCl (1M) after exposure to the cell extract did not alter its subsequent ability to capture pVHL. Likewise treatment of the HIF fusion protein after modification by exposure to extract, with phosphatase or DFO did not prevent pVHL capture. Overall
15 this suggested an enzyme-mediated modification of HIF-1 α that was not phosphorylation.

1.3: Study of the HIF-1 α recognition.

Figure 1A shows alignment of the known or putative pVHL binding domains amongst HIF- α homologues. The effects of selected point mutations in human HIF-
20 1 α on the ability to interact with pVHL were also tested. Wild type (WT) and modified full length HIF-1 α molecules bearing the mutations D556N, D558N, D560Q, P564G, Y565A, P567G, M568R, D569N and D570N were generated in reticulocyte lysate and examined for interaction with VHL.HA by anti-HA immunoprecipitation. Among the tested substitutions, mutation of conserved proline,
25 Pro564Gly totally abrogated interaction and Tyr565Ala reduced the interaction, whereas other mutations had little effect or even enhanced interaction.

Further studies were performed using synthetic polypeptides as inhibitors of the HIF-1 α /pVHL interaction(23). When added to an interaction mix of pVHL.HA and HIF-1 α IVTTs the 34 residue sequence encompassing amino acids 549-582 was
30 unable to block interaction. However, blocking activity was strikingly induced by exposure to cell extract supplemented with Fe(II) (Figure 1B). Induction of blocking

activity showed precisely the same characteristics as had been determined for promotion of interaction between pVHL and recombinant HIF proteins. Results for a series of polypeptides derived from this domain are summarized in Figure 1B and implicate a similar minimal interaction domain. One of the polypeptides shown to
5 have blocking activity following exposure to Fe(II) supplemented cell extract was 19:WT. This polypeptide comprises HIF-1 α polypeptide residues 556 to 574 and its ability to block binding following exposure to a variety of direct oxidation conditions was assessed. Exposure of polypeptide to Fe(II) (100 μ M) with hydrogen peroxide (1mM), NADH oxidase (1U/ μ l) with NADH (1mM), NADH-FMN oxidoreductase
10 (7mU/ μ l) with NADH (1mM) or metachlorobenzoic acid (1mM) did not promote the blocking activity whereas exposure to cell extract plus iron did.

In summary, no polypeptide could block the interaction without prior enzymatic modification, blocking activity could not be induced by a variety of direct oxidation systems, phosphorylation of Tyr565 had no effect on the ability of extract
15 to promote blocking activity, and the mutant sequence Pro564Gly did not block the HIF-1 α /pVHL interaction, even after exposure to extract.

Mass spectrometric analyses (24) (MALDI-Tof) of extract-treated synthetic polypeptide, and recombinant HIF (expressed in insect cells and subsequently treated with mammalian extracts to promote pVHL binding ability) implied several
20 oxidations as evidenced by +16 Da mass shifts in ions derived from this sequence. Further analyses by MS/MS (ESI-QTof) indicated oxidation affecting Pro564 and the nearby methionine residues. Since the methionine residues are either non-conserved or could be mutated without effect, and direct oxidation methods known to oxidize methionine efficiently could not mimic the enzymatic activity, we postulated that the
25 enzymatic oxidation that promoted interaction of this HIF-1 α sequence with pVHL was the oxidation of Pro564.

1.4 Hydroxyproline incorporation into a synthetic polypeptide.

We synthesized a polypeptide (HIF-1 α residues 556-574), containing a *trans*-4-hydroxy-*S*-proline residue at position 564 (19:Pro564Hyp), since the *trans*-4-
30 hydroxylation is the commonest enzymatic proline oxidation (25). A polypeptide blocking assay was carried out using the 19:Pro564Hyp modified polypeptide and, as

a control, the unmodified polypeptide without the *trans*-4-hydroxylation (19:WT). The 19:WT polypeptide was either incubated with cell extract before the binding assay or was untreated. The 19:Pro564Hyp polypeptide was added to a mixture of HIF-1 α and pVHL-HA IVTTs at a concentration of 1, 0.25, 0.05 or 0.01 μ M and the cell extract treated or untreated 19:WT polypeptide at a concentration of 0.5 μ M. Interaction of HIF-1 α with pVHL was then assayed by anti-HA immunoprecipitation. The hydroxyproline substituted polypeptide (19:Pro564Hyp) was highly effective at inhibiting the HIF-1 α /pVHL interaction without the need for modification by cell extract. The 19:WT control unsubstituted equivalent polypeptide showed the expected requirement for cell extract in order to inhibit interaction. Control reactions carried out with no blocking polypeptide showed the expected binding of HIF-1 α to pVHL-HA. A pVHL capture assay was then carried out using biotinylated synthetic polypeptides. The same polypeptides, 19:Pro564Hyp and 19:WT were assessed for the ability to capture wild type or mutant (Pro86His) pVHL. 19:WT control polypeptide captured pVHL only after incubation with cell extract, whereas 19:Pro564Hyp captured wild type pVHL without pre-treatment. In both cases capture was specific for wild type, as opposed to mutant, pVHL.

In summary, in striking contrast with previously tested polypeptides, 19:Pro564Hyp blocked the HIF-1 α /pVHL interaction without the need for exposure to cell extract. Moreover, a biotinylated version of 19:Pro564Hyp specifically captured wild type but not mutant pVHL, and its ability to capture pVHL was not increased further by incubation with cell extract (26). In comparison, the equivalent unmodified synthetic polypeptide 19:WT could not interact without prior incubation with cell extract.

These results reveal that the enzymatic activity promoting interaction of HIF-1 α with pVHL is a prolyl-4-hydroxylase, which we term HIF- α prolyl-hydroxylase (HIF-PH). All previously described prolyl-4-hydroxylases are members of the superfamily of 2-oxoglutarate-dependent, and related, dioxygenases (27). Consistent with the data presented above none of these enzymes have an absolute requirement for ATP or NAD(P) but they do have an absolute requirement for Fe(II) as a co-factor and dioxygen as a co-substrate (27). Structural studies within the class have defined a

non-haem iron centre co-ordinated by an HXD/E...H motif (28). Interestingly, and consistent with our findings, the Fe(II) is not firmly bound and can be readily removed by chelating agents, and enzyme inhibition occurs following substitution of Fe(II) with Co(II) or Ni(II)(25).

5 1.5: Effect of ascorbate supplementation and various inhibitors on HIF-PH activity.

The capture of labelled pVHL by different HIF substrates was monitored after exposure to various test conditions.

The effect of ascorbate on pVHL capture by a Gal/HIF-1 α 549-582/VP16 fusion protein substrate was monitored and ascorbate (2mM) was found to enhance
10 the modifying activity of cell extract, but have no effect in the absence of cell extract. Ascorbate therefore enhances the activity of HIF-PH.

We went on to test a series of 2-oxoglutarate analogues which act as competitive inhibitors of this class of enzyme (29) for ability to inhibit HIF-PH as assessed by the ability of cell extracts to modify either HIF polypeptide (19:WT) or a
15 HIF fusion protein (Gal/HIF-1 α 549-582/VP16) so as to promote pVHL capture. Concordant results were obtained with both sources of HIF sequence. In one such experiment the effect of N-oxalylglycine on pVHL capture by a biotinylated HIF polypeptides as substrate was monitored. The WT:19 and 19:Pro564Hyp polypeptides described above were used as substrates. N-Oxalylglycine (0.2-1mM)
20 was found to completely inhibit the modifying activity of cell extract on 19:WT. Inhibition by N-Oxalylglycine was overcome by addition of 5mM 2-oxoglutarate. As previously, 19:Pro564Hyp captured pVHL efficiently without modification by cell extract, and this was not influenced by exposure to N-oxalylglycine. Similar inhibition, also competed by 2-oxoglutarate, was observed with N-oxalyl-2S-alanine
25 but not the enantiomer N-oxalyl-2R-alanine, demonstrating that the effect was not due to simple Fe(II) chelation in solution. We also used a 2-oxoglutarate dependent dioxygenase, phytanoyl-CoA α -hydroxylase(30), together with a readily available unnatural substrate (isovaleryl CoA) (31) to deplete the cell extract of 2-oxoglutarate produced by the citric acid cycle; as predicted, this prevented the subsequent
30 modification of HIF polypeptide. The effect of dimethyl-oxalylglycine on HIF-1 α expression was also studied. HIF-1 α immunoblot analysis of extracts of Hep3B and


absence of doxycycline for 16 hours. Immunoblots were then performed on cell lysates to quantify levels of HIF-1 α and also to check for expression of PHD via the PK tag.

- Exposure of cells to doxycycline for 16 hours induced expression of PK tagged PHD1. The induced expression of PHD1 substantially reduced expression of HIF-1 α in modest hypoxia (3% oxygen) and also reduced expression to a lesser extent under total hypoxia (0% oxygen). Thus in this Example, the expression of endogenous HIF-1 is shown to be strikingly dependent on the activity of the specifically induced PHD1 isoform under the conditions of assay.

10 EXAMPLE 8

(S)-2-(Methoxyoxalyl-amino)-pentanedioic acid diethylester (IS12)

or: *Diethyl N-methoxyoxalyl-(L)-glutamate (IS12)*



- To a stirred solution of 10 mmol (2.40g) of diethyl (L)-glutamate hydrochloride in 10 ml of toluene, 10 mmol (1.23 g, 0.93 ml) of methyl oxalyl chloride was added and heated until no further HCl gas evolved (4-6 hr). The solvent was evaporated yielding 2.86 g (9.9 mmol, 99%) of IS12 as a yellowish oil, $[\alpha]_D^{25} -28.3^\circ$ (c 1 in methanol); ν_{max} (NaCl)/cm⁻¹ 1738, 1705 (C=O); δ_H (200 MHz; CDCl₃) 1.22, 1.26 (6 H,

EXAMPLE 6

In this example it is shown that enhanced new vessel growth can be stimulated in a murine subcutaneous sponge angiogenesis assay by injection of the HIF prolyl hydroxylase inhibitor, dimethyl oxalylglycine.

- 5 In previous studies we have demonstrated that N-oxalylglycine is an inhibitor of PHD activity *in vitro* but seems to be incapable of entering intact cells. Application of an esterified form, dimethyloxalylglycine, to tissue culture cells results in stabilisation of HIF alpha chains (Example 1) and activation of transcription of endogenous HIF target genes (Example 5).
- 10 Implantation of a polyurethane sponge subcutaneously in a mouse provides an inflammatory stimulus to angiogenesis and is a well established model for assessing the pro- and anti- angiogenic effects of compounds. To test the effects of dimethyl oxalylglycine *in vivo*, sterile 8mm sponge discs were inserted under the dorsal skin of C57 Black mice on Day 0. Test solutions were injected through the skin into the
- 15 sponges of mice once per day on days 1, 2, 4 and 5. Individual mice received 100 microlitres aliquots of either sterile dimethyloxalylglycine (0.1mM, 1mM or 10mM) or carrier solution. Animals were sacrificed on day 7 and the sponges removed. The sponges were fixed in 3.7% formaldehyde, paraffin embedded and stained immunohistochemically for von Willebrand factor to identify blood vessels.
- 20 Considerably more blood vessels were observed in sponges injected with dimethyloxalylglycine (1mM) than those receiving solvent alone.

EXAMPLE 7: The effect of PK-tagged PHD1 expression on HIF-1- α induction by hypoxia

- 25 In this Example it is shown that a recombinant PHD (PHD1) may be overexpressed in a tissue culture cell line in such a manner as to affect the metabolism of a HIF polypeptide.

U2OS cells were stably transfected with a binary system encoding a tetracycline operator fused to an activator, and a plasmid encoding C-terminal PK

30 epitope tagged PHD1 under control of a tetracycline response element. The transfected cells were incubated with 21%, 3% or 0% oxygen in either the presence or

but also a range of systemic functions such as the control of oxygen delivery through effects on angiogenesis, vasomotor control, and erythropoiesis. These complex requirements have argued against the concept of a single oxygen sensor. However, the existence in mammalian cells of (at least) three isoforms of HIF.PH, and (at least) two
5 isoforms of HIF- α , each with more than one site of prolyl hydroxylation (Masson et al., 2001 supra), may provide the potential for different physiological responses to oxygen availability to be generated through combinatorial interactions amongst these molecules.

The characterisation of the HIF PH enzymes described herein has various therapeutic
10 applications, in particular as targets in the development of pharmacological agents which modulate HIF- α levels in a cell.

EXAMPLE 5

In this Example it is shown that HIF-1 α protein and the endogenous HIF target gene encoding carbonic anhydrase 9 (CA-9) are induced by exposure of cells to
15 the PHD inhibitor, dimethyl oxalylglycine. In previous studies we have demonstrated that N-oxalylglycine is an inhibitor of PHD activity *in vitro*, but seems to be incapable of entering intact cells. The esterified form, dimethyloxalylglycine, has therefore been used to deliver the compound to tissue culture cells.

Hep3B and U2OS cells were exposed to either 0.1 mM or 1 mM
20 dimethyloxalylglycine for 6 hours, harvested and assayed by immunoblotting (Western blotting) for changes in the level of HIF-1 α and CA-9 expression. Controls, where no inhibitor was added, were also performed. Clear upregulation of both HIF-1 α and the HIF target gene product CA-9 are observed in the presence of dimethyloxalylglycine. Upregulation of HIF-1 α increased with increasing
25 concentration of dimethyloxalylglycine, whilst the level of CA-9 expression was similar after exposure to both 0.1 and 1 mM of dimethyloxalylglycine.

Treatment with cell extract in the absence of inhibitor showed strong binding of pVHL. Treatment with cell extract in the presence of oxalyl glycine (NK 87) showed reduced binding of pVHL, showing that oxalyl glycine inhibits the HIF prolyl hydroxylase. Treatment with cell extract supplemented with additional 2-oxoglutarate produced strong pVHL binding. Treatment with cell extract in the presence of NK87 and additional 2-oxoglutarate also produced strong binding, indicating that HK87 competes with the oxoglutarate co-substrate.

Treatment with cell extract in the presence of 1mM NMPG produces strong binding of pVHL. However, treatment in the presence of 5mM NMPG reduced the amount of pVHL binding.

As a positive control, pVHL was captured when Gal-HIF-VP16 substrate was synthesised in rabbit reticulocyte lysate in the presence of additional ferrous chloride. Other potential inhibitors as shown in Table 3 were screened for the ability to inhibit HIF hydroxylase activity as described above.

Of the compounds screened in this assay, reduced pVHL binding indicative of inhibition of HIF hydroxylation was observed for Is1, Is3, Is8, benzohydroxamic acid, ethyl dihydroxybenzoate, and NK45.

The present application relates to the characterization of a HIF-1/VHL prolyl hydroxylase system and the identification of a new functional group of 2-oxoglutarate dependent oxygenase that function as HIF prolyl hydroxylases (HIF-PHs). The critical role of these enzymes in the regulation of HIF is emphasised by analysis of *vhl-1* and *egl-9* mutant worms, which show essentially complete loss of regulation of HIF-1 by oxygen. The availability of recombinant HIF-PHs permits further investigation of the HIF/VHL system and an important challenge will be to determine the extent to which the complex demands of physiological oxygen homeostasis are met by the biochemical properties of these enzymes.

Identification of the HIF system in nematode worms that obtain oxygen directly by diffusion reveals that this system of gene regulation must have evolved before the development of complex systemic oxygen delivery systems, presumably to regulate responses to oxygen availability at the cellular level.

In mammals, the HIF system regulates not only cellular responses to oxygen,

overall inhibitor mediated increase in HIF activity.

Addition of dimethyloxalylglycine to Hep3B and U20S cells in tissue culture (0.1mM, 1mM) was also observed to increase intracellular levels of HIF-1 in Western Blot experiments.

- 5 Effects of forced expression of EGLN2 (PHD 1) or a naturally occurring splice variant lacking amino acids 369-389 (PHD4) on HIF Activity

Hep3b cells were co-transfected with a mixture of three plasmids; pHRE.luc, encoding a HIF responsive luciferase gene, pcDNA3 or pcDNA3.HIF, mammalian expression plasmids leading to expression of no product or full length human HIF-1
10 alpha, and pCMV.beta-gal, encoding a constitutively expressed betagalactosidase gene as a transfection control.

48 hours following transfection, cells were incubated in normoxia or 2% hypoxia overnight. Cell lysates were assayed for luciferase and beta galactosidase activity and the relative luciferase activity in each sample determined.

- 15 In all circumstances, relative luciferase activity was lower when co-expression included the full length EGLN2 (PHD1) rather than the deleted, non- functional version (PHD4)(figure 7), providing indication that expression of full length EGLN2 reduces functional HIF by enhancing the generation of the rapidly destroyed hydroxylated HIF protein.

- 20 The effect was even more prominent in circumstances where the level of HIF-1 α would be expected to be higher (e.g. when co-expressed from a plasmid or partially stabilised by modest hypoxia). This demonstrates that expression of these gene products is able to enhance HIF-1 degradation in vivo.

4.12 Effect of inhibitors of HIF prolyl hydroxylase activity

- 25 Modification and pVHL binding assays were performed as described above (and by reverse phase HPLC) to determine the effect of inhibitors on the ability of cell extract to modify Gal-Hif549-582-VPI6, thereby allowing capture of radiolabelled recombinant pVHL.

Binding of pVHL was determined using SDS-PAGE and autoradiography.

- 30 In the absence of treatment with cell extract, no binding of pVHL was observed, showing the Gal-Hif549-582-VPI6 was unmodified.

no insert, an insert encoding the full open reading frame of the human homologue of *C. elegans* Egl-9 known as EGLN-2, or an insert encoding the full open reading frame of the human homologue known as EGLN-3. Modification was performed in the absence of additives, in the presence of oxalylglycine, oxalylglycine plus 2-oxoglutarate, 200 μ M desferrioxamine or 200 μ M cobaltous chloride.

Enzyme activity was observed to be diminished by oxalylglycine, desferrioxamine and cobaltous ions. The inhibitory effect of oxalylglycine was partially competed by addition of excess 2-oxoglutarate. The family of 2-oxoglutarate dependent dioxygenases demonstrate a requirement for oxygen, iron and 2-oxoglutarate. The ability of these gene products to modify HIF-1 to a VHL binding form was examined in differing conditions of iron availability, 2-oxoglutarate availability and in the presence of a 2-oxoglutarate inhibitor. These results demonstrated an iron and 2-oxoglutarate dependence of activity in reticulocyte lysate.

4.11 Effects of dimethyl oxalyl glycine on HIF activity in vivo.

Hep3b and U20S cells were co-transfected with a mixture of three plasmids; pUAS.tk.luc, encoding a GAL 4 responsive luciferase gene, pgal-hif775-826, a mammalian expression plasmid leading to expression of a fusion between a 147 amino acid DNA binding domain of GAL 4 and the carboxy terminal transactivator of human HIF-1 alpha, and pCMV.beta-gal, encoding a constitutively expressed beta-galactosidase gene as a transfection control.

48 hours following transfection, cells were incubated in normoxia or 2% hypoxia overnight in the presence or absence of dimethyl oxalyl glycine as indicated. Cell lysates were assayed for luciferase and beta galactosidase activity and the relative luciferase activity in each sample determined.

In both cell lines, the presence of the HIF prolyl hydroxylase inhibitor resulted in enhanced activity of the carboxy terminal transactivator in both normoxia and hypoxia compared to the untreated samples (Figure 6). This result shows a potentiating action of this inhibitor on a domain of HIF which is not normally considered to be dependent on proteolytic destruction for its activity.

Potential of the action of the carboxy terminal transactivator coupled with inhibition of destruction via the oxygen dependent degradation domains enhances the

Modification and pVHL binding assays were performed as described above. Wheat germ lysate was programmed with pcDNA3 based plasmids containing no insert, an insert encoding the full open reading frame of the human homologue known as EGLN-2, or an insert encoding the full open reading frame of the human
5 homologue known as PHD-3 or EGLN-3.

HIF substrates for modification were wild type or contained mutation of proline 564 to glycine. The pVHL target for capture was wild type or contained the mutation of tyrosine 98 to histidine.

Binding of labeled pVHL was observed in assays using wild type HIF and
10 pVHL. No binding was observed in assays using mutant HIF or pVHL.

Both EGLN2 and EGLN3 were therefore able to modify wild type but not mutant HIF in a manner allowing capture of wild type but not mutant pVHL.

4.9 Oxygen Dependence of Modification of HIF by Recombinant Enzymes

Modification and pVHL binding assays were performed as described above
15 except that enzymes were generated by expression in COS cells or rabbit reticulocyte lysate.

Plasmids used to generate enzymes were as follows; pcDNA3 (without insert); pcDNA3 containing sequence encoding rat SM20 lacking the first 59 amino acids; pcDNA3 containing sequence encoding EGLN2 (PHD 1). Modification of HIF
20 substrate by enzymes was performed in either normoxia (21% O₂) or anoxia (2% O₂) conditions using a hypoxia workstation.

Given the regulation of HIF-1 by oxygen and the known substrate requirement of 2-oxoglutarate dependent dioxygenases for oxygen, the oxygen dependence of the HIF modifying activity was examined.

25 In anoxia, EGLN2(PHD1) or rat SM20 (lacking the amino terminal 59 amino acids) were unable to modify HIF for pVHL binding, in contrast to the clear modification at 21% O₂. This demonstrates oxygen dependence and a means of oxygen sensing in the regulation of HIF-1.

4.10 Effects of Oxalylglycine and 2-Oxoglutarate on EGLN2 and EGLN3 in vitro

30 Modification and pVHL binding assays were performed as described above. Rabbit reticulocyte lysate was programmed with pcDNA3 based plasmids containing

of 100 μ M ferrous chloride. The nucleotide sequence corresponding to these putative proteins were generated by PCR and their identity was confirmed by sequencing. The protein products generated conformed with their predicted molecular weights.

Protein containing HIF-1 sequence (amino acids 549-582) was generated in a
5 reticulocyte in vitro transcription translation reaction in the presence of 100
micromolar desferrioxamine, retrieved and then exposed to wheat germ translates
containing the putative enzymes as described above. These proteins contain the
critical proline (564) which can be modified by hydroxylation and which enables
recognition by von Hippel Lindau tumour suppressor protein (pVHL).

10 The modification of the HIF-1 sequence was assayed by the binding of the
HIF-1 to radiolabelled pVHL, generated by in vitro transcription and translation of a
pcDNA3 vector containing the human wild type pVHL open reading frame in a rabbit
reticulocyte lysate in the presence of 35 S-methionine.

SDS PAGE analysis indicated that the plasmids encoding the genes EGLN-2,
15 full length and truncated rat SM20 produced, in the presence of Fe^{2+} , a clear
modification of HIF-1 which allowed capture of labeled pVHL.

No pVHL binding was observed in the absence of Fe^{2+} for SM20, or EGLN-2
and only a low level of binding was observed for truncated SM20.

4.7 Mutation of EGLN2

20 Modification and pVHL binding assays were performed as described above.
Rabbit reticulocyte lysate was programmed with pcDNA3 based plasmids containing
no insert, an insert encoding the full open reading frame of the human homologue
known as EGLN-2, a mutant form of EGLN-2 with a Histidine to Alanine
substitution at amino acid residue 358 or a naturally occurring splice variant lacking
25 amino acids 369-389.

pVHL binding was observed only with the full length wild type PHD-3
polypeptide. The full length wild type enzyme was able to modify HIF -1 sequence
whilst neither the mutant form nor the deleted splice variant was able to do so. This
demonstrates that His358 and the region between residues 369 and 389 are necessary
30 for HIF hydroxylase activity.

4.8 Effects of HIF and pVHL Mutations

it has been predicted that HIF-PH would itself be induced by the transcriptional response to hypoxia.

RNase protection demonstrated that all three HIF-PH mRNAs are expressed in HeLa cells and that, in this cell line, transcripts for EGLN1 (PHD2) and EGLN3 (PHD3) but not EGLN2 (PHD1) are induced by hypoxia. In keeping with this, semi-quantitative analysis of lysates prepared from HeLa cells that had been grown in normoxia or exposed to hypoxia for 16 hours then assayed for HIF-PH activity *in vitro* using the pVHL capture assay, demonstrated induction of total HIF-PH activity that was blocked by actinomycin D.

Finally, we used pVHL capture assays to measure the activity *in vitro* of recombinant EGLN2 on a HIF-1 α 549-582 substrate at graded levels of hypoxia in a controlled hypoxia work-station. We first measured the effect of graded hypoxia on the HIF modifying activity of extracts of vhl-defective RCC4 cells that contain a relatively high level of total HIF-PH activity. A progressive reduction in activity was observed with graded hypoxia. Similar assays were then performed using EGLN2 produced in reticulocyte lysate by IVTT, or purified MBP/PHD-1 obtained by expression in *E. coli*. Closely similar progressive reductions in the activity of each preparation were observed with graded hypoxia. Thus, the oxygen dependent activity of recombinant PHD-1 from either source parallels that observed in crude cell extracts, and mirrors the progressive increases in HIF- 1 α protein and DNA binding that are observed when cells are exposed to graded hypoxia in culture (Jiang et al., (1996) Am. J. Physiol. 271 C1172-C1180).

4.6 HIF Prolyl Hydroxylase Activity

Full length rat SM20, a truncated form of rat SM 20 lacking the amino terminal 59 amino acids and the human homologue EGLN-2 were shown to modify HIF amino acids 549-582 in a manner which facilitates interaction with the VHL protein. This is known to depend on hydroxylation of proline 564.

Wheat germ lysate was programmed with pcDNA3 based plasmids containing no insert, an insert encoding the full open reading frame of rat SM20, a truncated form of rat SM20 lacking the amino terminal 59 amino acids or the human homologue known as EGLN-2 in the absence of exogenous iron or with the addition

that stably express HA tagged human pVHL, and tested for interaction by anti-HA immunoprecipitation.

With full length wild-type HIF-1 α , striking activity was observed with rat SM-20 and all three human gene products, but not a mutant EGLN1 bearing an H358A substitution at the predicted catalytic site, and not a different human 2-oxoglutarate dependent oxygenase (phytanoyl coenzyme A hydroxylase (Mukherji et al., 2001) that was tested as a negative control. Similar results were obtained with HIF-2 α .

Examination of HIF-1 α mutants bearing missense substitutions at the critical prolyl residues in the: HIF-1 α ODDD (Masson et al 2001) showed that that enzymes were differentially efficient at promoting interaction via the C-terminal (P564) and N-terminal (P402) prolyl hydroxylation sites. Whereas interaction through the C-terminal site could be promoted by all enzymes, VHL E3 capture was less efficient when only the N-terminal site (P402) was intact, and was only promoted by EGLN2 (PHD1) and EGLN1 (PHD2). No activity at all was observed with a double HIF-1 α mutant, P402A P564G, that ablates both hydroxylation sites.

In keeping with these results, all enzymes strongly promoted interaction of pVHL with isolated HIF-1 α sequences {residues 549-582} from the C-terminal site. Further analysis demonstrated that this activity was strongly inhibited by iron chelation, cobaltous ions, and the 2-oxoglutarate analogue N-oxalylglycine.

To confirm direct action on HIF- α sequences, we prepared purified EGLN2 as an MBP fusion protein in *E. coli* and assayed activity using either purified His-tagged HIF-1 α polypeptides containing the N-terminal (344-503) or C-terminal (530-698) hydroxylation sites, or a synthetic peptide consisting of the minimal HIF-1 α C-terminal substrate (B19Pro, residues 556-574). These experiments demonstrated activity by pVHL capture assays, HPLC/MS detection of the hydroxylated peptide product or derivatised 4-hydroxyproline, and by 2-oxoglutarate decarboxylation assays.

To verify expression of all three isoforms, we performed RNase protection analysis using riboprobes specific for each transcript. Since recently published work has indicated that the rate of HIF degradation in normoxia is enhanced by prior exposure of cells to a period of hypoxia (Berra et al. (2001) FEBS Letters 491 85-90)

MBP/ Δ N.EGL-9 or MBP. His₆/Gal4/HIF-1 substrates were retrieved by nickel affinity chromatography and aliquots assayed for ability to capture 35S-methionine labelled HA.VHL-1 using anti-Gal immunoprecipitation, or subjected to acid hydrolysis and HPLC analysis for the presence of phenylisothiocyanate derivatised 4-hydroxyproline.

In concordance with the HA.VHL-1 capture assay results, 4-hydroxyproline was produced in the wild type but not the mutant His₆/Gal4/HIF-1 substrate following exposure to active enzyme (Figure 12) These results therefore demonstrated the activity of EGL-9 as a prolyl hydroxylase that targets HIF-1 to VHL-1 in *C.elegans*.

4.5 Identification of a Series of Mammalian HIF-PH Isoforms.

Sequence similarity between EGL-9 and a rat gene product termed SM-20 (Wax et al., (1994) J. Biol. Chem. 269 13041-13047) has been noted previously though no functional connection was recognised (Darby et al.(1999) supra). Our sequence-structure search identified a larger series of homologies and predicted three closely related genes in each of the human and rodent genomes that bore striking homology to *egl-9*, in particular over the core putative catalytic domain.

Figure 9 illustrates sequence alignment of EGL-9 (acc. no. AAD56365) SM.20 (acc. no, AAA 19321) and the predicted human proteins defined by acc. nos. XP_040482, AAG33965, and NP_071356 (EGLN1-3), corresponding to Unigene clusters Hs.324277, Hs.6523, and Hs. 18878.

The human protein products have been termed EGLN1, 2, and 3 or 'Prolyl Hydroxylase Domain containing' (PHD) 2, 1 and 3 respectively. Note that the gene termed EGLN3 or PHD3 has previously been identified as human homologue of rat SM-20 (Dupuy et al. (2000) Genomics 69 348-354).

To test the role of these gene products in regulating the interaction between human HIF α sub-units and the VHL E3 ligase complex, we first produced the proteins by reticulocyte IVTT. Since unprogrammed lysate has a low level of HIF-PH activity (Jaakkola. et al., 2001 supra) we tested for enhanced ability of the programmed lysates to promote the HIF α /VHL E3 interaction. After incubation with relevant enzyme, HIF- α substrates were mixed with extracts from 786-0/VHL cells

To determine the effect of EGL-9 on the HIF-1 transcriptional response, we measured mRNA levels of a range of hypoxia inducible transcripts and found striking up-regulation in *egl-9* worms. A strongly inducible mRNA of unknown function (F22B6.4) was also identified. These findings demonstrated a critical function for
5 EGL-9 in the regulation of HIF-1 and provided further indication that EGL-9 functions as a HIF.PH that targets HIF-1 to VHL-1.

We produced recombinant EGL-9 and assessed its ability to catalyse the post-translational modification of HIF-1. HIF-1 was captured efficiently by HA.VHL-1 after incubation with EGL-9 programmed reticulocyte or wheat germ lysates, but not
10 unprogrammed lysate.

In contrast, IVTTs expressing recombinant *C.elegans* PHY-1, PHY-2 and the gene product of the predicted ORF T20B3.7 that also has significant homology to known prolyl hydroxylases, had no activity in these assays.

To test whether EGL-9 could act directly on HIF-1, further preparations were
15 made by baculoviral expression in insect cells and by expression as maltose binding protein (Map) fusion proteins in *E. coli*. Since full length MBP/EGL-9 protein was insoluble when expressed in *E. coli* we prepared an N-terminal truncation containing residues 359-723 (MBP/ Δ N.EGL-9) that preserved the predicted catalytic domain and had HIF.1 modifying activity when expressed as an IVTT. *C.elegans* HIF-1 substrates
20 were made as N-terminal Gal4 fusion proteins in either insect cells or *E.coli* and purified by anti-Gal immunoprecipitation. These substrates were incubated with lysates of insects cell expressing full length EGL-9 or purified MBP/ Δ N.EGL.9, and tested for ability to capture VHL-1.

Both forms of recombinant EGL-9 efficiently promoted modification of HIF.1
25 as indicated by HA.VHL-1 capture. Moreover analysis of this activity demonstrated 2-oxoglutarate, iron, and oxygen dependence, and direct inhibition by cobaltous ions.

To demonstrate that activity in the HA.VHL-1 capture assays corresponded to hydroxylation of the critical HIF-1 residue P621, we assayed modified HIF-1 polypeptides for 4-hydroxyproline content by HPLC. To provide larger quantities of
30 protein for this analysis, we co-transformed *E.coli* with wild type or the P621 to G mutant form of a His₆/Gal4/HIF-1(590-719) (HGH) fusion protein and either

results demonstrated that conservation of the HIF/pVHL system in *C.elegans* extends to the mode of regulation by prolyl hydroxylation.

4.4 The *C.elegans* egl-9 Gene Product is a Prototype HIF-PH.

The best characterised prolyl hydroxylases are the procollagen-modifying enzymes (Kivirikko and Myllyharju, (1998) Matrix Biol. 16 357-368). However, worms containing inactivating mutations in each of two isoforms of the catalytic subunits, dpy-1B (also termed *phy-1*) and *phy-2* (Friedman et al., 2000 supra: Winter and Page, 2000 supra) showed normal HIF-1 regulation, consistent with HIF-PH, being distinct from the collagen modifying enzymes.

We searched *C.elegans* and mammalian databases for additional HIF-PH candidate genes that were well conserved between these species and possessed a common β -barrel jelly roll motif.

Of particular interest was a family of genes related to the *C. elegans* gene *egl-9*, a gene of previously unknown function that was first identified on the basis of an egg-laying abnormal (*egl*) phenotype (Trent et al., (1983) Genetics 104 619-647).

Sequence analyses coupled with secondary structure predictions in the light of crystallographic data (Valegard et al. (1998) supra: Zhang et al, (2000) Nature Structural Biology 7 127-133) predicted that these genes would encode a family of enzymes conserved in *C.elegans* and mammals. The predictions suggested that the enzymes would contain not only the jelly roll motif, but also conserved iron and 2-oxoglutarate binding residues in the same relationship that they occur in crystallographically characterised enzymes e.g. the HXD...H iron binding motif on the second and seventh strands of the jelly roll motif.

Mutants worms containing defective *egl-9* alleles were therefore assessed for regulation of HIF-1 by immunoblotting. Three strains bearing inactivating mutant alleles of *egl-9*, (*sa307*, *sa330*, and *n571*) (Darby et al., (1999) PNAS 96 15202-15207; Trent et al.(1983) Genetics 104 619-647) all showed striking constitutive up-regulation of HIF-1 in normoxia and loss of induction by hypoxia. Moreover, a further temperature sensitive *egl-9* mutant, *n586* showed enhanced normoxic HIF-1 level at the non-permissive temperature.

A series of N-terminal truncations of HIF-1 linked to a Gal4 DNA binding domain was constructed. The Gal/HIF-1 fusion proteins were expressed in reticulocyte lysates, pre-incubated with worm extracts and then tested for interaction with HA.VHL-1. These experiments demonstrated that whilst N terminal truncations up to and including Gal4/HIF-1(590-719) were captured efficiently by HA.VHL-1, Gal4/HIF-1(641-719) was not, implicating HIF-1 amino acids 590-641 in the interaction.

Inspection of this region revealed homology to pVHL-binding domains in human HIF-1 α that have recently been shown to contain sites of prolyl hydroxylation (Ivan et al., (2001) Science 292 464-468; Jaakkola et al. Science (2001) 292 468-472). We therefore mutated the homologous prolyl residue in *C.elegans* HIF-1 (P621 to G) and found that this mutation ablated interaction with HA.VHL-1.

The demonstration of a critical conserved prolyl residue in *C.elegans* HIF-1, together with the need for pre-incubation with worm extract provided indication that the mechanism regulating the HIF-1/VHL-1 interaction through enzymatic prolyl hydroxylation might also be conserved in *C. elegans*. To verify this, N-oxalyl-2S-alanine, a 2-oxoglutarate analogue that inhibits this class of enzymes (Cunliffe et al.(1992) supra) was added to the worm extract during pre-incubation with HIF-1. This strongly inhibited activity in a manner that was competed by excess 2-oxoglutarate, as inhibition was antagonized by 5mM 2-OG.

To test whether hydroxylation of the critical P621 residue in *C.elegans* HIF-1 could indeed promote binding to VHL-1, we synthesised N-terminal biotinylated peptides corresponding to residues 607-634 of *C. elegans* HIF-1 that contained either a proline (B28Pro) or a (2S,4R)-*trans*-hydroxyproline residue (B28Hyp) at position 621. We found that B28Hyp but not B28Pro blocked capture of pretreated HIF-1 by HA.VHL-1, when added to the interaction mix.

Furthermore, B28Hyp but not B28Pro captured immunodetectable native VHL-1 when mixed with extracts from wild type but not *vhl-1* mutant worms. Finally, to test the importance of prolyl hydroxylation in regulating *C.elegans* HIF-1 *in vivo* we exposed worms to the cell-penetrating prolyl hydroxylase inhibitor, dimethyloxalylglycine. This strongly induced HIF-1 in normoxic worms. These

tested concentrations of 0.5% and 0.1% oxygen. In 0.1% oxygen, HIF-1 protein level was strongly induced within 4hrs, and sustained over 24hrs, but disappeared within minutes following re-oxygenation. In contrast, *hif-1* mRNA levels were unchanged by hypoxia. Thus, these experiments confirmed up-regulation of HIF-1 by hypoxia, and suggested a mode of regulation at the protein level similar to that described for mammalian HIF- α subunits.

4.2 Critical function of a pVHL homologue VHL-1 in the regulation of HIF-1 in *C.elegans*.

We compared HIF-1 expression in wild type and a series of mutant worms that were selected because of potential relevance to previously proposed models for oxygen sensing and signal transduction processes in the mammalian HIF system (Chandel et al., (2000) J. Biol. Chem. (Aug) 1-37; Ehleben et al. (1997) Kidney Int. 51 483-491; Zundel et al. (2000) Genes & Dev. 14 391-396) for review see (Semenza, (1999) Cell 98 281-284). These included mutants in the PTEN/insulin receptor/PI-3-kinase pathway (*daf-18*, *daf-1*, *age-1*), a mutant in a putative homologue of VHL (*vhl-1*), mutants affecting mitochondrial proteins (*mev-1*, *clk-1*, *gas-1*), a mutant that affects cytosolic catalase activity *ctl-1*, and others selected for resistance or sensitivity to oxidant stresses but where the mutant gene is not yet characterized (*mev-2*, *mev-3*).

With the exception of *vhl-1* all mutant worms showed preserved regulation of HIF-1 protein levels. In contrast, the *vhl-1* worms showed high levels of HIF-1 in normoxia that were essentially unregulated by oxygen. These results confirmed that proposed homology for *vhl-1* (Woodward et al.(2000) Genomics 65 253-265), and indicated a conserved role for *C.elegans* VHL-1 protein in the response to hypoxia.

4.3 Interaction of HIF-1 with VHL-1 is Regulated by Prolyl Hydroxylation.

To address the mechanism of regulation of HIF-1 by VHL-1, interaction between the two proteins was tested. ³⁵S-methionine labelled haemagglutinin (HA) tagged VHL-1 (HA.VHL-1), and HIF-1 were produced separately *in vitro* in coupled transcription translation reactions (IVTT) in reticulocyte lysate. IVTTs were then mixed and assayed for interaction by anti-HA immunoprecipitation. When produced this way, the proteins did not interact. However, when recombinant HIF-1 was pre-incubated with worm extract, a clear interaction was observed.

The reaction was incubated in an end-over-end rotator for 2 hours in cold room, then spun at 2.000rpm for 1 minute and the supernatant removed (radioactive liquid waste). The beads were washed with 500 μ l of EBC buffer and 100 μ M DFO and spun again at 2.000 rpm for 1 minute. The wash steps were repeated a total of 5 times. The supernatant was removed from the final wash and eluted in 15 μ l of 2x SDS sample buffer.

Samples were stored at -20°C and examined by SDS-PAGE.

DNA and Protein Manipulation

DNA manipulation and cloning and protein expression and analysis by SDS-Polyacrylamide Gel Electrophoresis (SDS-PAGE) were performed according to standard techniques which are well known to those of skill in the art and described in detail in Molecular Cloning: a Laboratory Manual: 2nd edition, Sambrook et al., 1989, Cold Spring Harbor Laboratory Press and Current Protocols in Molecular Biology, Ausubel et al. eds., John Wiley & Sons, 1992.

15

4.1: Identification and Characterization of a HIF-1 α Homologue in *C.elegans*

A tBLASTn enquiry with the HIF- α human sequence was used to identify HIF- α subunit homologues in the *C.elegans* EST database. An EST contig was identified which was identical to an open reading frame (ORF: F38A6.3). This ORF was predicted following determination of the *C.elegans* genome sequence, with the exception of a 104 amino acid amino terminal extension in the latter. No further ESTs or PCR products corresponding to the extension were identified and RACE-PCR products contained a putative *trans* spliced leader sequence. These findings predict that F38A8.3 encodes a 119 amino acid polypeptide that lacks the proposed amino terminal extension.

25

To characterize the regulation of the putative HIF- α homologue (HIF-1), we raised antisera to a recombinant polypeptide and immunoblotted worm extracts. Extracts were prepared from worms exposed to hypoxia, or the cell penetrating iron chelator 2, 2' dipyridyl.

30

Immunoblotting showed a striking induction of HIF-1 by both stimuli. Induction by hypoxia was progressively below 5% oxygen and maximal at the lowest

Gal/549-582/VP16 In vitro transcription translation (IVTT) was prepared using 20 μ l Promega TnT Quick Coupled Retic lysate (Promega, Madison, USA) 1 μ l DNA (1 μ g/ μ l), 2 μ l 1mM desferrioxamine (DFO) and 2 μ l cold methionine (supplied with IVTT kit). For a positive control, the 2 μ l DFO was replaced with 2 μ l 1 mM FeCl₂ (freshly made). The IVTT reaction was incubated at 30°C for 90 min

5 Beads were prepared using 20 μ l gal beads (Santa Cruz no. sc-S10 AC), 5 μ l IVTT, & 100 μ l EBC + 100 μ M DFO and incubated in an End-Over-End rotator for 30-60min. The beads were then spun at 2,000 rpm for 1 minute, the supernatant removed and the beads washed in 1ml of EBC (no EDTA or DFO). This was repeated
10 three times.

The beads were then re-suspended in 1000 μ l HEB (hypotonic extraction buffer: 20mM Tris pH7.5, 5mM KCl, 1.5mM MgCl₂, 1mM dithiothreitol) for each reaction. 100 μ l of the re-suspended beads were transferred into fresh microfuge tubes containing 500 μ l of HEB + DTT. The tubes were spun at 2000rpm for 1 minute and
15 the supernatant removed. Beads were then incubated at room temperature for ten minutes in an end-over-end rotator with a lysate sample under modification conditions as described below then spun at 2000rpm for 1 minute.

Supernatant was removed and the beads washed three times in 500 μ l EBC (50mM Tris pH 7.5, 150mM NaCl, 0.5% v/v Igepal, 0.5mM EDTA) + DFO (In the
20 case of incubation with neat retic lysate, removal of supernatant was facilitated by addition of 500 μ l of EBC + DFO prior to the first spin). The supernatant was then removed from the final wash and the beads used for pull down assays.

VHL Capture Assays

VHL capture or 'pull-down' assays on the Gal/549-582/VP16 beads modified
25 as described above were performed on ice. VHL-HA (T2.1) IVTT performed by mixing 20 μ l Promega TnT Quick Coupled Retic lysate (Promega) with 3 μ l H₂O, 1 μ l DNA (1 μ g/ μ l) and 1 μ l (0.37MBq) ³⁵S-methionine (Amersham Redivue no. AG1094) and incubating at 30°C for 90 minutes. VHL-HA IVTT was then diluted in 100 μ l of EBC + 100 μ M DFO, for each set of beads to be assayed. To the modified, washed
30 gal/ODD/PI6 beads, 100 μ l of the VHL IVTT (T2.1) were added in EBC buffer + 100 μ M DFO.

For VHL capture assays using synthetic biotinylated HIF1- α peptides, peptide was preincubated as indicated for 30min at 37°C, then bound to streptavidin beads, washed, mixed with recombinant VHL or extract, re-captured using beads, and bound VHL analysed by PAGE.

- 5 For HIF-1 α capture assays using 786-0/VHL cell extract, HIF-1 α polypeptides were produced by IVTT, pre-incubated with enzyme, then interacted with cell extract under conditions (10mM Tris pH7.5, 0.25M NaCl, 0.5% NP40, at 4°C) that do not permit modification of HIF1- α (Masson et al.(2001) EMBO), then immunoprecipitated with anti-HA and analysed by PAGE.

- 10 Details of the capture assay protocol are provided below.

HPLC Analyses

- Hydroxylation of the HIF-1 α peptide B19Pro (residues 556-574) was analysed by reverse phase HPLC using a Phenomenex Hypersil 5 μ C18(octadecylsilane) 250 x 4.6mm column and a 5% to 95% acetonitrile gradient in 0.1%TFA at 1 ml/min as the
15 mobile phase. A Gilson HPLC system using 306 pumps and 115 UV detector controlled by Gilson 715 software was used. Standards were unmodified B19Pro and a synthetic peptide (B19Hyp) bearing a hydroxyproline substitution at Pro564.

- Assays were performed with 2.5mM ascorbate, 1.25mM DTT, 50 μ M α KG,
20 1.25mM Fe(II), 25 μ M peptide, 0.66mg/ml catalase, 1.75 mg/ml EGLN2pMAL, in 50mM Tris/HCl, 1.5mM MgCl₂, 5mM KCl. All cofactors were mixed simultaneously by the addition of enzyme to separate drops and incubation was at 37°C for 30 mins. Assays stopped with methanol (70 μ l) and frozen on dry-ice before centrifugation and injection.

- 25 For analysis of hydroxyproline, peptides or proteins were subject to acid hydrolysis, derivatisation with phenylisothiocyanate and HPLC using standard methods.

- Decarboxylation assays were performed using 1-[¹⁴C]-2-oxoglutarate purified polypeptide substrates at approximately 25 μ M, and a purified EGLN2 (PHD1) fusion
30 as described in Mukherji et al. (2001) supra.

On Bead Modification

Mutations were generated using a site directed mutagenesis system (Stratagene). All plasmid sequences were verified by DNA sequencing.

Protein Expression.

³⁵S-labelled or unlabelled proteins were generated in TNT reticulocyte lysate or wheat germ lysate (Promega). Protein expression in insect cells was performed using the Bac-to-Bac/Sf9 system (Gibco BRL). Bacterially expressed proteins were produced in *E. coli* strain BL21 (DE3). Proteins were used in lysates or purified using amylose resin, DEAE-Sepharose, nickel affinity chromatography, or anti-Gal antibodies, as appropriate.

10 Interaction Assays

Assays for interaction between recombinant VHL and HIF polypeptides conformed to the following experimental design. Recombinant VHL and HIF polypeptides were produced separately *in vitro*. The HIF polypeptide was then pre-incubated with extract or a recombinant enzyme as described below, then mixed with VHL and incubated in EBC buffer (50mM Tris pH 7.5, 150mM NaCl, 0.5% v/v Igepal, 0.5mM EDTA) at 4°C for 1 hour, before immunoprecipitation with anti HA antibodies (for HA tagged VHL) or anti Gal antibodies (for Gal4HIF fusions) and analysis by PAGE (Jaakkola et al. (2001) supra).

A schematic of the on bead modification assay is shown in Figure 5.

20 Preincubation of *C.elegans* HIF-1 with worm extract or recombinant EGL-9 was for 30min at 25°C. Preincubation of mammalian HIF- α polypeptides with cell extract or recombinant enzymes was at 37°C for 10-30min unless otherwise stated. For assays of recombinant enzymes, 2-oxoglutarate (2mM), iron (100 μ M), and ascorbate (2mM) were added to the reaction buffer unless otherwise indicated.

25 Reactions performed in hypoxia were in the stated atmospheric oxygen concentration (balance nitrogen) obtained using a controlled environment Invivo₂ 400 hypoxia work-station (Ruskin Technologies) and buffers pre-equilibrated with the appropriate atmosphere. Reactions (50 μ l) were performed in open Eppendorf tubes with mixing and stopped by the addition of 20 volumes desferrioxamine (100 μ M).

30 For peptide blocking experiments peptides (final conc. 1 μ M) were pre-incubated with VHL-1 for 15min before addition to the interaction.

assays were performed as described (Wiesener et al., (1998) Blood 92 2260-2268) using total RNA prepared from a mixed population of worms using Tri-Reagent (Sigma), or total RNA prepared from HeLa cells using RNazolB (Biogenesis).

Plasmid *C. elegans* cDNAs.

5 The *hif-1* cDNA was assembled from 4 overlapping cDNA clones, yk510h7, yk4a2, yk383g1 and yk272d11 (Yuji Kohara, National Institute of Genetics, Japan), and inserted into pcDNA1AMP (Invitrogen). The *vh1-1* cDNA and the cDNA encoding the predicted ORF of T20B3.7 were obtained by RT-PCR of worm RNA and inserted into pcDNA3 (with linkers that encoded an N-terminal HA tag), and
10 pSP72 (Promega) respectively. The *egl-9* cDNA was subcloned into pcDNA1 from yk130h5 (Yuji Kohara). *Phy-1* and *phy-2* cDNAs were subcloned in pCR-Script (Winter and Page, (2000) Mol. Cell. Biol. 20 4084-4093) Gal4/HIF-1 fusion proteins were generated by PCR and inserted into pcDNA3Gal (O'Rourke et al., (1999) J. Biol. Chem. 274 2060-2071).

15 For insect cell expression, sequences encoding Gal4/HIF-1 (289-790) and EGL-9 (1-723) were subcloned into pFastBac1 (Gibco BRL). For bacterial expression, sequences encoding Gal4/HIF-1 (590-790) and EGL-9 (359-723) were subcloned into pET-28a (Novagen), and pMAL-p2X (NEB) respectively.

Mammalian cDNAs

20 The cDNAs encoding the human polypeptides designated EGLN-2 (PHD1), EGLN1 (PHD2), and EGLN3 (PHD3) were obtained by PCR amplification and/or restriction endonuclease digestion from publicly available cDNA banks (The I.M.A.G.E consortium, end NEDO human cDNA sequencing project) or a human colonic cDNA library. Products were ligated into pcDNA3 for expression in
25 reticulocyte lysate IVTTs, or into pMAL-c2X for expression in *E. coli* as maltose binding protein fusions. pPDS15 (Lipscomb et al. (1999) J. Neurochem. 73 429-432) was used for expression of rat SM-20 in reticulocyte lysate IVTT; sequences encoding amino acids 60-355 were subcloned into pTYB11 (NEB) for expression in *E. coli*.

30 For bacterial expression, human HIF-1 α sequences encoding amino acids 344-503 or 530-698 were subcloned into pET28a.

Experimental for Example 4

Materials and Methods

C.elegans Culture, Strains and Extract Preparation.

Worms were cultured using standard methods. Exposure to hypoxia was in
5 bell jars gassed with humidified air or certificated nitrogen / oxygen mixes (British
Oxygen Company). Exposure to 2,2 dipyridyl (200µm), or dimethyl-oxalyglycine (1
mM) was performed during growth in a liquid medium. Wild type worms were
Bristol strain (N2). Mutant strains were obtained from the Caenorhabditis Genetics
Centre and are as indicated in table 5. A deletion mutant in the *vhl-1* gene (ok1610)
10 was generated using trimethylpsoralen. The *vhl-1* strain CB5603 was constructed by
backcrossing ok161 twice against wildtype (N2), then constructing a triple mutant
with markers on either side of *vhl-1* (genotype: *dpy-6* (e2062) *vhl-1* (ok161) *unc--9*
(8101), and then removing these markers by further crosses against N2.

Worm extracts were prepared by homogenisation (Ultraturax T20, IKA
15 Labortechnik) in 4 volumes extraction buffer (100mM NaCl, 1mM EDTA, 50mM
Tris pH7.5, 1% NP-40, 1% sodium deoxycholate) for immunoblotting or in 2
volumes of hypotonic extraction buffer, HEB (20mM Tris pH7.5 5mM KCl, MgCl₂
1mM DTT) for modification reactions.

Mammalian Cells and Extract Preparation

20 HeLa and RCC4 cells were cultured in DMEM. Cell extracts were prepared in
HEB.

Antibodies for Immunoblotting and Immunoprecipitation.

For detection of native *C.elegans* HIF-1 and VHL-1 proteins, antisera were
produced in rabbits immunised with either a glutathione-S-transferase fusion protein
25 expressing amino acids 360-467 of HIF-1, or a maltose binding protein fusion linked
to full length (1-174) VHL-1. Recombinant proteins were expressed in *E.coli*.
Antisera were tested for reactivity using extracts of appropriately transfected Cos7
cells, and purified by ammonium sulphate precipitation. Mouse anti-HA antibody was
12CAS (Roche), and mouse anti-Gal4 antibody was RK5C1 (Santa Cruz).

30 Riboprobes and RNase Protection.

Details of riboprobe templates are provided in table 4. RNase protection

by the von Hippel-Lindau protein. Proceedings of the National Academy of Sciences, USA, 1996. 93: p. 10595-10599.

6. Ohh, M., et al., *Ubiquitination of hypoxia-inducible factor requires direct binding to the beta-domain of the von Hippel-Lindau protein*. Nat Cell Biol, 5 2000. 2(7): p. 423-427.

7. O'Rourke, J.F., et al., *Oxygen-regulated and transactivating domains in endothelial PAS protein 1: comparison with hypoxia inducible factor-1 alpha*. Journal of Biological Chemistry, 1999. 274: p. 2060-2071.

8. Lewis, J.A. and Fleming J.T. *Basic culture methods* (1995) In Methods 10 in Cell Biology, Vol 48 (ed. H.F. Epstein and D.C. Shakes) p.3 Academic Press, San Diego, California.

be ubiquitylated by VHL E3 in S100 extract), this indicates that determinants other than the conserved core residues are important. It is important to understand the key determinants both for oxygen-dependent proline hydroxylation and for subsequent interaction with VHL E3. Particularly since database searches identify "LXXLAP" motifs in a wide variety of cellular proteins.

Although the two sites have functional differences, they both seem to be regulated by the same enzymatic modification. Hydroxylation at proline 564 is the key modification controlling activity of the 3' VHL E3 site. The corresponding proline in the 5' VHL E3 site is also critical for function and polypeptide competition experiments implicate regulatory hydroxylation. Direct evidence of this will come from mass spectrometric analysis. Also of interest is whether the same enzyme is responsible for oxygen-dependent proline hydroxylation at both sites. Sequence differences in the target sites may allow recruitment of different enzymes which in turn may allow graded or cell-type specific differences in the oxygen response. S100 extract was found to be incapable of supporting VHL-dependent ubiquitylation at the 5' site. This may be due to removal of a 5' site-specific enzyme. Alternatively it may be due to removal of a bridging protein proposed to act between the 5' VHL E3 target site and VHL E3. The bridging protein may be an unknown protein or an already identified component of the VHL E3 ligase.

REFERENCES

- Cockman, M.E., *et al.*, *Hypoxia inducible factor- α binding and ubiquitylation by the von Hippel-Lindau tumor suppressor protein*. J Biol Chem, 2000. 275: p. 25733-41.
2. Brenner, S., *The genetics of Caenorhabditis elegans*. Genetics, 1974. 77: p. 71-94.
3. Wood, S.M., *et al.*, *Selection and analysis of a mutant cell line defective in the hypoxia-inducible factor-1 α -subunit (HIF-1 α)*. Journal of Biological Chemistry, 1998. 273: p. 8360-8368.
4. Huang, L.E., *et al.*, *Regulation of hypoxia-inducible factor 1 α is mediated by an oxygen-dependent domain via the ubiquitin-proteasome pathway*. Proceedings of the National Academy of Sciences, USA, 1998. 95: p. 7987-7992.
5. Iliopoulos, O., *et al.*, *Negative regulation of hypoxia-inducible genes*

reaction were then used as substrates in an *in vitro* ubiquitylation assay in the presence or absence of VHL E3 ligase. The 19mer wild-type polypeptide (P) was found to completely ablate the cytoplasmic extract effect. In contrast a polypeptide in which the critical proline is mutated to glycine (P-G) was found to have no effect.

- 5 The 3' VHL E3 target site polypeptide can therefore compete the cytoplasmic extract-dependent modification at the 5' site in a manner dependent upon integrity of proline 564. Pre-hydroxylation of proline 564 rendered the polypeptide unable to compete for modification at the 5' VHL E3 target site presumably because it is no longer a substrate for the enzymatic modification which is occurring at the 5' VHL E3 target
10 site. Thus proline hydroxylation appears to be involved in regulating VHL-dependent ubiquitylation at both the 5' and 3' E3 target sites.

3.9 Discussion.

- Through the use of *in vitro* ubiquitylation assays we have identified 2 independent regions of HIF-1 α targeted by the VHL E3 ligase. Both target sites are
15 located within the ODDD and are functional *in vivo*. Identification of the two VHL E3 target sites is consistent with published data which implied the existence of more than one oxygen-dependent degradation domain within HIF-1 α . Residues 532-585 of HIF-1 α encompassing the 3' VHL E3 target site has previously been shown to be a target for VHL-dependent ubiquitylation. Identification of a second VHL E3 target
20 site provides further evidence of the critical role played by VHL in HIF-1-mediated oxygen-sensing.

- Although HIF-1 α possesses two target sites for VHL E3, they appear to be functionally different. The 3' VHL E3 target site corresponds to the previously identified VHL-binding site. This region of HIF-1 α appears to be targeted directly by
25 VHL acting as the recognition component of the VHL E3 ligase. In contrast we have no evidence that the 5' VHL E3 site can bind VHL directly although it can interact with the complete VHL E3 ligase complex. This may be because the interaction of VHL with the 5' site is indirect or weak compared to the 3' site and difficult to detect by the *in vitro* binding assay used. Both target sites contain a potential consensus
30 motif "LXXLAP" but differ in the sequences surrounding the motif. Since the sites also differ functionally (i.e. in their ability to interact with VHL and their ability to

not direct. After treatment of the Gal-Hif-1 alpha fusion proteins with cytoplasmic extract both the 5' and 3' VHL E3 target sites can be captured by the anti-HA immunoprecipitation. Interaction is not seen when the Gal-Hif-1 alpha fusion protein contains the P402A mutation known to disrupt function of the 5' site.

5 In a previous domain analysis of HIF-1 α the 5' VHL E3 target site was not detected (Ohh *et al*). We wondered whether this was due to the use of S100 extract. VHL-dependent ubiquitylation of both the 5' and 3' E3 target sites was compared using the standard cytoplasmic extract or S100. 35S-methionine labelled GAL-HIF-1 alpha amino acids 344-553 fusion protein and GAL-HIF-1 alpha amino acids 554-698
10 fusion protein substrates were generated by IVTT. Ubiquitylation was performed in fresh cytoplasmic extract, cytoplasmic extract which had been left at 4 degrees C for 4 hours or the S100 supernatant of cytoplasmic extract from RCC4 cells, which lack pVHL or RCC4 cells stably transfected with pcDNA3 VHL.HA. The S100 extracts clearly enabled VHL dependent ubiquitylation of GAL-HIF-1 alpha amino acids 554-
15 698 fusion protein but not GAL-HIF-1 alpha amino acids 344-553 fusion protein. A factor specifically required for recognition of the 5' VHL E3 target site is either lost or inactivated during S100 preparation.

3.8 The 5' VHL E3 target site is also regulated by proline hydroxylation.

It has been shown above that the 3' VHL E3 target site responds to oxygen
20 level via hydroxylation at proline residue 564. This proline residue forms part of a potential motif conserved between the 5' and 3' target sites. Mutation of the corresponding proline residue (P402A) in the 5' target site also results in functional inactivation. It was possible therefore that proline residue 402 was also a target for regulatory hydroxylation. To test this we asked whether polypeptides corresponding
25 to the 3' VHL E3 target site could interfere with the cytoplasmic extract-dependent modification of the 5' VHL E3 target site. 35S-methionine labelled GAL-HIF-1 alpha amino acids 344-553 fusion protein substrate was generated by IVTT and incubated *in vitro* in buffer or cytoplasmic extracts from RCC4 cells in the presence of wild-type 19mer peptide representing HIF-1 alpha amino acids 556-574 (12.5 μ M); a
30 polypeptide where the critical proline is mutated to glycine (P564G); or a polypeptide where the proline is modified to a hydroxy-proline (P-OH). The products of this

1 α wild type and mutant substrates were generated by IVTT and subjected to *in vitro*
ubiquitylation in cytoplasmic extracts from RCC4 cells, which lack pVHL (RCC4),
or RCC4 cells stably transfected with pcDNA3 VHL.HA (RCC4/VHL) in the
presence or absence of exogenous ubiquitin. The double mutant P402A+P564G was
5 found to show no VHL dependent ubiquitylation, but isolated mutations of the critical
prolines at each individual VHL E3 target site did not ablate ubiquitylation. Thus
when these mutations are introduced individually the mutant HIF-1 α proteins still
remain targets for VHL-dependent ubiquitylation (presumably because each retains
an active VHL E3 target site). HIF-1 α therefore appears to contain two, and only two
10 target sites for VHL-dependent ubiquitylation. To assay importance *in vivo*, the single
and double VHL E3 target site mutants were transfected into the HIF-1 α deficient cell
line KA13 (Wood *et al*) and tested for their ability to mediate oxygen-dependent
transcriptional regulation (Fig. 4). The transfected wild-type HIF-1 α protein displayed
oxygen-dependent regulation. However the P 402 A, and P 564G point mutants were
15 transcriptionally active under normoxic conditions and showed very little
upregulation in hypoxia (Fig. 4). The P 402 A + P 564G double mutant was
essentially constitutive under normoxic conditions (Fig. 4).

3.7 The 5' and 3' VHL E3 target sites differ in their functional requirements.

The ability of pVHL to interact directly with both the 5' and 3' E3 target sites
20 was tested *in vitro*. The 35S-methionine labelled GAL-HIF-1 α fusion proteins
GAL 344-553 P402A, GAL 344-553, GAL 652-826, GAL 554-698 were made by
IVTT, incubated in buffer or cell extract from RCC4 cells lacking pVHL at 30
degrees C for 1 hour. Samples were then cooled and incubated with extract from 786-
0 cells stably transfected with pcDNA3 VHL.HA for 90 minutes on ice prior to
25 immunoprecipitation with anti-HA antibodies and protein G beads. Input samples of
the GAL-HIF-1 α fusion proteins and retrieved immunoprecipitates were
analysed by SDS/PAGE and autoradiography. The 3' VHL E3 target site is already
known to bind VHL in an *in vitro* interaction assay (Cockman *et al*) and the results
obtained confirmed this. In contrast the 5' VHL E3 target site (represented by GAL
30 344-553) does not appear to bind pVHL in this assay. Either the interaction of pVHL
with the 5' E3 target site is transient and too weak to be detected, or the interaction is

(GAL 344-400 and GAL 344-379), whereas deletions made at the N-terminus retained activity. The minimal functional domain defined by this analysis was HIF-1 α residues 380-417. Although the output ubiquitylation signal was reduced, GAL 380-417 was still a target for VHL-dependent ubiquitylation and still displayed enhanced ubiquitylation following cytoplasmic extract pre-treatment.

3.5 Identification of a potential functional motif conserved between the 5' and 3' VHL E3 target sites.

The HIF-1 α 380-417 sequence was analysed in an attempt to identify residues critical to the functional effect. The HIF-1 α sequence was aligned with the corresponding region of HIF-2 α and the VHL-binding site. Within the VHL-binding site, hydroxylation at proline 564 is identified in Example 1 above as a key regulatory event. Interestingly, a potential conserved motif encompassing this proline can be identified between the two VHL E3 ligase target sites (Fig. 3A). Mutations of this potential motif were assayed in the context of GAL 344-417. 35S-methionine labelled GAL-HIF-1 alpha amino acids 344-417 wild type and mutant substrates (comprising the mutation P402A or the double mutation LL397, 400A) were generated by IVTT, incubated in buffer or cytoplasmic extract prior to *in vitro* ubiquitylation in the presence or absence of the VHL E3 ligase. The double mutation of leucines 397 and 400 to alanine (LL 397,400 AA) was found to ablate VHL-dependent ubiquitylation. The point mutation of proline 402 to alanine (P 402 A) also ablated VHL-dependent ubiquitylation.

Mutations of the 344-417 region were then tested for their effects on oxygen-dependent regulation *in vivo*. The HIF-1 α 344-417 region is known to confer oxygen-dependent regulation on a GAL-VP16 fusion (O'Rourke). The C-terminal deletion (344-400) and the P 402 A mutation were tested in this context and both were found to abolish oxygen-dependent regulation *in vivo* (Fig. 3B).

3.6 Identification of critical point mutations.

Identification of critical point mutations allows these two VHL E3 target sites to be assayed within the full-length HIF-1 α molecule. The P 402 A mutation was introduced to ablate activity of the 5' VHL E3 target site and the P 564 G mutation to ablate activity of the 3' VHL E3 target site. 35S-methionine labelled full length HIF-

E3 ligase. Iron was found to enhance the ubiquitylation of Gal-HIF-1 α 344-417 fusions in the presence of cytoplasmic extract. Cytoplasmic extract enhanced the ubiquitylation of Gal-HIF-1 α 554-698 although the effect was less pronounced than that of GAL344-417. These data suggested that the two independent VHL E3
5 ligase target sites may be regulated by a similar mechanism.

3.4 Mapping of 380-417 as a minimal domain targeted by cytoplasmic extract and VHL E3.

To begin to understand the mechanism it was necessary to define a minimal functional domain. Residues 344-553 correspond to exons 9-11 of HIF-1 α and so an
10 exon-based deletional strategy was used. 35S-methionine labelled GAL-HIF-1 α amino acids 344-553 fusion protein substrate was generated by IVTT, incubated in buffer or cytoplasmic extract prior to *in vitro* ubiquitylation in the purified component assay in the presence or absence of the VHL E3 ligase. The GAL 344-503 fusion (corresponding to exons 9 and 10 of HIF-1 α) still displayed enhanced VHL-
15 dependent ubiquitylation following cytoplasmic extract pre-treatment. Exons 9 and 10 were then assayed individually by generating fusions carrying GAL-HIF-1 α amino acid residues 344 to 503, 344 to 417 or 418 to 503. The only fusion which was not ubiquitylated was that carrying residues 418 to 503. Thus both ubiquitylation and the cytoplasmic extract effect were found to localise to exon 9, represented by GAL 344-
20 417. VHL dependent extract enhanced ubiquitylation therefore clearly depends on HIF-1 α amino acids 344-417.

The corresponding exon in HIF-2 α was then assayed. The substrates used were Gal-HIF-2 α fusion comprising amino acids 344-417 or 345-416. Residues 345-416 of HIF-2 α were also found to be a target for VHL-dependent ubiquitylation and
25 also exhibited enhanced ubiquitylation following cytoplasmic extract pre-treatment. The function of this region is therefore conserved between HIF-1 α and HIF-2 α and sequence comparisons will help to identify critical residues.

Deletional analysis was further extended to screen the HIF-1 α 344-417 region. Gal-HIF-1 α fusions comprising amino acids 344 to 417, 344 to 400, 344 to
30 379, 360 to 417 or 380 to 417 of HIF-1 α were individually assessed as above. Deletions made at the C-terminus completely ablated VHL-dependent ubiquitylation

converting the GAL 344-553 substrate into a strong target for VHL-dependent ubiquitylation. Accompanying this effect a marked mobility shift of the GAL 344-553 substrate was seen due to phosphorylation in the cytoplasmic extract.

Phosphorylation is known to play an important role in regulating recognition
5 of substrates by the SCF E3 ligase. HIF-1 α is known to be a phosphoprotein, although an oxygen-dependent phosphorylation event has not been identified. A potential link between HIF-1 α phosphorylation and ubiquitylation was therefore of interest. Pre-incubation of the GAL 344-553 substrate with nuclear extract also resulted in a phosphorylation-induced mobility shift, but this was not accompanied by
10 increased VHL-dependent ubiquitylation.

To clarify the role of phosphorylation in the cytoplasmic extract effect, hexokinase treatment was used. 35S-methionine labelled GAL-HIF-1 alpha amino acids 344-553 fusion protein substrate was generated by IVTT, incubated in buffer, cytoplasmic extract, cytoplasmic extract that had been depleted of ATP by pre-
15 incubation with hexokinase or cytoplasmic extract which had been heat denatured. Enhanced VHL dependent substrate ubiquitylation was found to persist in the absence of ATP (and consequent absence of phosphorylation) but not following heat denaturation. As the ATP-depleted extract can no longer support GAL 344-553 phosphorylation but is still capable of supporting enhanced VHL-dependent
20 ubiquitylation, phosphorylation of GAL 344-553 is therefore not the key event mediating interaction with VHL E3. As heat-treated cytoplasmic extract was unable to support enhanced VHL-dependent ubiquitylation this suggests that a protein factor may be involved either in binding to, or modifying the GAL 344-553 substrate.

The demonstration in Example 1 above that interaction of VHL with the VHL
25 binding site in HIF-1 α is promoted by cytoplasmic extract and iron led us to test the effect of cytoplasmic extract on ubiquitylation of the GAL 554-698 substrate and to test the effect of iron on ubiquitylation of GAL 344-417. 35S-methionine labelled GAL-HIF-1 α fusions comprising amino acids 344-417 or 554-698 of HIF-1 α substrates were generated by IVTT, incubated in buffer, cytoplasmic extract,
30 cytoplasmic extract supplemented with 100 μ M iron chloride prior to *in vitro* ubiquitylation in the purified component assay in the presence or absence of the VHL

correspond to ubiquitylated forms of GAL344-698 as their production is E1-, E2- and VHL E3-dependent.

In vitro ubiquitylation was then performed on a variety of GAL-HIF-1 α fusions. 35S-methionine labelled immunopurified GAL-HIF-1 α fusions comprising amino acid residues 344 to 698, 344 to 553, 554 to 698 or 652 to 826 of HIF-1 α were used as substrates using reaction mixtures containing E1, E2, VHL E3 ligase, ubiquitin and ATP or reaction mixtures where ubiquitin or VHL E3 ligase were omitted to act as controls. VHL E3 ligase dependent ubiquitylation was clearly seen when the substrate contained HIF-1 α amino acids 344-698, 344-553 or 554-698 but not for substrates containing residues 652-826 of HIF-1 α . The fusion containing residues 652-826 of HIF-1 α acted as a control as residues 652-826 of HIF-1 α show no oxygen-dependent regulation at the protein level *in vivo* (O'Rourke) and do not interact with pVHL *in vitro* (Cockman *et al*). As the GAL 344-553 and GAL 554-698 substrates were both found to be targets for the VHL E3 but GAL652-826 are not, the results obtained using the purified component assay concurs with the cytoplasmic extract assay in identifying HIF-1 α residues 344-553 and 554-698 as independent VHL E3 targets *in vitro*.

3.3 Cytoplasmic extract enhances functional interaction of the VHL E3 ligase with the 5' target site in HIF-1 α .

It was noted however that VHL-dependent ubiquitylation of the GAL 344-553 substrate differed greatly between the two assays. In the cytoplasmic extract assay GAL 344-553 is a much better substrate for VHL-dependent ubiquitylation than GAL 554-698, but in the purified component assay the position is reversed with GAL 344-553 an extremely weak substrate. We wondered whether cytoplasmic extract was important for recognition of the 344-553 region by VHL E3. To test this 35S-methionine labelled GAL 344-553 substrate was generated by IVTT, incubated in buffer, cytoplasmic extract or nuclear extract prior to *in vitro* ubiquitylation in the purified component assay in the presence or absence of the VHL E3 ligase. Treatment with cytoplasmic extract dramatically enhanced the VHL dependent ubiquitylation of the substrate. Thus, whilst the buffer-treated substrate remains an extremely weak target for VHL E3, pre-treatment with cytoplasmic extract has a dramatic effect,

for Western blotting were prepared by resuspending cell pellets in 7M urea, 10% glycerol, 1% SDS, 10mM Tris pH6.8, containing 50 μ M phenylmethylsulfonyl fluoride and leupeptin, pepstatin and aprotinin all at 0.1 μ g/ml, followed by disruption using a hand-held homogenizer (Ultra-Turrax T8 with 5G dispersing tool; Janke & Kunkel GmbH). Following SDS-PAGE, proteins were transferred onto Immobilon-P membrane (Millipore) and processed for western blotting using the indicated antibody.

3.1 The VHL E3 ligase can interact functionally with two distinct regions of the HIF-1 α ODDD in vitro.

In order to understand more about the interactions of pVHL with HIF-1 α we analysed VHL-dependent ubiquitylation of the HIF-1 α ODDD in an *in vitro* assay using cytoplasmic extracts as a source of ubiquitylation enzymes. 35S-methionine labelled GAL-HIF-1 alpha fusion proteins containing the amino acids 344 to 698, 344 to 553, 554 to 698 or 504 to 554 of HIF-1 α were generated by IVTT and subjected to *in vitro* ubiquitylation in cytoplasmic extracts from RCC4 cells, which lack pVHL (RCC4), or RCC4 cells stably transfected with pcDNA3 VHL.HA (RCC4/VHL) in the presence or absence of exogenous ubiquitin. PVHL dependent ubiquitylation, resulting in a strong signal of decreased mobility at the top of the lane, was clearly observed when the substrate contained HIF-1 alpha amino acids 344-698, 344-553 and 554-698, but not amino acids 504-554. HIF- α residues 344-553 and 554-698 are both capable of oxygen-dependent regulation *in vivo* (O'Rourke *et al*) and when analysed *in vitro* here both regions exhibit VHL-dependent ubiquitylation. This indicates that the VHL E3 ligase can interact functionally with at least two sites in HIF-1 α .

3.2 Requirements for functional interactions.

To investigate this further, it was necessary to develop a ubiquitylation assay using purified components. 35S-methionine labelled GAL-HIF-1 α amino acids 344-698 fusion protein was generated by IVTT, immunopurified with anti-Gal antibody conjugated agarose and subjected to *in vitro* ubiquitylation with purified components. This resulted in the production of high molecular weight GAL344-698-related species in a ubiquitin and ATP-dependent manner. These high molecular weight species

Affiniti Research (Exeter, UK). VHL E3 was obtained by anti-HA immunoprecipitation from stably transfected 7860-VHLHA cell lysates (Iliopoulos *et al*). GAL-HIF-1 α substrate was prepared by anti-GAL immunoprecipitation from [³⁵S] methionine-labeled TnT rabbit reticulocyte (Promega) translates. Each 40 μ l
5 ubiquitylation reaction consisted of 4 μ l of 5mg/ml ubiquitin, 4 μ l of 10xATP regenerating system (20mM Tris pH7.5, 10mM ATP, 10mM magnesium acetate, 300mM creatine phosphate, 0.5mg/ml creatine phosphokinase), 2 μ l E1, 3 μ l E2, 6 μ l VHL E3 immunopurified on protein G sepharose, 6 μ l GAL-HIF-1 α substrate immunopurified on agarose beads. Reactions were incubated at 30°C for 2h with
10 occasional mixing, stopped by the addition of SDS sample buffer and analysed by SDS-PAGE and autoradiography. Cytoplasmic extract-based ubiquitylation assays have been previously described (Cockman).

In vitro interaction assays - TnT rabbit reticulocyte (Promega) translates (4 μ l [³⁵S] methionine-labeled) were mixed either in 70 μ l hypotonic extraction buffer (20mM
15 Tris pH7.5; 5mMKCl; 1.5mMMgCl₂; 1mM DTT) or RCC4 cytoplasmic extract at 30 degrees C for 1 hour. Samples were then cooled and incubated with 400 μ l extract from 786-0 cells stably transfected with pcDNA3 VHL.HA for 90 minutes on ice prior to immunoprecipitation with excess anti-HA antibodies and protein G beads. Input samples of the GAL-HIF-1 alpha fusion proteins and retrieved
20 immunoprecipitates were analysed by SDS/PAGE and autoradiography.

Luciferase and beta-Galactosidase Assays - Luciferase activities in cell extracts were determined using a commercially available luciferase assay system (Promega) and a TD-20e luminometer (Turner Designs). Relative beta-galactosidase activity in
extracts were measured using o-nitrophenyl-beta-D-galactopyranoside (0.67 mg/ml)
25 as substrate in a 0.1 M phosphate buffer (pH 7.0) containing 10mM KCl, 1mM MgSO₄ and 30mM beta-mercaptoethanol incubated at 30°C for 15-45 min. The A₄₂₀ was determined after stopping the reaction by the addition of 0.4M sodium carbonate (final concentration).

Cell extract preparation and Western Blotting - Cytoplasmic extract for ubiquitylation
30 assays was prepared as previously described (Cockman *et al*). S100 extract was obtained by an additional ultracentrifugation step at 100,000g at 4°C for 4h. Extracts

Plasmid constructs - His₆-E1 -tagged mouse E1 cDNA in pRSET was a kind gift of T.Hunt. pcDNA3-VHLHA has been previously described Cockman *et al.* pGAL 344-417VP16 has been previously described (O'Rourke). Plasmids bearing mutations were generated using a site-directed mutagenesis kit (QuickChange; Stratagene) and
5 mutagenic oligonucleotides designed according to the manufacturer's recommendations. All PCRs were performed using *pfu* DNA polymerase (Stratagene).

Cell culture and transient transfection - 7860, U2OS and RCC4 cells were maintained in Dulbecco's modified Eagle's medium supplemented with 10% fetal
10 calf serum, glutamine (2mM), penicillin (50 IU/ml) and streptomycin sulfate (50µg/ml). Ka13 cells (Wood *et al*) were grown in Ham's F12 medium with the same supplements.

Cell extract preparation and Western Blotting - Cytoplasmic extract for ubiquitylation assays was prepared as previously described (Cockman *et al*). S100
15 extract was obtained by an additional ultracentrifugation step at 100,000g at 4°C for 4h. Extracts for Western blotting were prepared by resuspending cell pellets in 7M urea, 10% glycerol, 1% SDS, 10mM Tris pH6.8, containing 50 µM phenylmethylsulfonyl fluoride and leupeptin, pepstatin and aprotinin all at 0.1 µg/ml, followed by disruption using a hand-held homogenizer
20 (Ultra-Turrax T8 with 5G dispersing tool; Janke & Kunkel GmbH). Following SDS-PAGE, proteins were transferred onto Immobilon-P membrane (Millipore) and processed for western blotting using the indicated antibody.

Antibodies - Anti-HA antibody (12CA5) was from Roche Molecular Biochemicals, anti-GAL4(DBD) (RK5C1) agarose conjugate from Santa Cruz Biotechnology and
25 anti-HIF-1α clone 54) antibody from Transduction Laboratories.

Ubiquitylation enzymes and assays - The E1 activating enzyme used in ubiquitylation assays was either obtained from Affiniti Research (Exeter, UK) or purified from BL21 (DE3) *E.coli* transfected with plasmid expressing His₆-tagged mouse E1. His₆-E1 was purified by Ni²⁺-agarose affinity chromatography. After
30 dialysis against phosphate buffered saline, glycerol was added to 10% (vol/vol) and 25ng/µl aliquots stored at -80°C. Human CDC34 recombinant E2 enzyme was from

In this Example we show that two independent regions of the HIF-1 α ODDD are targeted for ubiquitylation by VHL E3 in a manner dependent upon proline hydroxylation. However these two VHL E3 target sites differ in their overall sequence, their ability to bind VHL directly and their requirement for other cellular factors. These data reinforce the critical role for pVHL in HIF- α regulation, but
5 implicate a more complex model for pVHL/HIF- α interactions.

Immunoprecipitation and band shift assays show that VHL and HIF- α subunits are physically associated in a wide range of cell types, consistent with a general role for VHL in oxygen-dependent regulation of HIF- α subunits. At the same
10 time biochemical studies show that VHL exists as a multiprotein complex with elongins B and C, CUL-2 and RBX1. This complex is homologous to the SCF (Skp-1-Cdc53/Cullin-F-box) class of E3 ubiquitin ligases. Like SCF E3, the VHL complex has inherent ubiquitin ligase activity. VHL itself is thought to play a role analogous to the F-box substrate recognition component. HIF- α subunits are therefore clear
15 candidate substrates for VHL E3 and have since been shown to be ubiquitylated in a VHL-dependent manner *in vitro*.

Example 1 above demonstrates that degradation of HIF-1 α mediated by the VHL binding site occurs through oxygen-dependent hydroxylation at proline 564. It is currently unclear whether oxygen-dependent degradation of HIF- α subunits is
20 solely VHL-dependent. In renal cell carcinoma lines and in CHO cells VHL appears to be the critical mediator. However only one VHL binding site has been identified in HIF-1 α and regions outside this site can confer oxygen-dependent regulation *in vivo*. To investigate the mechanisms underlying this we have employed *in vitro* ubiquitylation assays which provide evidence of functional interaction with the VHL
25 E3 ligase. We find that two independent regions of the HIF-1 α ODDD are targeted for ubiquitylation by VHL E3 in a manner dependent upon proline hydroxylation. However these two VHL E3 target sites differ in their overall sequence, their ability to bind VHL directly and their requirement for other cellular factors. These data reinforce the critical role for pVHL in HIF- α regulation, but implicate a more
30 complex model for pVHL/HIF- α interactions.

Materials and methods

Protein expression and interaction assays. 35S labelled proteins were generated in reticulocyte lysates (Promega) programmed with plamids encoding wild type ceHIF (pcDNA1cehif), mutant ceHIF (pcDNA1cehif.P621G) or c-terminal HA tagged ceVHL (pcDNA3ceVHL-HA). pcDNA1cehif.PxxxG was generated from
5 pcDNA1cehif using a site directed mutagenesis system (Stratagene) and the following forward and reverse primers:
5'GATTTATCGTGCTTGGCAGGATTCGTTGACACTTATG (forward)
5'GTGTCAACGAATCCTGCCAAGCACGATAAATCAGGC (reverse).
pcDNA3ceVHL.HA was obtained by RT-PCR amplification of nucleotides 1 to 525
10 of the predicted ORF of sequence F08G12.4 from *C.elegans* RNA, and exchange for human VHL sequence in pcDNA3-VHL.HA. For interaction assays 1µl of each programmed lysate was mixed in EBC buffer at 4°C for 1hr before anti-HA immunoprecipitation as described in Cockman *et al.* Pretreatment of ceHIF with worm extract was for 30min at 25°C with 10µl of extract derived by hypotonic
15 extraction of a worm homogenate in 20mM Tris pH7.5, 5mMKCl, 1.5MgCl₂, 1mMDTT.

Worm strains and experimental conditions *C. elegans* strains were cultured as described by Brenner[Brenner, 1974 #1]. Exposure to hypoxia was in bell jars gassed with humidified air or certificated nitrogen/ oxygen mixes (British
20 Oxygen Company). Exposure to iron chelators worms was by growth in a liquid medium as described previously [Lewis, 1997 #2] with or without 200 µM 2,2 Dipyridyl. Wild type worms were Bristol strain (N2). ok161 was generated by Dr. Robert Barstead, Oklahoma Medical Foundataion, using ultraviolet and psoralen mediated utagenesis. PCR using oligonucleotides from the flanking genomic
25 sequence was used to select worms bearing a deletion at the FO8G12.4 (vhl) locus. Confounding mutations in ok161 were removed by backcross selection using visible markers that flank the VHL locus (dpy-6 unc-9).

Example 3: The VHL E3 Ligase Complex Interacts with Two Independent Regions of
30 HIF-1α

In mammalian systems the HIF/pVHL system has important functions in the regulation of oxygen delivery through effects on angiogenesis, vasomotor control and erythropoiesis. Conservation in *c. elegans* indicates that the HIF/pVHL system of oxygen regulated gene expression antedates the development of these complex oxygen delivery systems and that the system must have a critical function in other responses to oxygen availability. The effects observed already on the expression of metabolic enzymes may provide clues to such functions. However though the viability of both *vhl* mutant and *hif* mutant worms in the laboratory suggests that the critical functions that have directed the evolution of this system are likely to be observed under other, presumably more stressful, conditions.

2.5: Methods

Identification of *c.elegans hif* cDNA. *C.elegans* EST database searches were performed using the tBLASTn programme and the human HIF-1 α sequence as a probe. The putative *c.elegans hif* cDNA was assembled from 4 overlapping cDNA clones, yk510h7, yk4a2, yk383g1, and yk272d11 (kindly provided by Yuji Kohara, National Institute of Genetics, Mishima, Japan), and inserted into the polylinker of pcDNA1AMP (invitrogen) to create pcDNA1*cehif* using standard methods.

Antibody Generation and Immunoblotting. DNA encoding amino acids 360 to 497 of ceHIF was inserted into pGEX-4t-1 and the corresponding GST/ceHIF fusion protein was expressed in *E. coli*. The protein was purified using glutathione agarose and used to raise antisera in rabbits. Antisera were tested for reactivity using extracts of Cos7 cells transfected with pcDNA1*cehif*, and purified by ammonium sulphate precipitation. Worm extracts used in immunoblotting were prepared from washed worms by homogenisation in 4 volumes extraction buffer (150mM NaCl, 1mM EDTA, 50mM Tris pH 7.5, 1% NP-40 1% sodium deoxycholate) using an Ultraturax T20 homogeniser.

Riboprobes and RNase Protection Riboprobe templates were generated from total *c.elegans* RNA using RT-PCR. Details of the primers, and sequences are provided in supplementary information. RNase protection assays were performed as described in (ref) using 10-50mg total RNA prepared from a mixed population of worms using Tri-Reagent (Sigma).

Clear induction by hypoxia was observed for mRNA encoding lactate dehydrogenase-A and an isoform of carbonic anhydrase, and in each case the mRNA was strikingly upregulated in *vhl* worms.

Second we tested for induction by hypoxia among a subset of pVHL dependent differentially expressed gene defined by array screening. Of eight genes
5 demonstrated by RNase to be upregulated in *vhl* worms five were strongly inducible by hypoxia in wild type worms.

Oxygen homeostasis is a fundamental physiological problem in all organisms that can live in an aerobic environment, and genetic studies in bacteria and yeast have
10 defined specific sensing systems that regulate gene expression in accordance with oxygen availability. However efforts to link these systems to responses in mammalian cells have so far been unsuccessful, and database analysis has not reveal a HIF homologue in the *s. cerevisiae* genome or sequenced prokaryotic genomes. The current work therefore provides the clearest analysis to date of homology with a
15 primitive organism that has been developed for genetic analysis. Given recent advances in large scale analysis of gene expression gene function in *c.elegans* the findings provide important new opportunities to understand cellular responses to oxygen availability.

In mammalian cells transcriptional activation of HIF is believed to be a multi-
20 step process involving separate regulatory steps in nuclear localization, DNA binding, and co-activator recruitment as well as different systems of ubiquitin mediated proteolysis. Somewhat surprisingly, in VHL defective renal carcinoma cell lines HIF- α subunits are constitutively stabilised and hypoxia inducible mRNAs are constitutively upregulated in normoxic cells, indicating that at least in this cell
25 background pVHL has a dominant non-redundant function in the regulation of the HIF transcriptional response. Both ceHIF protein and its transcriptional target mRNAs also showed striking up-regulation in normoxic *vhl* mutant worms. Importantly this indicates that a critical non-redundant function of VHL in regulation of HIF extends outside the cell background of VHL associated tumours, and most
30 likely operates generally in higher eukaryotes.

Table A

Sequence Name	Gene Description	Regulated by Hypoxia	Regulated by VHL
5 F13D12.2	Lactate de hydrogenase	+	+
F54D8.4	Putative carbonic anhydrase	-	-
T28F2.3	Putative carbonic anhydrase	-	-
R01E6.3	Putative carbonic anhydrase, strong similarity to human CA2	+	+
R173.1	Putative carbonic anhydrase	-	-
10 K05G3.3	Putative carbonic anhydrase, strong similarity to human CA7	-	-
B0412.2	daf-7/ member of the TGF β superfamily	-	-
C14F5.1	nip 3/ bcl-2	-	-
B0432.5	Putative tyrosine hydroxylase	-	-

15 Table B

Sequence Name	Gene Description	Regulated by Hypoxia	Regulated by VHL
F22B5.4	Protein of unknown function	+	+
F35G2.4	Prolyl 4-hydroxylase alpha subunit	+	+
20 C55B7.4	Member of the acyl-CoA dehydrogenase protein family	+	+
K09E4.4	Strong similarity to human alpha T-acetylglucosaminidase	-	-
T05B4.2	Member of the nuclear hormone receptor/Zinc finger protein family	+	+
H14N18.4	Member of the gamma-glutamyltransferase (tentative) protein family	-	-
25 C16C10.3	<u>Piwi related protein</u>	+	+

extracts were made from wild type worms grown in normoxic (21% oxygen) conditions in the presence and absence of dimethyloxallylglycine (1mM).

Immunoblots for *ceHIF* were performed after separation on SDS/PAGE. The band representing *ceHIF* was identified and it could clearly be seen that inhibitor treatment
5 clearly results in a substantial increase in the amount of immunodetectable *ceHIF* in normoxia.

2.4: Expression of HIF target genes.

We wished to test directly for effects of the HIF/pVHL system on patterns of gene expression in *c.elegans*. First we tested for hypoxia inducible expression
10 amongst a set of *c.elegans* homologues of mammalian genes that are known HIF targets, and compared the upregulation of mRNA upon hypoxic exposure of wild type worms with that observed in the *vhl* mutant worm. The results obtained are shown in Tables A and B below.

Table A summarises results for a subset of genes selected for analysis on the
15 basis of putative homology to mammalian HIF target genes and tested for regulation by hypoxia and VHL in *c.elegans*. Table B summarises results for a subset of genes confirmed as regulated by *vhl* by RNase protection after identification in comparative array screening of wild type and *vhl* mutant worms, and subsequently tested for regulation by hypoxia. The full gene array dataset from which these genes were
20 identified are available at <http://genome-www4.stanford.edu/cgi-bin/SMD/login.pl>.

(21% oxygen) or hypoxic (0.1% oxygen) gas mixtures for 6 hrs. As expected the *vhl* mutant worms were found to have high levels of *ceHIF* protein regardless of oxygen tension. The mutants *daf-18* (encoding a PTEN homologue), *daf-2* (encoding an insulin receptor homologue) and *age-1* (encoding a PI3-kinase homologue), in contrast with the *vhl* mutant worms, all showed regulation of *ceHIF* by oxygen that was similar to wild type. The other mutants screened for effects on *ceHIF* were selected on the basis of known effects on ROS metabolism, or altered phenotypic sensitivity to oxidant stress included several mutants affecting mitochondrial proteins (*mev-1*, *clk-1*, *gas-1*), a *ctl-1* mutant that affects cytosolic catalase activity and others (*mev-2*, *mev-3*) where the product is not yet characterized and again regulation was similar to wild type suggesting that this a distinct oxygen sensing system that in *C. elegans* is not tightly linked to general systems of oxidant defence.

In view of the data presented here demonstrating a critical role for enzymatic hydroxylation of prolyl residues within HIF in its normoxic recognition by pVHL and subsequent ubiquitylation and destruction by the proteasome in the mammalian system we also tested worms bearing mutations in known prolyl hydroxylases (*dpy-18* and *phy-2*), and genes containing sequence motifs compatible with a function as a prolyl hydroxylase (*egl-9* – located at F22E12.4). The effect of prolyl hydroxylase mutants on HIF activity was studied by blotting. Extracts were made from wild type and mutant worms grown in normoxic (21% oxygen) and hypoxic (0.1% oxygen) conditions. Immunoblots for *ceHIF* were performed after separation on SDS/PAGE. The band representing *ceHIF* was identified. No detectable *ceHIF* was seen in an extract from normoxic wild type worms. In contrast in normoxic extracts from *egl-9* deficient worms *ceHIF* is easily detected (allele MT 1201; allele MT 1216 grown at 25 degrees C), at levels comparable with those seen in extracts from these strains grown in hypoxic conditions. As the *egl-9* deficient worms have high normoxic levels of *ceHIF*, this suggests that this gene product is involved in the normal degradation of *ceHIF*. The *dpy-18* and *phy-2* deficient worms showed normal *ceHIF* levels.

We also used dimethyloxalylglycine (a cell permeant alpha ketoglutarate analogue known to block this family of dioxygenases) and demonstrated an increased abundance of *ceHIF* in normoxia in the presence of the inhibitor. In these experiments

mammalian interaction and replaced it with glycine. Whilst wild type ceHIF could interact with tagged pVHL, the ceHIF Pro621-Gly mutant form was unable to interact with pVHL mirroring the findings with mammalian HIF.

To pursue the functional importance of the interaction between ceVHL and ceHIF, we next employed a viable homozygous deletion mutant worm lacking
5 ceHIF, and assayed worm extracts for ceHIF by immunoblotting. In normoxic ceVHL worms ceHIF levels were strikingly upregulated and were essentially unregulated by oxygen, being similar in hyperoxia (80% O₂), air, and hypoxia (0.1% O₂). Thus a critical function for pVHL in the response to oxygen appears also to be
10 conserved. Surprisingly, ceVHL deficient worms are phenotypically relatively normal, with only slightly slower growth rates and mildly reduced reproductive capacity compared to wild type.

This tight conservation of the HIF/pVHL system indicates that *C. elegans* provides a new model for analysis of the oxygen sensing and signalling pathways that
15 regulate HIF, and for the analysis of downstream effects on patterns of gene expression. As a first step in exploring this potential we assessed ceHIF induction by hypoxia in a mutant worms selected to test candidate molecules in the sensing/signalling pathway. A number of studies support the involvement of oxygen radicals though the source and mode of interaction with the HIF/pVHL complex is
20 unclear. Other studies have suggested the involvement of particular growth factor signalling pathways in HIF regulation, but the relation of these findings to the oxygen sensitive signal is uncertain. In one line of investigation it has been found that insulin and insulin-like growth factors can activate HIF in normoxic cells, that the tumour suppressor PTEN acts as a negative regulator of HIF, and that the downstream target
25 of PTEN, Akt shows oxygen dependent phosphorylation, suggesting the involvement of an insulin receptor/PI3-kinase pathway in HIF regulation. This pathway is conserved in *C. elegans*. and interestingly has been implicated in ROS metabolism. We therefore tested several mutants to determine their effect on the interaction of ceHIF with VHL.

30 The level of ceHIF in wild type and a series of mutant worms was determined by immunoblotting. Worms were grown on plates in bell jars flushed with normoxic

elegans. Worms were cultured in liquid media in the presence or absence of the penetrant bidentate iron chelator 2', 2' dipyridyl (200 μ M) for 6 or 16 hrs. A striking Induction of *ceHIF* by iron chelation was observed at both 6 and 16 hours and the level of induction was equivalent to that observed in severe hypoxia. *ceHIF* was not induced in the absence of iron chelation.

2.3 Conserved role for VHL.

Regulation of mammalian HIF- α subunit protein levels occurs through a one or more systems of ubiquitin mediated, oxygen regulated proteolysis. To date the most clearly defined of these involves the von Hippel-Lindau tumour suppressor protein (pVHL), which physically interacts with specific HIF- α residues, acting as the recognition component of an E3 ubiquitin ligase. In VHL defective renal carcinoma cells HIF- α subunits are constitutively stabilised leading to greatly increased steady-state levels in normoxia. Recently a putative pVHL homologue in *c.elegans* has been proposed on the basis of database analysis and a sequence alignment showing 23% amino acid identity. The analysis of HIF regulation in *c.elegans* performed here shows a conserved role for pVHL.

To determine whether pVHL function in HIF regulation might also be conserved we first tested for interaction. 35-S labelled *ceHIF* and HA tagged pVHL were synthesised by IVTT in rabbit reticulocyte lysate, the *ceHIF* and/or tagged pVHL were then added to EBC buffer with or without worm extract, prior to immunoprecipitation with an anti-HA antibody. Co-immunoprecipitation of *ceHIF* with pVHL was observed, but only when the recombinant *ceHIF* IVTT was preincubated with worm extract. Interestingly, though mammalian HIF- α produced in reticulocyte lysates will interact with pVHL, we have found that this is dependent on a factor in the reticulocyte lysate that can be substituted by other mammalian cell extracts, but not the *c.elegans* extract. Though the human and *c.elegans* systems appear homologous, this suggests the existence of a species specific modifying factor that promotes the HIF/pVHL interaction. Mammalian pVHL recognises HIF- α through a subsequence within a transferrable oxygen dependent degradation domain (ODDD) that shows short regions of conservation with *ceHIF*. To test the functional importance of this we mutated a conserved proline residue that is critical for the

reading frame, and raised antisera to a bacterially expression recombinant protein containing amino acids 360 to 497 of the putative protein. The antisera recognised a single species of the appropriate mobility in Cos7 cells transfected with an expression vector expressing the full length cDNA. Total RNA and protein extracts were
5 prepared from populations of worms exposed to normobaric hypoxia by incubation in bell jars flushed with premixed gases of specified oxygen content balanced with nitrogen. Immunoblotting of worm extracts showed a striking induction of *ceHIF* under hypoxia.

Immunoblots of *ceHIF* levels in extracts of *C. elegans* were carried out to
10 monitor regulation by hypoxia and iron chelation. Firstly, the oxygen dependence of protein induction was analysed. Worms were grown on plates in bell jars flushed with air (N), or with oxygen/balance nitrogen having an oxygen concentration of 5%, 1%, 0.5% or 0.1% for 18 hrs. A graded increase in protein level was seen as the oxygen level was reduced below 5% with the highest level of induction at 0.5 and 0.1%
15 oxygen concentration. The time course of protein induction was then studied. Worms were grown in bell jars flushed with a 0.1% oxygen/balance nitrogen mixture for 0, 4, 8, 16 or 24 hrs before preparation of extracts. The results showed strong induction within 4 hours which was sustained over a 24 hour period. The time course of protein decay on re-oxygenation was then assessed. Worms were grown in bell jars flushed
20 with air (N) or a 0.1% oxygen / balance nitrogen mixture. Extracts were made either immediately, or after 4 and 8 minutes of re-oxygenation. Decay of *ceHIF* protein was very rapid on re-oxygenation. Protein levels were clearly reduced after 4 minutes and undetectable after 8 minutes of re-oxygenation of the culture. A time course RNase protection assay showing *cehif* mRNA levels in worms exposed to 0.1%
25 oxygen/balance nitrogen for 0, 4, 8, 16 and 24 hrs was carried out. No induction of *ceHIF* mRNA by hypoxia was seen. Thus *ceHIF* expression was strongly induced by hypoxia at the protein level, but not at the mRNA level, in a manner very similar to that described for mammalian HIF- α subunits. In mammalian cells HIF- α protein is also strongly induced by iron chelating agents as well as hypoxia, a characteristic that
30 has suggested that an interaction of iron and oxygen is central to the underlying mechanism of oxygen sensing. Induction by iron chelation was also studied in *C.*

37. E. E. Patton, A. R. Willems, M. Tyers, *Trends Genet.* **14**, 236-243 (1998).
38. D. E. Richard, E. Berra, E. Gothie, D. Roux, J. Pouyssegur, *J. Biol. Chem.* **274**, 32631-32637 (1999).
39. P. W. Conrad, T. L. Freeman, D. Beitner-Johnson, D. E. Millhorn, *J. Biol. Chem.* **274**, 33709-33713 (1999).
40. M. Rechsteiner, S. W. Rogers, *Trends Biol. Sci.* **21**, 267-271 (1996).

Example 2. Identification of hypoxia inducible factor and von Hippel-Lindau tumour suppressor homologues in *C. elegans*.

10 In this Example we define a HIF homologue in *c.elegans* and demonstrate that both the transcriptional response to hypoxia, and an important mode of regulation through interaction with the von Hippel-Lindau tumour suppressor are conserved.

2.1 Identification of a homologue.

We sought homologues to HIF- α subunits in the *c.elegans* EST database using
15 an tBLASTn enquiry with the human sequence. Prior to completion of the *c.elegans* sequencing programme an EST was found with significant homology to HIF- α in the basic-helix-loop-helix region, and we assembled a contig of ESTs covering the putative homologue. Complete determination of the *c.elegans* sequence revealed a further six predicted PAS proteins but no closer matches to mammalian HIF- α . The
20 EST contig we had identified corresponds to a predicted open reading frame (ORF) on chromosome V (F38A6.3) that is identical except for a 104 amino acid amino terminal extension in the latter. Extensive searching of the EST database has not revealed any cDNAs that map to this putative 5' extension. No PCR products corresponding to the extension could be identified and RACE-PCR products did
25 contain a putative *trans* spliced leader sequence. These findings argue against the predicted N-terminal extension and support the presence of a 719 amino acid protein encoded by 9 exons. Figure 2 shows an alignment of the human and *c.elegans* sequences.

2.2: Regulation of HIF in *c.elegans*.

30 To characterize the putative *c.elegans* HIF homologue (ceHIF), we constructed a riboprobe encompassing nucleotides 1366 to 1496 of the predicted open

detection mode with a cone voltage of 30 V. The collision induced dissociation for polypeptide sequencing by MS/MS was performed with argon gas at 20 – 40 eV and a 3 Da quadrupole resolution.

25. K. I. Kivirikko, R. Myllylä, in *The Enzymology of Post-translational*
5 *Modification of Proteins* R. B. Freeman, H. C. Hawkins, Eds. (Academic Press, London, 1980) pp. 53-104.
26. For pVHL capture assays using biotinylated polypeptides, the polypeptide was interacted with VHL.HA for 30 minutes at 4°C, and precipitated with streptavidin beads. Pre-incubation with cell extract or buffer under test conditions was for 30
10 minutes at 30°C.
27. K. I. Kivirikko, J. Myllyharju, *Matrix Biol* 16, 357-68. (1998).
28. C. J. Schofield, Z. Zhang, *Curr Opin Struct Biol* 9, 722-31. (1999).
29. C. J. Cunliffe, T. J. Franklin, N. J. Hales, G. B. Hill, *J Med Chem* 35, 2652-8. (1992).
- 15 30. G. A. Jansen, et al., *J Lipid Res* 40, 2244-54. (1999).
31. M. Mukherji, M. D. Lloyd et al. unpublished observations.
32. Prolyl 4-hydroxylase activity was assayed by a method based on the hydroxylation-coupled decarboxylation of 2-oxo[1-¹⁴C]glutarate (Kivirikko, K. I., Myllylä, R.: Posttranslational enzymes in the biosynthesis of collagen: intracellular
20 enzymes. *Methods Enzymol.*, 82, 245-304, 1982) using recombinant human type I and II prolyl 4-hydroxylases expressed in insect cells (26). 0.5 or 1.0 mg of polypeptide was used in each reaction. The assay was performed by Dr. Johanna Myllyharju at the Collagen Research Unit, Department of Medical Biochemistry, University of Oulu, Finland.
- 25 33. W. Ehleben, T. Porwol, J. Fandrey, W. Kummer, H. Acker, *Kidney Int.* 51, 483-491 (1997).
34. E. R. Stadtman, *Annu. Rev. Biochem.* 62, 797-821 (1993).
35. Y. Takahashi, S. Takahashi, Y. Shiga, T. Yoshimi, T. Miura, *J Biol Chem* 275, 14139-46. (2000).
- 30 36. D. Skowrya, K. L. Craig, M. Tyers, S. J. Elledge, J. W. Harper, *Cell* 91, 209-219 (1997).

with beads pre-coated with anti-Gal4 antibody RK5C1 (Santa Cruz). After washing with NETN buffer, experimental exposures were to hypotonic extraction buffer (HEB: 20 mM Tris pH7.5, 5mM KCl, 1.5 mM MgCl₂, 1mM DTT) or cell lysate prepared in HEB. Incubations were for 60 minutes at 22°C unless otherwise stated, following which the beads were washed with NETN containing DFO, and incubated for 2 hours at 4°C in NETN+DFO with 5µl rabbit reticulocyte lysate programmed with pcDNA3.VHL.HA.

22. Baculoviral PK.HIF-1α (1-826) or PK.HIF-1α (344-698) were immunoprecipitated with anti-PK antibody (Serotec). Bead bound

10 immunoprecipitates were washed, then incubated with test cell lysates, following which the immunoprecipitates were washed again with NTEN containing DFO, incubated with pVHL, and assayed for interaction.

23. For polypeptide inhibition assays, polypeptides were added to NETN buffer containing a mixture of HIF-1α and pVHL.HA. Final concentration of polypeptide was 1µM unless otherwise stated. Pre-incubation of polypeptide in-cell extract or other conditions was for 60 minutes at 30°C.

24. Samples for mass spectroscopic analyses were either biotinylated synthetic polypeptides 19:WT (residues 556-574), or 34:WT (residues 549-582), or PK-tagged HIF-1α retrieved from insect cell lysates. After modification by mammalian cell lysates the material was purified either by streptavidin/biotin capture (synthetic polypeptides) or anti-PK immunoprecipitation and SDS-PAGE. Proteolytic digestion was performed either on the beads or in-gel with trypsin and V8 protease at pH7.8, or V8 protease at pH4.5. Samples were lyophilised, and dissolved in aqueous 0.1 % TFA. Polypeptides were concentrated, desalted on a 300 µm ID/5 mm length C18 PepMap column (LC Packings, San Francisco, CA, USA) and eluted with 80 % acetonitrile. The HPLC (CapLC, Waters, Milford, MA, USA) was coupled via a Nano-LC inlet to a Q-Tof mass spectrometer (Micromass, Manchester, UK) equipped with a nanoelectrospray Z-spray source. The eluted polypeptide mixture was analysed by tandem mass spectrometric sequencing with an automated MS-to-MS/MS switching protocol. Online determination of precursor-ion masses was performed over the m/z range from 300 to 1200 atomic mass units in the positive charge

11. M. E. Cockman, et al., *J Biol Chem* (2000).
12. M. Ohh, et al., *Nat Cell Biol* 2, 423-7. (2000).
13. T. Kamura, et al., *Proc Natl Acad Sci U S A* 97, 10430-5. (2000).
14. K. Tanimoto, Y. Makino, T. Pereira, L. Poellinger, *Embo J* 19, 4298-309.
5 (2000).
15. M. A. Goldberg, S. P. Dunning, H. F. Bunn, *Science* 242, 1412-1415 (1988).
16. G. L. Wang, G. L. Semenza, *Blood* 82, 3610-3615 (1993).
17. G. L. Semenza, *Cell* 98, 281-284 (1999).
18. Hypoxia (< 0.1% oxygen) was obtained in a workstation with O₂, CO₂ and
10 temperature control (Ruskin Technologies, Leeds, UK). For hypoxic harvest,
buffers were preincubated in the chamber overnight. RCC4-VHL.HA, labelling
conditions and co-immunoprecipitation assays have been described previously (11);
in the current study 12.5µM MG132 was used for proteasomal inhibition. For
standard harvest, the cells were removed from the chamber after hypoxic exposure,
15 prior to cell lysis. Co-immunoprecipitation assays on all lysates were performed at
4°C under ambient oxygen conditions. Parallel experiments established that adding
desferrioxamine (100µM) to the lysis and immunoprecipitation buffers did not alter
the protein species co-precipitated with pVHL.
19. pcDNA3.VHL.HA and pcDNA3.HIF-1α.PK were used to program TNT
20 reticulocyte lysate (Promega). When programming in hypoxia, reaction mix was
preincubated in the workstation for 10 minutes before addition of the DNA template.
An aliquot was removed from the workstation for transcription/translation under
ambient oxygenation. Interaction assays were as described previously (11).
20. Protein expression systems used were wheatgerm lysate (Promega)
25 programmed with pcDNA3 based vectors, insect cell expression using recombinant
baculovirus (pFastBac1, (GibcoBRL) encoding PK.HIF-1α(344-698) and PK.HIF-
1α(1-826)) bacterial expression as glutathione-S-transferase (GST-VBC complex)
and maltose binding protein fusions (pMAL-HIF-1α(344-698)). For insect cell
expression, Sf9 cells (GibcoBRL) were infected 60 hours prior to harvest.
- 30 21. pGal/HIF-1α549-582/VP16 was used to program reticulocyte lysate in the
presence of unlabelled methionine. The fusion protein product was immunopurified

ligases. Furthermore, it is of interest that evolutionarily conserved proline residues are observed at a number of other sites in HIF- α subunits, and that other internal regions of HIF-1 α can convey oxygen-dependent destruction(6). In other studies we have defined a second subdomain within the N-terminal portion of the HIF-1 α

5 oxygen dependent degradation domain that supports pVHL dependent ubiquitylation and contains a functionally critical proline residue. Furthermore, we have established the existence of a functionally conserved pVHL/HIF system in *C.elegans* (see below) and demonstrated the critical importance of a conserved proline residue in the ceVHL/ceHIF interaction (indicated in figure 1A).

10 Overall this suggests that similar marking modifications may occur elsewhere in HIF- α molecules and could contribute to the oxygen sensitive properties of other domains. Whether proline hydroxylation occurs in other molecules on residues that form part of so-called "PEST" domains that are associated with rapid turnover is also clearly now of interest(40). Equally, if the prolyl modification is relatively specific to
15 pVHL-mediated ubiquitylation then the new findings may help define other substrates that are important in pVHL tumor suppressor function.

REFERENCES & NOTES FOR EXAMPLE 1

1. G. L. Semenza, *Genes Dev* 14, 1983-91. (2000).
2. N. V. Iyer, et al., *Genes Dev.* 12, 149-162 (1997).
- 20 3. E. Maltepe, J. V. Schmidt, D. Baunoch, C. A. Bradfield, M. C. Simon, *Nature* 386, 403-407 (1997).
4. G. L. Wang, B.-H. Jiang, E. A. Rue, G. L. Semenza, *Proc. Natl. Acad. Sci. USA* 92, 5510-5514 (1995).
5. S. Salceda, J. Caro, *J. Biol. Chem.* 272, 22642-22647 (1997).
- 25 6. L. E. Huang, J. Gu, M. Schau, H. F. Bunn, *Proc. Natl. Acad. Sci. USA* 95, 7987-7992 (1998).
7. M. S. Wiesener, et al., *Blood* 92, 2260-2268 (1998).
8. P. H. Maxwell, et al., *Nature* 399, 271-275 (1999).
9. K. Iwai, et al., *Proc. Natl. Acad. Sci. USA* 96, 12436-12441 (1999).
- 30 10. J. Lisztwan, G. Imbert, C. Wirbelauer, M. Gstaiger, W. Krek, *Genes Dev* 13, 1822-33 (1999).

5 sulphur clusters) do not exchange in this way it was proposed that such a protein must be turning over rapidly. Second, it has been postulated that cobaltous ions and iron chelators might act by interfering with Fenton chemistry and signalling through reactive oxygen species(17, 33). For instance non-enzymatic 'metal catalysed oxidation' systems that oxidatively modify specific amino acids by local Fenton chemistry can also be inhibited by iron chelators and non-iron transition metal ions (34) providing an alternative hypothesis for effects of these substances on the HIF system. Clearly the labile iron centres associated with prolyl-4-hydroxylases can accommodate the original iron centre substitution hypothesis without the need to propose rapid turnover of the sensor. In contrast we were repeatedly unable to promote specific interactions of HIF- α sequences with pVHL by a variety of non-enzymatic oxidation systems and our evidence clearly indicates an enzymatic mechanism of proline hydroxylation. Our findings do not exclude direct oxidation processes or other oxygen sensing systems impinging on HIF at other sites, on other molecules involved in HIF signal transduction, or indeed on components of the enzymatic prolyl hydroxylation complex. Though our evidence indicates that HIF-PH is distinct from the [α 1 and α 2] prolyl-4 hydroxylases associated with collagen modification, it is interesting that these enzymes employ protein disulfide isomerase as a β subunit, thus providing a potential link to sulfhydryl redox chemistry. Also of interest, P4HA1 has recently been shown to be HIF responsive(35), suggesting that similar hypoxic induction of HIF-PH activity could down-regulate HIF in prolonged hypoxia, contributing to accommodation of the HIF response.

The pVHL multi-protein complex belongs to the SCF class of ubiquitin ligases, with pVHL acting as the F-box like substrate recognition component (36, 37). To date, characterised examples of recognition by F-box proteins have been regulated by phosphorylation of the target sequence. Furthermore, HIF-1 α is a phosphoprotein, and phosphorylation has been implicated in HIF regulation(38, 39). While our findings do not exclude the possibility that HIF- α phosphorylation could influence pVHL recognition, they demonstrate that the key event in recognition of the minimal interaction domain studied here is enzymatic hydroxylation of Pro564. This defines a novel mechanism of regulating substrate recognition for the F-box class of ubiquitin

U2OS cells exposed to dimethyl-oxalylglycine (0, 0.1 or 1.0mM) for 6 hours was carried out. HIF-1 α was seen to be strongly induced under normoxic culture conditions.

Prolyl-4-hydroxylases have been identified in many organisms. In mammalian cells these form $\alpha_2\beta_2$ tetramers in which the β -subunit is identical with the multifunctional protein disulphide isomerase(27). These enzymes function in collagen modification in the endoplasmic reticulum, and are reported to have a strict substrate specificity for prolyl residues in collagen repeat sequences, typically (Pro-Pro-Gly) $_n$ (27). When tested as substrate for recombinant [$\alpha 1$ or $\alpha 2$] human prolyl-4-hydroxylase, the HIF polypeptide showed no activity(32). Taken together these findings lead us to postulate that HIF-PH is a novel prolyl-4-hydroxylase which marks HIF promoting recognition by the pVHL ubiquitination ligase system. Since such enzymes utilise molecular oxygen as a co-substrate this predicts a mechanism for direct sensing of oxygen. To test this we examined the effect of hypoxia in the presence of supplements of other co-factors, on HIF-PH activity as assessed by ability to modify the Gal/HIF-1 α 549-582/VP16 fusion protein so as to promote pVHL capture. HIF substrate was incubated with cell extract (supplemented with 2mM ascorbate and 10 μ M Fe(II)) for 1, 2, 5 or 10 mins at 30°C under ambient conditions or in the hypoxic workstation. The reaction was stopped by washing with DFO, and the HIF substrate assayed for ability to interact with pVHL. A time-dependent increase in capture was seen in normoxia and a marked suppression of activity by hypoxia.

1.6: Summary of Example 1.

Our findings therefore demonstrate a novel method of protein modification that regulates interaction with pVHL ubiquitylation complexes and indicate that enzymatic prolyl hydroxylation may act directly as a sensor of molecular oxygen. The known properties of 2-oxoglutarate dependent oxygenases readily explain the classical features of mimicry of hypoxia by iron chelators or cobaltous ions. Two explanations have been advanced previously for these findings. First, it has been proposed that cobaltous ions might substitute for ferrous ions at an oxygen sensing iron centre (15). Since most iron centres (e.g. haem and the large majority of iron


2 t, $^3J_{\text{HH}}$ 7.3, OCH_2CH_3), 1.95-2.47 (3 H, m, CHCH_2CH_2), 3.88 (3 H, s, OCH_3), 4.10, 4.20 (4 H, 2 quart, $^3J_{\text{HH}}$ 7.3, OCH_2CH_3), 4.60 (1 H, ddd, $^3J_{\text{HH}}$ 8.1, $^3J_{\text{HH}}$ 8.1, $^3J_{\text{HH}}$ 4.8, CH), 7.76 (1 H, d, $^3J_{\text{HH}}$ 8.1, NH); δ_{C} (50 MHz; CDCl_3) 14.1 (OCH_2CH_3), 27.0, 30.1 (CH_2CH_2), 52.1, 53.6 (CH, OCH_3), 60.8, 62.0 (OCH_2CH_3), 156.1, 160.4, 170.5, 172.4 (C=O); $m/z(\text{AP}^+)$ 290 (MH^+ , 68%).

EXAMPLE 9

(S)-2-(Oxalyl-amino)pentanedioic acid (IS13)

10 or: *N*-Oxalyl-(*L*)-glutamate (IS13)

15



3 mmol (0.87 g) of
 20 IS12 was heated with 5.0 ml of 2 N aqueous sodium hydroxide solution ensuring 1.1 equivalents of sodium hydroxide for the sum of the ester functions to be cleaved in the compound. The reaction was percolated through a column of 'Amberlite IR 120 H' ion exchange resin (previously washed with water to about pH 4) and eluted with water until pH raised to 4 again. The water evaporated in vacuo and the residue dried
 25 in vacuum. This yielded 0.65 g (2.9 mmol, 97%) of IS13 as a yellowish hygroscopic


solid, mp ca. 60°C; $[\alpha]_D^{25}$ -2.2 (c 1 in methanol); ν_{\max} (NaCl, MeOH)/cm⁻¹ 1697 (C=O); δ_H (200 MHz; D₂O) 1.77-2.35 (3 H, m, CHCH₂CH₂), 4.30 (1 H, dd, $^3J_{HH}$ 9.1, $^3J_{IH}$ 5.0, CH); δ_C (50 MHz; D₂O) 25.8, 30.3 (CH₂CH₂), 52.6, (CH), 161.4, 162.9, 174.5, 177.3 (C=O); m/z (AP-) 218 (M - H⁺, 5%), 168 (M - H⁺ - oxalyl, 85%).

5

EXAMPLE 10

(S)-2-(Methoxyoxalyl-amino)-propionic acid (IS68)


or: *Methyloxalyl-L-alanine (IS68)*



- 10 This compound was prepared as in Example 8 using 10 mmol (0.89 g) of (L)-alanine and 10 mmol (1.23 g, 0.93 ml) of methyl oxalyl chloride yielding 1.96 g crude yellow oil. The crude product was chromatographed over silica gel (ethyl acetate eluent) resulting in 1.42 g of IS68 as a yellowish oil, which still contained traces of impurities. A pure sample was obtained from recrystallization from a mixture of ethyl
- 15 acetate and diethyl ether (0.39 g, 2.2 mmol, 22%), mp 129-130 °C; $[\alpha]_D^{25}$ 1.1° (c 1 in MeOH); ν_{\max} (NaCl, MeOH)/cm⁻¹ 1744, 1693 (C=O); δ_H (200 MHz; DMSO-*d*₆) 1.35 (3 H, d, $^3J_{HH}$ 7.3, CHCH₃), 3.43 (1 H, br, COOH), 3.81 (3 H, s, OCH₃), 4.28 (1 H, pseudo-quint, $^3J_{HH}$ 7.4, CH), 9.16 (1 H, d, $^3J_{HH}$ 7.5, NH); δ_C (50 MHz; DMSO-*d*₆) 17.3 (CHCH₃), 48.8 (CH), 53.7 (OCH₃), 157.6, 161.8, 173.8 (C=O).

EXAMPLE 11


(R)-2-(Methoxyoxalyl-amino)-propionic acid (IS69)
or: Methyloxalyl-D-alanine (IS69)



- 5 This compound was prepared as for IS68 but with (L)-alanine substituted by (D)-alanine yielding 0.36 g (2.1 mmol, 21%) of IS69 as a colourless solid, mp 131-132 °C; $[\alpha]_D^{25} +1.9^\circ$ (c 1 in MeOH). Analytical data except optical rotation corresponded to those of IS68.

10 **EXAMPLE 12**

(S)-2-(3-Mercapto-propionylamino)-propionic acid (IS37)
or: N-(3-Mercaptopropanoyl)-(L)-alanine (IS37)



- 15 Prepare according to literature procedure: M. A. Ondetti, D. W. Cushman, USP 4053651, 1977, E. R. Squibb & Sons (*Chem. Abstr.*, Volume 88, 136977). No analytical details but melting point were given in the literature work. A solution of 4


mmol (1.13 g) of **IS20** in 2 ml of water was treated with 1.6 ml of conc. aqueous ammonia solution for one hour at room temperature, while a colourless precipitate formed. The mixture was diluted with water and the solids filtered off. The filtrate was washed with ethyl acetate, the aqueous phase was acidified with conc.

- 5 hydrochloric acid and extracted with ethyl acetate. The combined organics were washed with water, dried over magnesium sulfate and evaporated in vacuo resulting in 0.49 g of crude **IS37**. Recrystallization from a mixture of ethyl acetate and *n*-hexane yielded 0.32 g (1.8 mmol, 45%) of **IS37** as a colourless solid, mp 78-79 °C; $[\alpha]_D^{25}$ -39.4 (*c* 1 in methanol); ν_{\max} (NaCl, MeOH)/cm⁻¹ 1728, 1638 (C=O); δ_H (200
- 10 MHz; DMSO-*d*₆) 1.28 (3 H, d, $^3J_{HH}$ 7.3, CH₃), 2.31 (1 H, t, $^3J_{HH}$ 7.9, SH), 2.40-2.48, 2.60-2.73 (4 H, 2 m, CH₂CH₂), 4.22 (1 H, quint, $^3J_{HH}$ 7.3, CH), 8.26 (1 H, d, $^3J_{HH}$ 7.3, NH), 12.56 (1 H, br s, COOH); δ_C (50 MHz; DMSO-*d*₆) 18.0 (CH₃), 20.8 (CH₂CH₂, second signal covered by DMSO, recording in CDCl₃ revealed it at 40.0), 48.3, (CH), 171.0, 175.1 (C=O); *m/z*(AP-) 176 (M - H⁺, 100%).

15

EXAMPLE 13

(R)-2-(3-Mercapto-propionylamino)-propionic acid (**IS38**)
or: *N*-(3-Mercaptopropanoyl)-(*D*)-alanine (**IS38**)



20


The title compound was prepared as for **IS37** but with **IS20** substituted by **IS21** yielding 0.18 g (1.0 mmol, 25%) of **IS38** as a colourless solid, mp 64 °C; $[\alpha]_D^{25}$ +39.5 (*c* 1 in methanol). Analytical data except optical rotation correspond to those of **IS37**.

25

EXAMPLE 14

(S)-2-(3-Benzoylsulfanyl-propionylamino)-propionic acid (IS20)
or: *N*-(3-Benzoylthiopropionyl)-(*L*)-alanine (IS20)

5




Prepared according to literature procedure: M. A. Ondetti, D. W. Cushman, USP 4053651, 1977, E. R. Squibb & Sons (*Chem. Abstr.*, Volume 88, 136977). No analytical details but melting point were given in the literature work.

- 10 In 16.7 ml of 1 N aqueous sodium hydroxide solution, 16.7 mmol (1.48 g) of (*L*)-alanine were dissolved. After adding another 9 ml of 2 N sodium hydroxide solution at ice temperature, 16.7 mmol (2.85 g) of 3-bromopropionic acid were added and the reaction was stirred for 3.5 h at room temperature. A mixture of 18.1 mmol (2.50 g) of thiobenzoic acid and 11.6 mmol (1.6 g) of potassium carbonate in 16.7 ml of water and
- 15 5 ml of THF was then added to the reaction, which was then stirred overnight. The resultant mixture was acidified with conc. hydrochloric acid, stirred for 30 min. and extracted with ethyl acetate. The combined organics were dried and the solvents evaporated in vacuo. The remaining thick (5.15 g) yellow oil was crystallized from ether yielding 1.83 g (6.5 mmol, 39%) of IS20 as a colourless powder, mp 98-99 °C; $[\alpha]_D^{25}$
- 20 -19.1 (*c* 1 in methanol); ν_{\max} (NaCl, MeOH)/cm⁻¹ 1730, 1660 (C=O); δ_H (200 MHz; CDCl₃) 1.44 (3 H, d, $^3J_{HH}$ 7.1, CH₃), 2.66, 3.32 (4 H, 2 d, $^3J_{HH}$ 7.1, CH₂CH₂), 4.61 (1 H, quint, $^3J_{HH}$ 7.1, CH), 6.77 (1 H, d, $^3J_{HH}$ 7.1, NH), 7.37-7.61, 7.89-7.96 (5 H, 2 m, ar), 10.08 (1 H, br s, COOH); δ_C (50 MHz; CDCl₃) 18.0 (CH₃), 24.6, 36.1 (CH₂CH₂), 48.3, (CH), 127.2, 128.7, 133.6, 136.7 (ar), 171.5, 176.0, 192.4 (C=O); *m/z*(AP-) 280 (M - H⁺,
- 25 10%).

EXAMPLE 15

(R)-2-(3-Benzoylsulfanyl-propionylamino)-propionic acid (IS21)
or: *N*-(3-Benzoylthiopropionyl)-(D)-alanine (IS21)

5



This compound was prepared as for IS20 but with (L)-alanine substituting for (D)-alanine yielding 1.78 g (6.3 mmol, 38%) of IS20 as a colourless powder, mp 98-99 °C;

$[\alpha]_D^{25} +19.1$ (*c* 1 in methanol). Analytical data except optical rotation corresponded to

10 those of IS20.

Example 16 - Peptide blockade of HIF- α degradation modulates cellular metabolism and angiogenesis

16.1 Introduction

Ischaemia is a major cause of morbidity and mortality and effective molecular
15 therapies are being intensively sought ^{1 2}. The transcription factor hypoxia-inducible factor-1 (HIF) is a master regulator of the hypoxic response, controlling genes involved in diverse processes that balance metabolic supply and demand within tissues ^{3 4 5}. Modulation of HIF activity therefore provides an attractive approach for the treatment of ischaemic disease. Furthermore, HIF driven angiogenesis produces more mature and
20 less leaky vessels than those generated by individual growth factors ^{6 7 8}.

Regulation of HIF is mediated at multiple levels via its α chain ^{9 10 11 12 13}. It has been reported that PR39, a macrophage derived peptide, results in HIF accumulation and angiogenesis ¹⁴. Analysis of the HIF α oxygen-dependent degradation domains (ODD) by transient transfection studies ^{11 12 15 16 17 18} suggested a possible specific, alternative
25 approach to HIF stabilisation. We have used peptides containing the sites of regulated

prolyl hydroxylation identified as necessary in the previous Examples for proteasomal destruction in the presence of oxygen mediated by the von Hippel-Lindau E3 (VHL E3) ubiquitin ligase complex^{19 20,21 22}. Despite the multiple steps involved in HIF activation we demonstrate unequivocally that peptides from two regions of the ODD not only
5 stabilise HIF α but produce a transcriptional response that modulates normoxic angiogenesis and metabolism *in vivo*, suggesting that the peptides affect mechanisms that are common to all activation steps. These results indicate that these polypeptides, or molecules based on them, provide a possible therapeutic approach for ischaemic tissues.

10 16.2 Overexpression of CODD and NODD polypeptides can induce HRE-dependent reporter gene expression

Since normoxic HIF α degradation is saturable¹⁷ and depends on sub-regions within the ODD we tested whether peptides encoding the HIF-1 α amino terminal ODD (NODD) and carboxy terminal ODD (CODD)²¹ could affect HIF activity as measured by hypoxia response element (HRE)-dependent reporter gene expression. The proposed
15 model by which the NODD and CODD peptides inhibit HIF activity is shown in Figure 13. Briefly, Degradation is prevented by the NODD and CODD polypeptides competing for prolyl hydroxylation and/or VHL binding, thereby blocking subsequent ubiquitination. We constructed plasmids encoding NODD (HIF-1- α aa 343-417) or CODD (HIF-1- α aa 549-582) linked to nuclear localisation sequences and a c-myc
20 epitope tag. These plasmids were transiently co-transfected into U2OS and Hep3B cells with an HRE-dependent luciferase reporter plasmid. Under normoxic conditions expression of either ODD derived polypeptide increased relative luciferase activity after 24 hours to levels comparable to those induced in the absence of peptide by hypoxia (results shown in Figure 13). Transfection of the NODD and CODD expression plasmids
25 had no effect on luciferase activity from reporter plasmids lacking functional HRE's, demonstrating that the effect was mediated via the HRE (results not shown). To confirm that polypeptide action was being mediated via the endogenous HIF pathway we repeated this experiment in a mutant Chinese hamster ovary cell line lacking HIF α chains (Ka13), and a HIF-1 α complemented transfectant (KH-1)²³. Transfection of the NODD and
30 CODD plasmids led to enhanced luciferase activity in KH-1 but not in Ka13 cells

(results shown in Figure 13).

We tested shorter fragments of NODD and CODD peptides, defining amino acids 390-417 and amino acids 556-74 as minimal domains capable of HRE-dependent luciferase activation. (results shown in Figure 13). A sequence alignment of these
5 minimal domains with the equivalent regions from mouse HIF-1- α is shown in Figure 13.

HIF α chain degradation depends on recognition by the VHL E3 ubiquitin ligase following oxygen-dependent enzymatic hydroxylation of prolyl residues at positions 402 and 564^{19 20 21}. Using mutated expression plasmids (P402A or P564G) we demonstrated
10 complete ablation of HRE-dependent induction of luciferase activity (results shown in Figure 13), showing the critical role of these residues for polypeptide function. This suggested that expression of the NODD and CODD fusion proteins interfered with degradation of endogenous HIF α chains, most likely by interfering with VHL recognition or prolyl hydroxylation in normoxic cells, and hence provides a potential
15 route to the therapeutic manipulation of the HIF system.

16.3 Stable NODD and CODD polypeptide expression results in endogenous HIF-1 α accumulation

To explore further the therapeutic potential of NODD and CODD polypeptides to activate endogenous HIF we stably transfected U2OS cells with doxycycline-inducible
20 NODD (doxNODD) and CODD constructs (doxCODD) encoding identical sequences to those used for transient transfections.

Cells stably transfected with NODD or CODD controlled by the tetracycline-inducible system were exposed to doxycycline. Cell extracts were prepared 0, 16, 24 and 48 hours after exposure to doxycycline. The extracts of doxNODD (HIF-1 α aa343-417),
25 doxCODD (HIF-1 α aa549-82) and control cells (empty vector) were then immunoblotted for HIF-1 α protein. Increased HIF-1 α signals were detected from 16-48 hours following doxycycline administration in doxNODD and doxCODD cells but not in empty vector cells. Levels of HIF-1 α induced by hypoxia were also measured for comparison. The endogenous HIF-1 α induction was doxycycline dose (0.2-3.2 μ g/ml) and time

dependent, peaking after 48 hours. Maximal levels were about 20% and about 70% of HIF-1 α levels seen following hypoxic or DFO treatment of the doxNODD and doxCODD cells respectively. Doxycycline did not induce HIF-1 α protein in cells transfected with empty vector despite its weak ability to chelate iron.

5 Immunofluorescence microscopy allowed visualisation of both the c-myc tag of the expressed fusion proteins and endogenous HIF-1 α . In doxycycline activated doxCODD cells both were located in nuclei. HIF-1 α expression varied considerably from cell to cell. In cells which had not been exposed to doxycycline strong staining was only seen from the endogenous HIF-1- α .

10 Combined treatment of doxCODD cells with doxycycline and optimal DFO (75 μ m) or hypoxic stimuli (1% O₂) did not lead to further increases in HIF-1 α signals on immunoblots confirming that the peptides had no additional action when endogenous HIF α chains were fully induced by physiological stimuli.

To test directly whether the polypeptides prevented cellular HIF-1 α targeting by
15 the VHL-ubiquitin-proteasome system we showed that ubiquitination of exogenous ³⁵S-methionine labelled HIF-1 α was markedly reduced in the presence of doxCODD extracts compared with control cell extracts lacking the peptide transfected with empty vector.

HIF and HIF-dependent target gene expression has been suggested to be subject to a number of negative feedback controls. To investigate the consequences of
20 continuous activation of the system we exposed doxCODD cells to doxycycline for two, four, six or eight days. HIF-1 α protein levels were significantly elevated on days 2 and 4 but decreased thereafter. Switching off the system by removing doxycycline from the medium for 48 hours prior to re-exposure resulted in re-induction of elevated HIF-1 α protein levels, indicating that the suppressive effects were reversible. This phenomenon
25 will need to be considered in using the HIF system to modulate complex physiological downstream effects for example through the administration of modulators of the invention at spaced intervals or alternatively by inducing the constructs of the invention at spaced intervals.

16.4 NODD and CODD fusion proteins induce target gene mRNA and protein levels

Results thus far presented indicate that under normoxic conditions NODD and CODD polypeptide expression results in stabilisation of endogenous HIF-1 α chains and consequent activation of transiently transfected artificial HRE-dependent promoters. Expression of natural HIF target genes in chromosomal DNA may be constrained by other factors. We therefore investigated peptide modulation of endogenous genes known to be HIF targets.

Carbonic anhydrase IX (CAIX) is transcriptionally up-regulated under hypoxic conditions²⁴. We used a ribonuclease protection assay to measure CAIX mRNA at intervals following doxycycline treatment in doxCODD and empty vector transfected cells. Levels of mRNA were measured at 0, 24 and 48 hours following treatment. SnRNA (small nuclear RNA) was also probed to ensure equivalent loading. Doxycycline markedly induced mRNA levels in doxCODD cells after 24 and 48 hours to levels similar to those obtained by hypoxic incubation. Cells transfected with empty vector showed no induction of CAIX. Immunoblots demonstrated an associated increase of CAIX protein, paralleling detection of the CODD peptide, visualised by immunoblotting using the c-myc tag. To test the generality of this effect we performed comparable experiments on glucose transporter-1 (Glut-1) mRNA expression, obtaining similar results. When doxycycline was repeatedly added to cell culture medium Glut-1 mRNA, detected by ribonuclease protection, measured after 0, 2, 4, 6 and 8 days Glut-1 mRNA levels increased for the first 4 days and then declined in parallel with the HIF-1 α protein levels as observed above. Maximal Glut-1 mRNA levels were comparable to those induced following exposure to 75 μ M desferrioxamine.

To test for the physiological relevance of increased Glut-1 expression we conducted glucose uptake experiments. ³H-glucose uptake was measured. An enhanced uptake of ³H-glucose was measured in doxCODD cells compared with empty vector transfected cells after 24 hours induction with doxycycline. In contrast, basal levels in cells untreated with doxycycline and hypoxically induced levels (hypoxia) of ³H-glucose uptake were comparable between cell lines. (*: $P < 0.01$; Error bars represent the SEM

of 3 replicates.)

Thus, in contrast to control cells, expression of the CODD polypeptides mimicked the effect of hypoxia by inducing glucose uptake in stably transfected cells.

16.5 *tat*-NODD and *tat*-CODD fusion proteins enter cultured cells and induce HIF-1 α under normoxic conditions

Experiments presented above show that oxygen-dependent gene expression can be modulated in normoxia by plasmid based expression of NODD and CODD polypeptides. To extend this approach we chose to study the effects of transducing comparable peptides into cells. The transduction domain of HIV *tat*-protein delivers fused proteins across cell membranes in a transporter independent mechanism^{25 26}. We fused the NODD and CODD peptides to the *tat*-sequence in combination with HA and HIS tags to facilitate detection and nickel affinity purification. We did not include exogenous nuclear localisation sequences because the *tat* sequence itself is sufficient for nuclear entry²⁷.

We performed VHL E3 interaction assays²¹ with *tat*-NODD and *tat*-CODD, demonstrating their ability to undergo the necessary modifications for interaction with VHL. ³⁵S-Methionine labelled IVTT products of *tat*-ODD expression vectors were tested for their ability to bind to VHL E3 ligase. Concordant with the ubiquitination assays discussed above NODD (HIF-1 α 343-417) and CODD (HIF-1 α 549-582) polypeptides, but not their corresponding proline mutants (HIF-1 α 343-417/P402A and HIF-1 α 549-582/P564G), bound to VHL E3 ubiquitin ligase after modification by cell extracts. The ³⁵S-methionine labelled recombinant polypeptides therefore interacted with VHL, supporting the results of the ubiquitination experiments discussed above. The interaction was enhanced by the presence of cell extracts which promote hydroxylation of the prolines at positions 402 and 564¹⁹. In contrast, no binding occurred using peptides in which prolines were mutated.

We next tested, by immunoblotting, if these *tat*-fusion proteins could traverse cell membranes and induce HIF-1 α . Two hours following addition of *tat*-NODD or *tat*-CODD fusion proteins to cell cultures intact peptide was detectable in whole cell protein

extracts by immunoblotting for the HA tag. The HA tag of *tat*-NODD (*tat*-343-417) and *tat*-CODD (*tat*-549-582) polypeptides were detected in cell extracts, following repetitive polypeptide administration indicating their uptake by the cells. HIF-1 α protein was induced by the *tat*-NODD and the *tat*-CODD polypeptides (0.5 μ M), but not by the corresponding proline mutants. Maximal levels were comparable to those induced following exposure to 75 μ M desferrioxamine (DFO). In experiments with repetitive polypeptide administration, endogenous HIF-1 α was detectable in normoxia 20 hours after initial exposure of the cells to fusion proteins. In controls, using the corresponding mutant peptides lacking the prolines, we detected no HIF-1 α signals. It has been reported that denaturation enhances uptake of *tat*-fused proteins²⁵. Denatured *tat*-NODD and *tat*-CODD peptides were still able to enter cells, but were inactive in mediating HIF-1 α upregulation, perhaps because they were no longer capable of being hydroxylated.

16.6 Endothelial activation and *in vivo* angiogenesis assays

Artificial activation of the HIF signalling pathway using the methods of the invention should induce angiogenesis and will therefore be of potential therapeutic use in ischaemic disease. We tested the effect of polypeptide induced HIF stabilisation in an *in vitro* angiogenesis assay, co-culturing human microvascular endothelial cells (HMEC-1) with empty vector transfected or doxCODD cells. In view of the possibility of sustained activation inducing a negative feedback loop we opted to test the effects of intermittent induction. In doxCODD, but not in empty vector transfected cells, intermittent exposure to doxycycline over a period of 5 days led to assembly of co-cultured endothelial cells into complex tubular structures visualised by immunostaining for von Willebrand factor but not in control cells. As a positive control epidermal Growth Factor (EGF; 5 ng/ml), which is known to induce growth of HMEC-1, was used. To extend these observations into an *in vivo* model we assayed the effects of injecting *tat*-fusion proteins into polyurethane sponges implanted subcutaneously in mice. Intermittent injections on days 1, 2, 4 and 5 led to a markedly accelerated angiogenic response assayed on day 7 when compared with sponges injected with proline mutant fusion proteins, excluding a contribution from the *tat* component²⁸. Immunohistochemistry for von Willebrand factor revealed increased vessel density in sponges explanted after 7 days

following treatment with *tat*-CDD, but not with mutant peptide (*tat*-CDD/P564G). Staining for VEGF and Glut-1 was enhanced in *tat*-NODD or *tat*-CDD treated animals compared to controls. Cells surrounding the sponge showed particularly intense staining. The vessel endothelium within sponges was surrounded by cells expressing smooth muscle actin.

16.7 Summary

Hypoxia-inducible factor-1 (HIF) is a transcription factor known to regulate pro-angiogenic genes and modulate metabolism in response to hypoxic stress. Modulation of HIF activity therefore provides an attractive theoretical route to ameliorating ischaemic disease. Under normoxic conditions HIF α chains are ubiquitinated and destroyed by the proteasome following enzymatic hydroxylation of critical prolyl residues. Here we demonstrate use of polypeptides bearing these prolyl residues to stabilise endogenous HIF, thereby up-regulating HIF target genes. Peptide expression in cell cultures affects physiologically important functions such as glucose transport and leads to tubule formation by co-cultured endothelial cells. Subcutaneous injection of polypeptides results in a markedly accelerated local angiogenic response and induction of glucose transporter-1 gene expression. These results demonstrate the feasibility of utilising these polypeptides to enhance normoxic HIF activity, opening new therapeutic avenues for ischaemic diseases.

In this Example we have described the use of polypeptides which stabilise the hypoxia-regulated transcription factor HIF-1 α . We provide evidence that complex physiological systems like glucose uptake and angiogenesis can be induced strongly, even under normoxic conditions.

Related molecular approaches to treating ischaemic disease include use of single growth factors ² or gene therapy with HIF based sequences lacking the degradation domains ^{6 29}. The approach used here has advantages over the former in that it co-opts the entire physiological response resulting in metabolic adaptation as well as angiogenesis and provides an alternative to gene therapy that should be easier to apply.

Influences of HIF on cancer growth and apoptosis ^{30 31} lead to concerns that long-term HIF activation might have deleterious effects, including pro-neoplastic actions.

However, these processes probably require additional events beyond HIF activation and are likely to have a much longer time course than that required for therapeutic angiogenesis. Furthermore, the peptides used here are inherently unstable and act locally, allowing circumscribed dosing schedules that avoid continued and general exposure.

5 The NODD and CODD polypeptides were effective alone and in combination. Mechanisms of polypeptide action within cells include competition for HIF prolyl hydroxylase activity or VHL binding capacity. Three lines of evidence suggest the latter is more probable. Firstly, we have demonstrated that these NODD and CODD fusion proteins bind to VHL, presumably following their own hydroxylation. Secondly, the
10 action of either peptide is sufficient to stabilise HIF α even though it contains both prolyl residues, which can be targetted by different hydroxylase isoforms ²². Thirdly, concentrations of synthetic peptide necessary to quench HIF prolyl hydroxylase activity in cell extracts are unlikely to be produced in cells.

Comparison of the NODD and CODD sequences coupled with structural studies
15 clarifying the nature of their interactions with VHL and different prolyl hydroxylase isoforms will allow further refinements to these agents. Use of other protein transduction domains and/or tissue specific targetting sequences, including tripeptides such as GFE for lung or RDV for retina, will lead to new formulations with lower risks of side effects
32 33

20 The polypeptides reported here are exciting reagents, allowing controlled activation of the HIF pathway in normoxia. Animal models of ischaemia may be used to demonstrate the net therapeutic benefits of the peptides followed by clinical trials.

16.8 Methods

Plasmids, transient and stable transfections

25 *Plasmid constructs:* For reporter gene assays DNA fragments encoding HIF-1 α amino acids 343-417, 380-417, 390-417, 390-410, 343-400, 530-95, 530-82, 549-82, 556-74 and 530-62 were generated by PCR using oligonucleotides containing 5' SacII or 3' AscI sites and inserted into a pCMV/myc/nuc (Invitrogen) derivative bearing these sites in frame with the NLS and epitope tag. Site directed mutagenesis (QuikChange,
30 Stratagene) was used to mutate the constructs containing HIF-1 α aa343-417 or aa549-82 at aa402 [cca to gca] or aa564 [cca to ggc] converting prolines to alanine or glycine

respectively. To generate tet-operator dependent plasmids the open reading frames from the aa343-417 and aa549-82 constructs were subcloned into pUHD 10³⁴. Fragments coding for HIF-1 α aa343-417 and aa530-82 (with and without P402A and P564G mutants) were subcloned into *ptat*-HA²⁵. All constructs were confirmed by DNA sequencing.

Reporter gene assays: Cells were co-transfected with an HRE containing reporter gene, pCMV/myc/nuc constructs and a constitutively expressed beta-galactosidase gene using Fugene6 (RocheMolecular)³⁵. Transfectants were maintained in normoxia for 24 hours or in hypoxia for the final 16 hours. Luciferase activities in cell extracts were determined using a commercial kit (Promega) and a TD-20e luminometer (Turner Designs). Beta-galactosidase activity was measured spectrophotometrically using *o*-nitrophenyl-beta-D-galactopyranoside as substrate.

Stably transfected cell lines were generated by transfecting U2OS cells bearing the reverse tetracycline responsive transactivator³⁴ and the tetKRAB silencer construct³⁶ with pUHD/HIF plasmids. Following selection in G418 (1 mg/ml) individual colonies were picked. *DoxNODD (F21)* and *doxCODD (myc19)* clones expressed pUHD/HIF-1aa343-417/3NLS/c-myc and pUHD/HIF-1aa549-82/3NLS/c-myc respectively.

mRNA and protein detection

RNA analysis: Total RNA extracted using RNeasy B (Biotec Laboratories) was analysed by ribonuclease protection using ³²P-GTP labelled Glut-1, CAIX and snRNA (as internal control) riboprobes using templates previously described^{10 24}.

Immunoblotting: Cell extracts were prepared in buffer (8M urea, 10% glycerol, 1% SDS, 5 mM DTT, 10 mM Tris/pH 6.8), separated by SDS-PAGE and transferred to Immobilon-P membrane (Millipore). Primary antibodies against HIF-1 α , c-myc tag and HA tag were from Transduction Laboratories, Innogenex and Roche Molecular respectively.

Ubiquitination and interaction assays

Empty vector and doxCODD cells grown to confluence were induced with doxycycline (0.8 μ g/ml) for 48 hours and ubiquitination assays performed using cytoplasmic extracts as described previously³⁷.

For VHL E3 interaction assays ³⁵S-methionine labelled HIF-1 α substrates were prepared by transcription/translation using TnT7 rabbit reticulocyte lysate (Promega). 100

μ M DFO was added to the reaction to suppress prolyl hydroxylase activity of the reticulocyte lysate. Substrate modification was achieved by incubation of HIF-1 α translate with RCC4 cell lysate in the presence of ferrous chloride (100 μ M). Interaction with VHL E3 was analysed as described previously ²¹.

5 Glucose uptake

Empty vector and doxCODD cells were grown to confluence, exposed to 1%, 21% oxygen or 0.8 μ g/ml doxycycline for 16h, washed with glucose-free DMEM and incubated for 10 min with 1 μ Ci/ml 2-deoxy-D ³H-glucose (Amersham, UK), before lysis in 0.5% NP-40, 0.25 M NaCl, 10 mM HEPES/pH 7.6. Glucose uptake was determined by liquid scintillation counting ³⁵.

tat-Protein synthesis and purification

HIF-1 α -*tat*-fusion proteins were purified by sonication of transformed BL21pLysS (Novagen) in lysis buffer (0.5% Tween-20, 50 mM NaH₂PO₄, 300 mM NaCl, 5 mM imidazole) after 4 hours induction with 1mM IPTG. Lysates were spun down at 10³g before loading on a Ni-NTA column (Qiagen). Proteins were eluted with 100 mM imidazole and desalted on a PD10 column (Amersham)²⁵ in 10 mM Tris/pH 7.0 or in 10 mM Tris/pH 8.0, 30 mM KCl. Aliquots were snap frozen in liquid nitrogen. *Tat*-proteins (0.5 μ M final concentration) were given to cultures in DMEM/1%FCS at the beginning and 8 hours after starting the experiments before harvesting the cells after 24 hours.

20 Angiogenesis Assays

Tubule Formation Assay: doxCODD or empty vector cells were co-cultivated with human microvascular endothelial cells (HMEC-1) in a ratio of 2:1 in DMEM 10% doxycycline free FCS (Clontech), 2 mM Glutamine and 100 U/ml Penicillin/100 μ g/ml Streptomycin. Stimulation with doxycycline (0.8 μ g/ml) or epidermal growth factor (5 ng/ml) (Sigma) or control medium was renewed every second day. On day 5 cells were fixed with 70% ethanol, pre-blocked with 1% BSA/PBS. Endothelial cells were detected using antibodies to von Willebrand factor (vWF) (Dako).

Murine sponge model: Sterile polyurethane sponges (8 mm diameter) were inserted subcutaneously under the dorsal skin of anaesthetised black C57 female mice on day 0. On the 1st, 2nd, 4th and 5th days 100 μ l of *tat*-fusion proteins (1 μ M) in Tris buffer (10 mM, pH 7.0) were injected into the sponges. On day 7 mice were sacrificed and

sponges were excised with surrounding tissue and fixed in 3.5% paraformaldehyde.

Immunohistochemistry: Paraffin embedded sponges were cut into 6 µm sections, dewaxed with xylene, rehydrated and stained with vWF (Dako), Glut-1 (Alpha Labs), VEGF (Santa Cruz) and smooth muscle cell actin (DAKO) antibodies. Antigen retrieval,
5 blocking of sections, secondary, HRP labelled antibodies and chromogenic reactions were performed according to the manufacturers' recommendations (DAKO Envision System and Vector Labs ABC Vectastain).

References

1. Marti, H.H. & Risau, W. Angiogenesis in ischemic disease. *Thromb Haemost* 82
10 **Suppl 1**, 44-52. (1999).
2. Ferrara, N. & Alitalo, K. Clinical applications of angiogenic growth factors and their inhibitors. *Nat Med* 5, 1359-64. (1999).
3. Semenza, G.L. Hypoxia-inducible factor 1: master regulator of O₂ homeostasis. *Curr Opin Genet Dev* 8, 588-94. (1998).
- 15 4. Ratcliffe, P.J., O'Rourke, J.F., Maxwell, P.H. & Pugh, C.W. Oxygen sensing, hypoxia-inducible factor-1 and the regulation of mammalian gene expression. *J Exp Biol* 201, 1153-62. (1998).
5. Semenza, G.L. Surviving ischemia: adaptive responses mediated by hypoxia-inducible factor 1. *J Clin Invest* 106, 809-12. (2000).
- 20 6. Elson, D.A. et al. Induction of hypervascularity without leakage or inflammation in transgenic mice overexpressing hypoxia-inducible factor-1alpha. *Genes Dev* 15, 2520-32 (2001).
7. Yancopoulos, G.D. et al. Vascular-specific growth factors and blood vessel formation. *Nature* 407, 242-8. (2000).
- 25 8. Carmeliet, P. & Jain, R.K. Angiogenesis in cancer and other diseases. *Nature* 407, 249-57. (2000).
9. Kallio, P.J. et al. Signal transduction in hypoxic cells: inducible nuclear translocation and recruitment of the CBP/p300 coactivator by the hypoxia-inducible factor-1alpha. *Embo J* 17, 6573-86. (1998).
- 30 10. Maxwell, P.H. et al. The tumour suppressor protein VHL targets hypoxia-inducible factors for oxygen-dependent proteolysis. *Nature* 399, 271-5. (1999).

11. Salceda, S. & Caro, J. Hypoxia-inducible factor 1alpha (HIF-1alpha) protein is rapidly degraded by the ubiquitin-proteasome system under normoxic conditions. Its stabilization by hypoxia depends on redox-induced changes. *J Biol Chem* **272**, 22642-7. (1997).
- 5 12. Huang, L.E., Gu, J., Schau, M. & Bunn, H.F. Regulation of hypoxia-inducible factor 1alpha is mediated by an O₂- dependent degradation domain via the ubiquitin-proteasome pathway. *Proc Natl Acad Sci U S A* **95**, 7987-92. (1998).
13. Arany, Z. et al. An essential role for p300/CBP in the cellular response to hypoxia. *PNAS* **93**, 12969-73. (1996).
- 10 14. Li, J. et al. PR39, a peptide regulator of angiogenesis. *Nat Med* **6**, 49-55. (2000).
15. Srinivas, V., Zhang, L.P., Zhu, X.H. & Caro, J. Characterization of an oxygen/redox-dependent degradation domain of hypoxia-inducible factor alpha (HIF-alpha) proteins. *Biochem Biophys Res Commun* **260**, 557-61. (1999).
16. Yu, F., White, S.B., Zhao, Q. & Lee, F.S. Dynamic, site-specific interaction of
15 hypoxia-inducible factor-1alpha with the von Hippel-Lindau tumor suppressor protein. *Cancer Res* **61**, 4136-42 (2001).
17. O'Rourke, J.F., Tian, Y.M., Ratcliffe, P.J. & Pugh, C.W. Oxygen-regulated and transactivating domains in endothelial PAS protein 1: comparison with hypoxia-inducible factor-1alpha. *J Biol Chem* **274**, 2060-71. (1999).
- 20 18. Pugh, C.W., O'Rourke, J.F., Nagao, M., Gleadle, J.M. & Ratcliffe, P.J. Activation of hypoxia-inducible factor-1; definition of regulatory domains within the alpha subunit. *J Biol Chem* **272**, 11205-14. (1997).
19. Jaakkola, P. et al. Targeting of HIF-alpha to the von Hippel-Lindau ubiquitylation complex by O₂-regulated prolyl hydroxylation. *Science* **292**, 468-72. (2001).
- 25 20. Ivan, M. et al. HIFalpha targeted for VHL-mediated destruction by proline hydroxylation: implications for O₂ sensing. *Science* **292**, 464-8. (2001).
21. Masson, N., Willam, C., Maxwell, P.H., Pugh, C.W. & Ratcliffe, P.J. Independent function of two destruction domains in hypoxia-inducible factor-alpha chains activated by prolyl hydroxylation. *Embo J* **20**, 5197-206 (2001).
- 30 22. Epstein, A.C. et al. C. elegans EGL-9 and Mammalian Homologs Define a Family of Dioxygenases that Regulate HIF by Prolyl Hydroxylation. *Cell* **107**, 43-54 (2001).

23. Wood, S.M. et al. Selection and analysis of a mutant cell line defective in the hypoxia-inducible factor-1 alpha-subunit (HIF-1 alpha). Characterization of hif-1 alpha-dependent and -independent hypoxia-inducible gene expression. *J Biol Chem* 273, 8360-8. (1998).
- 5 24. Wykoff, C.C. et al. Hypoxia-inducible expression of tumor-associated carbonic anhydrases. *Cancer Res* 60, 7075-83. (2000).
25. Nagahara, H. et al. Transduction of full-length TAT fusion proteins into mammalian cells: TAT-p27Kip1 induces cell migration. *Nat Med* 4, 1449-52. (1998).
26. Schwarze, S.R. & Dowdy, S.F. In vivo protein transduction: intracellular delivery of biologically active proteins, compounds and DNA. *Trends Pharmacol Sci* 21, 45-8. 10 (2000).
27. Truant, R. & Cullen, B.R. The arginine-rich domains present in human immunodeficiency virus type 1 Tat and Rev function as direct importin beta-dependent nuclear localization signals. *Mol Cell Biol* 19, 1210-7 (1999).
- 15 28. Albin, A. et al. The angiogenesis induced by HIV-1 tat protein is mediated by the Flk-1/KDR receptor on vascular endothelial cells. *Nat Med* 2, 1371-5 (1996).
29. Vincent, K.A. et al. Angiogenesis is induced in a rabbit model of hindlimb ischemia by naked DNA encoding an HIF-1alpha/VP16 hybrid transcription factor. *Circulation* 102, 2255-61. (2000).
- 20 30. Maxwell, P.H., Pugh, C.W. & Ratcliffe, P.J. Activation of the HIF pathway in cancer. *Curr Opin Genet Dev* 11, 293-9. (2001).
31. Carmeliet, P. et al. Role of HIF-1alpha in hypoxia-mediated apoptosis, cell proliferation and tumour angiogenesis. *Nature* 394, 485-90. (1998).
32. Ho, A., Schwarze, S.R., Mermelstein, S.J., Waksman, G. & Dowdy, S.F. 25 Synthetic protein transduction domains: enhanced transduction potential in vitro and in vivo. *Cancer Res* 61, 474-7. (2001).
33. Rajotte, D. et al. Molecular heterogeneity of the vascular endothelium revealed by in vivo phage display. *J Clin Invest* 102, 430-7 (1998).
34. Gossen, M. et al. Transcriptional activation by tetracyclines in mammalian cells. 30 *Science* 268, 1766-9. (1995).
35. Vaux, E.C. et al. Selection of Mutant CHO Cells with Constitutive Activation of

the HIF System and Inactivation of the von Hippel-Lindau Tumor Suppressor. *J Biol Chem* 276, 44323-30 (2001).

36. Deuschle, U., Meyer, W.K. & Thiesen, H.J. Tetracycline-reversible silencing of eukaryotic promoters. *Mol Cell Biol* 15, 1907-14. (1995).

- 5 37. Cockman, M.E. et al. Hypoxia inducible factor- α binding and ubiquitylation by the von Hippel-Lindau tumor suppressor protein. *J Biol Chem* 275, 25733-41. (2000).

Example 17:

Effect of iron chelation on N-oxalylglycine inhibitory activity and direct 10 comparison of the inhibitory activity of a pair of enantiomers.

To determine whether N-oxalylglycine inhibits HIF-1 α modification via iron chelation we performed capture assays using a Gal/HIF-1 α /VP16 fusion protein expressing HIF-1 α residues 549-582 in the presence of varying concentrations of inhibitor and iron.

15 The unlabelled HIF-1 α substrate was immunopurified on beads, washed, and aliquots incubated in the presence of RCC4 cell extract with 0, 20, 50, 200, 500 or 2000 μ M N-oxalylglycine and either 5 or 100 μ M FeCl₂. After washing, the beads were assayed for their ability to interact with 35-S labelled pVHL IVTT, which was then visualised by fluorography. The results obtained are shown in Figure 14A. The amount
20 of pVHL captured is expressed as relative counts per band. Inhibition of pVHL capture by N-oxalylglycine increased with the concentration of inhibitor and was the same regardless of iron concentration. As iron supplementation has little or no effect on inhibition this shows that the inhibition of HIF-1 α modification by N-oxalylglycine is not mediated by iron chelation but by an alternative mechanism.

25 The inhibitory effect of the pair of enantiomers, N-oxalyl-2S-alanine and N-oxalyl-2R-alanine, on HIF-1 α modification was also studied. This was done using the same pVHL capture assay described above again using the Gal/549-582/VP16 fusion protein as a substrate. The effect of 0, 20, 50, 200, 500 and 2000 μ M concentrations of each enantiomer on pVHL capture was then assessed. The results obtained are shown in
30 Figure 14B. Again, the amount of labelled pVHL captured is expressed as relative counts per band. The results show that there is approximately one log difference in the ability

of the enantiomers to inhibit modification of the HIF substrate and hence pVHL capture. N-oxalyl-2S-alanine enantiomer had the higher inhibitory activity of the two enantiomers.

Example 18

5

In vitro screening of potential inhibitors of HIF modification was performed using a capture assay. A Gal/HIF-1 α /VP16 fusion protein expressing HIF-1 α residues 549-582 was prepared by IVTT and used as a substrate in the assay. The unlabelled substrate was immunopurified on beads, washed, and aliquots incubated in the presence of RCC4 cell
10 extract, with 100 μ M FeCl₂ and 2mM of the potential inhibitor. The inhibitors were either dissolved in DMSO or Tris as indicated. Controls, where no inhibitor but the equivalent amount of DMSO or Tris was added, were also performed. After washing, the beads were assayed for their ability to interact with 35-S labelled pVHL IVTT. Hydroxylation of the
15 fusion protein by HIF hydroxylase present in the cell extract leads to the ability to capture the labelled pVHL and the amount of labelled protein bound to the fusion protein can then be measured to determine relative HIF hydroxylase activity. Figures 15 - 20 show HIF hydroxylase activity in the presence of a particular inhibitor relative to that seen in the absence of the inhibitor (the DMSO/Tris control).


Name	Species	Nucleotide Accession	Protein Accession(s)
Egl-9	C. elegans	AF178536	GI5923812
		GI5923811	
CG1114	D. melanogaster	AE003603	AAF52050
5 Clorf12	H. sapiens	AF229245	NP071334
PHD2		AJ310543	
		gi14547145	Gi14547146
	M. musculus	AJ310546	
PHD1	H. sapiens	BC01723	NP071334
		AJ310543	
		gi14547147	Gi14547148
10 FALKOR	M. musculus	AF340231	Gi13649965
		gi13649964	
FLJ21620	H. sapiens	AK025273	BAB15101
			Colo7838
PHD3	H. sapiens	AJ310545	Gi14547150
		gi14547149	
	M. musculus	AJ310548	Gi14547243
		gi14547242	

<i>In-vitro</i> inhibitor	Disruption of HIF-VHL interaction	<i>In-vivo</i> esterified equivalent	Induction of HIF in tissue culture
NK80	No		
NK81	Yes	Methylmethoxalyl-D/L-alanine	No
NK82	No	Methylmethoxalyl-L/D-alanine	No
NK87	Yes	Methylmethoxalyl glycine	Yes
2, 4 pyridine dicarboxylic acid	Yes	2, 4 diethylpyridine dicarboxylate	No
2, 5 pyridine dicarboxylic acid	Not tested	2, 5 diethylpyridine dicarboxylate	No
2, 6 pyridine dicarboxylic acid	Not tested	2, 6 diethylpyridine dicarboxylate	No

TABLE 2


177

IS 5




5

IS 6




10

IS 7



15

IS 8




20

TABLE 3


178

IS 9




5

NK5




10

NK36



15


NK45



20


179

NK46




5

NK47



10

15 NK84




20

TABLE 3 CONTINUED


180

EDB




5

Benzohydroxamic acid (BHAA)




10

N-oxaloylamino-L-alanine



15

Dimethyloxaloylglycine (MMOG)




20


TABLE 3 CONTINUED

181

Protocatechuic acid




IS68




IS69

5




IS70


182




IS71



C8




C9




183

C10




C11




5

C14



C15




10


Table 3 continued

184


IS12



IS13




IS20




5

IS21




185

IS37



5

IS 38



10

Table 3 (end)

5

10

Designation	Accession No.	Protected Fragment
EGLN1/PHD2	BC 001723	1136-1481
EGLN2/PHD1	AF 229245	4050-4213
EGLN3/PHD3	AK 025273	817-1046
F22B5.4(C.elegans)		210-359
HIF-1(C.elegans)		1366-1496

15

TABLE 4

Gene	Strain	Allele
<i>daf-18</i>	CB1375	<i>e1375</i>
<i>daf-2</i>	CB1370	<i>e1370</i>
<i>age-1</i>	TJ1052	<i>hx546</i>
<i>mev-1</i>	TK22	<i>kn1</i>
<i>clk-1</i>	CB4876	<i>e2519</i>
<i>gas-1</i>	CW152	<i>fc21</i>
<i>ctl-1</i>	TU2463	<i>u800</i>
<i>mev-2</i>	TK93	<i>kn2</i>
<i>2mev-3</i>	TK66	<i>kn10</i>
<i>dpy-18</i>	CB364	<i>e364</i>
<i>phy-2</i>	JK2757	<i>ok177</i>
<i>egl-9</i>	MT1201	<i>n571</i>
<i>egl-9</i>	MT1216	<i>n586</i>
<i>egl-9</i>	JT307	<i>sa307</i>
<i>egl-9</i>	JT330	<i>sa330</i>
<i>vhl-1</i>	CB5603	<i>ok161</i>

Table 5

CLAIMS

1. An assay method for identifying an agent which modulates the interaction of a hypoxia inducible factor (HIF) hydroxylase with a substrate of the hydroxylase, the method comprising:
 - 5 - contacting a HIF hydroxylase and a test substance in the presence of a substrate of the hydroxylase under conditions in which the hydroxylase interacts with the substrate in the absence of the test substance; and
 - determining the interaction, or lack of interaction of, the hydroxylase and the substrate.
- 10 2. An assay method according to claim 1, wherein the interaction of the hydroxylase with the substrate in the presence of the test substance is determined by measuring the hydroxylase activity of the hydroxylase.
3. An assay method according to claim 2, wherein the hydroxylase and the test substance are contacted in the presence of 2-oxoglutarate and the
- 15 hydroxylase activity is determined by determining the turnover of said 2-oxoglutarate to succinate and CO₂.
4. An assay method according to claim 2 or 3, wherein the hydroxylase activity is determined by determining the hydroxylation of one or more proline residues in the substrate.
- 20 5. An assay method according to any one of the preceding claims, wherein the substrate is a HIF polypeptide.
6. An assay method according to claim 5, wherein the HIF polypeptide comprises HIF-1 α , a fragment thereof comprising a proline equivalent to Pro 564 and/or Pro 402 of HIF-1 α or a peptide analogue of HIF-1 α or fragment thereof
- 25 comprising a proline equivalent to Pro 564 and/or Pro 402 of HIF-1 α , and wherein hydroxylation of Pro 564 and/or Pro 402 of or HIF-1 α or of a said equivalent proline is determined.
7. An assay method according to any one of the preceding claims, wherein the substrate comprises a polypeptide sequence comprising residues 556-574,
- 30 such as 549 to 582 of the human HIF-1 α sequence.

8. An assay method according to any one of the preceding claims, wherein the HIF hydroxylase is a prolyl hydroxylase.

9. An assay method according to any of the preceding claims, wherein the HIF hydroxylase comprises:

- 5 (a) the amino acid sequence of SEQ ID NO: 2, 4, 6 or 8;
(b) a variant thereof having at least 60% identity to the amino acid sequence of SEQ ID NO: 2, 4, 6 or 8 and having HIF hydroxylase activity; or
(c) a fragment of either thereof having HIF hydroxylase activity.

10. An assay method according to any one of the preceding claims, wherein the HIF hydroxylase is selected from SEQ ID NO: 2, 4, 6 or 8, FLJ21620 (BAB15101) or Clorf12(NP071334).

11. An assay method according to any one of claims 5 to 10, comprising providing a cell expressing a HIF polypeptide; expressing the HIF hydroxylase recombinantly in the cell; contacting the cell with a test substance and monitoring for HIF polypeptide abundance or stability or HIF-associated activity in the cell to determine the effect of the test substance on HIF hydroxylase activity.

12. An assay method according to claim 11, wherein the HIF associated activity is selected from interaction with von Hippel-Lindau protein (VHL), HIF-mediated transcription or expression of a reporter gene driven by a HIF regulated promoter.

13. An assay method according to any one of the preceding claims, wherein the HIF hydroxylase is a human HIF hydroxylase.

14. An assay method according to any one of the preceding claims, wherein the HIF polypeptide is human HIF-1 α or HIF-2 α .

15. A method for providing a pharmaceutical composition, which method comprises:

- (a) identifying an agent by an assay method as defined in any one of claims 1 to 14; and
(b) formulating the agent thus identified with a pharmaceutically acceptable excipient.

16. An agent identified by an assay method according to any one of claims 1 to 14.
17. A polypeptide comprising:
- (a) the amino acid sequence of SEQ ID NO: 2, 4, 6 or 8;
 - 5 (b) a variant thereof having at least 60% identity to the amino acid sequence of SEQ ID NO: 2, 4, 6 or 8 and having HIF hydroxylase activity; or
 - (c) a fragment of either thereof having HIF hydroxylase activity.
18. A polynucleotide comprising a sequence which encodes a polypeptide as defined in claim 17.
- 10 19. A polynucleotide according to claim 18 which comprises
- (i) SEQ ID NO: 1, 3, 5 or 7 or a complementary sequence thereto;
 - (ii) a sequence which hybridises under stringent conditions to the sequence defined in (i);
 - (iii) a sequence which is degenerate as a result of the genetic code to a
 - 15 sequence as defined in (i) or (ii);
 - (iv) a sequence having at least 60% identity to a sequence as defined in (i);
- or
- (v) a fragment of any of the sequences (i), (ii), (iii) or (iv), and which
 - 20 encodes a polypeptide having hydroxylase activity or capable of generating antibodies specific for a HIF hydroxylase.
20. A polynucleotide according to claim 18 or 19 which is DNA.
21. An expression vector, comprising a polynucleotide as defined in any one of claims 18 to 20.
22. An antibody or a fragment thereof capable of specifically binding a
- 25 polypeptide as defined in claim 17.
23. An agent according to claim 16, a polypeptide according to claim 17, a polynucleotide according to any one of claims 18 to 20, a vector according to claim 21 or an antibody according to claim 22 for use in a method of treatment of the human or animal body by therapy.
- 30 24. A pharmaceutical composition comprising an agent according to claim 16, a polypeptide according to claim 17, a polynucleotide according to any one of


claims 18 to 20, a vector according to claim 21 or an antibody according to claim 22 and a pharmaceutically acceptable excipient.

25. Use of a modulator of HIF hydroxylase activity, an agent according to claim 16, a polypeptide according to claim 17, a polynucleotide according to any one of claims 18 to 20, a vector according to claim 21 or an antibody according to claim 22 in the manufacture of a medicament for the treatment of a condition associated with increased or decreased HIF levels or activity or the treatment of a condition where it is desirable to modulate HIF levels or activity.

26. Use according to claim 25, wherein said condition is selected from the group consisting of ischaemia, wound healing, auto-, allo-, and xeno- transplantation, systemic high blood pressure, cancer, and inflammatory disorders.

27. Use according to claim 25 or 26, wherein the modulator of HIF hydroxylase activity is a collagen prolyl hydroxylase (CPH) inhibitor which inhibits the biological activity of a HIF hydroxylase and wherein the condition is one associated with reduced HIF levels or activity.

28. Use according to claim 27, wherein said inhibitor has the following formula:




where the group R^1 is capable of forming an electrostatic interaction with the sidechain of the arginine which binds the 5-carboxylate of 2-oxoglutarate during catalysis; $A*B$ is a chain of two atoms which are, independently, carbon, oxygen, nitrogen or sulphur, which chain can be functionalised; A, B, C and D being linked to one another by single and/or double and/or triple bonds, y is 0 or 1; and $C*D$ is a chain of two atoms which are, independently, carbon, oxygen, nitrogen, or sulphur, which chain C-D can be functionalised, such that when y is 0 or 1 at least one of the atoms of said chain C-D is capable of chelating with a metal group and when y is 1 said chain C-D is attached to R^2 which is capable of chelating with a metal group.

29. Use according to claim 28, wherein R^1 is $-CO_2H$, $-SO_3H$, $-B(OH)_2$ or $-PO_3H_2$.

30. Use according to claim 28 or 29, wherein y is 0 and C and/or D is attached to $=S$, $=O$, $-SH$ or $-OH$.

31. Use according to claim 28 or 29, wherein y is 1 and R² is -SH, -OH, -CO₂H, -SO₃H, -B(OH)₂, -PO₃H₂, -NHOH, -CONHR³, -CONHOR³, -CONR³OH or CONR³OR³ where in R³ is a branched or straight chain alkyl group of 1 to 6 carbon atoms which can be functionalised.


5 32. Use according to any one of claims 28 to 31, wherein said inhibitor has the following formula:



where R¹ and R may independently be H, a branched or straight C₁ to C₆ alkyl chain which can be functionalised, any natural amino acid side chain, a 4 to 7 membered
10 heterocyclic ring optionally containing 1 or 2 N, S, O or P atoms or a 5 or 6 membered aromatic ring, optionally containing 1 or 2 N, O or S atoms, which can be fused to another ring; and X is NH, NR" where R" is OH, a branched or straight C₁ to C₆ alkyl chain which can be functionalised or O; and Y is O or S.

33. Use according to claim 32, wherein said inhibitor is
15 dimethyloxalylglycine.


34. Use according to any one of claims 28 to 31, wherein said inhibitor has the following formula:



where R¹ and R may independently be H, a branched or straight C₁ to C₆ alkyl chain,
20 any natural amino acid side chain, a 4 to 7 membered heterocyclic ring optionally containing 1 or 2 N, S, O or P atoms or a 5 or 6 membered aromatic ring, optionally

containing 1 or 2 N, O or S atoms, which can be fused to another ring; and Y is O or S.

35. Use according to any one of claims 28 to 31, wherein said inhibitor has the following formula:



5

where R^I to R^V may independently be H, a branched or straight chain alkyl of from 1 to 6 C atoms, a halogen group (i.e. fluoro-, chloro-, bromo- or iodo-), a carboxylate group, a 4 to 7 membered heterocyclic ring optionally containing 1 or more N, S, O or P atoms, a 5 or 6 membered aromatic ring, optionally containing 1 or more N, O or S atoms, which can be fused to another ring or a C(=O)XR group in which:


X is O, NH, NR, where R is H, OH, a branched or straight chain alkyl of from 1 to 6 C atoms, which can be functionalised, alkoxy containing a branched or straight chain alkyl of from 1 to 6 C atoms which can be functionalised, a 4 to 7 membered heterocyclic ring containing 1 or 2 N, S, O or P atoms, a 5 or 6 membered aromatic ring, optionally containing 1 or 2 N, O or S atoms, which can be fused to another ring or O, and m is 0 or 1.

10

15

20


36. Use according to any one of claims 28 to 31, wherein said inhibitor has the following formula:



where R^{III} to R^V may independently be H, a branched or straight C₁ to C₆ alkyl chain.


optionally with 1, 2, 3, 4 or 5 halo substitutions, a 4 to 7 membered heterocyclic ring optionally containing 1 or 2 N, S, O or P atoms, or a 5 or 6 membered aromatic ring, optionally containing 1 or 2 N, O or S atoms, which can be fused to another ring.

37. Use according to anyone of claims 28 to 31, wherein said inhibitor has the following formula:




- where R¹, R², R³ and R⁴ may independently be H, a branched or straight C₁ to C₆ alkyl chain, OH, O-alkyl having a branched or straight C₁ to C₆ alkyl chain optionally containing 1 or 2 N, O or S atoms, COOH, a branched or straight C₁ to C₆ alkyl ester, a 4 to 7 membered heterocyclic ring optionally containing 1 or 2 N, S, O or P atoms or a 5 or 6 membered aromatic ring, optionally containing 1 or more N, O or S atoms, which can be fused to another ring.

38. Use according to any one of claims 28 to 31, wherein said inhibitor has the following formula;



where R^I to R^V may independently be H, a branched or straight C₁ to C₆ alkyl chain, optionally with 1, 2, 3, 4 or 5 halo substitutions, a 4 to 7 membered heterocyclic ring optionally containing 1 or 2 N, S, O or P atoms, or a 5 or 6 membered aromatic ring, optionally containing 1 or 2 N, O or S atoms, which can be fused to another ring.

39. Use according to any one of claims 28 to 31, wherein said inhibitor has the formula:



where A*B*C*D are as defined in claim 28 and R²¹ is hydrogen or acyl.

40. Use according to claim 25, wherein the modulator of HIF hydroxylase is a selective inhibitor of the HIF hydroxylase polypeptide and the condition is one associated with reduced HIF levels or activity.

41. A method of treating a condition associated with increased or reduced HIF activity or a condition in which is desired to modulate HIF activity comprising administering a therapeutically effective amount of an agent according to claim 16, a polypeptide according to claim 17, a modulator of the activity of a polypeptide according to claim 17, a polynucleotide according to any one of claims 18 to 20, a vector according to claim 21 or an antibody according to claim 22 to an individual in need thereof.

42. A method of modulating the amount of HIF polypeptide in a cell comprising contacting the cell with a substance which inhibits the prolyl hydroxylase activity of a HIF hydroxylase.

43. A method according to claim 41 or 42, wherein the modulator is an agent according to claim 16.

44. A method according to any one of claims 41 to 43, wherein the modulator inhibits the biological activity of a HIF hydroxylase but does not inhibit biological activity of a collagen prolyl hydroxylase.

45. A method of making a pharmaceutical composition comprising admixing an agent according to claim 16, a polypeptide according to claim 17, a modulator of the activity of a polypeptide according to claim 17, a polynucleotide according to any one of claims 18 to 20, a vector according to claim 21 or an antibody

according to claim 22 with a pharmaceutically acceptable excipient.

46. A method of producing a HIF hydroxylase comprising:

- (a) causing expression from nucleic acid which encodes a HIF hydroxylase in a suitable expression system to produce the polypeptide recombinantly; and
- (b) determining the hydroxylation of a HIF polypeptide by said recombinantly produced polypeptide.

47. An assay method for identifying a HIF hydroxylase comprising,

- (a) providing a test polypeptide,
- (b) bringing into contact an HIF polypeptide and the test polypeptide under conditions in which the HIF polypeptide is hydroxylated by a HIF hydroxylase; and

(c) determining hydroxylation of the HIF polypeptide.

48. Use of an polypeptide according to claim 17 for the prolyl hydroxylation of a HIF polypeptide.

49. An isolated polypeptide of SEQ ID NO: 9.

DLDLEMLAP*YIPMDDDFQL (SEQ ID NO: 9.)

wherein P* is 4-hydroxy proline.

50. An isolated polypeptide which is a variant of SEQ ID NO: 9, wherein such variants comprises from 1 to 4 amino acid substitutions of any amino acid apart from P*, said variant retaining the ability to antagonise the interaction of a HIF- α subunit with VHL.

51. An assay for a modulator of the interaction between VHL and HIF- α comprising contacting a polypeptide according to claim 49 or claim 50 with a HIF-1 α binding domain of VHL in the presence of a test substance and determining the interaction of the polypeptides.

52. An assay for an inhibitor of VHL-mediated HIF- α destruction, which comprises providing a HIF- α , or fragment thereof which includes a VHL-binding portion, together with its cognate prolyl-hydroxylase under conditions suitable for the hydroxylation of a proline residue in the HIF- α VHL-binding domain;

providing a putative modulator of hydroxylation; and

determining whether the amount of hydroxylation of said proline residue has been modulated by said putative modulator.

53. An assay for a promoter of hydroxylation of a proline residue in HIF- α , which comprises:

5 providing HIF- α , or fragment thereof which includes a VHL-binding portion, under hypoxic conditions, said HIF- α or fragment thereof containing a proline residue in the VHL-binding domain;

providing a putative hydroxylation promoting agent; and

determining whether said agent provides for hydroxylation of said proline.

10 54. A method of identifying a substrate of a polypeptide as defined in claim 17, comprising contacting a test polypeptide substrate with a HIF hydroxylase according to claim 17, and monitoring for hydroxylation of said test polypeptide substrate.

55. A polynucleotide comprising:

- 15 (a) an antisense polynucleotide to the polynucleotide of claim 19 or
(b) an RNAi to the polynucleotide of claim 19.

1/23

HIF-1 α (556-574)	DLDLEMLAPYIPMD-DDFQL
HIF-2 α	ELDLETLAPYIPMDGEDFQL
X.laevis	DLDLEMLAPYIPMD-DDFQL
D.melanogaster	FEAFAMRAPYIPID-DDNPL
C.elegans	EPDLSCLAPFVDY-DMMQM

Figure 1A

POLYPEPTIDE	INHIBITION	INHIBITION WITH EXTRACT
PFSTQDLDLDLEMLAPYIPMDDDFQLRSFDQLSP	-	++
DLDLEMLAPYIPMDDDFQL	-	++
LEMLAPYIPMDDDFQL	-	+
LAPYIPMDDDFQL	-	-
DLDLEMLAPYIPMD	-	-
DLDLEMLAGYIPMDDDFQL	-	-
DLDLEMLAPY*IPMDDDFQL	-	++

Figure 1B

2/23

MEGAGGAN DKKNISSERRKEKSDAARSRRSSSEVEFEYELAHQLF hHIF1Alp
 NE-----DNRNRNMENHRETESHREPDERSES DIEDD LKMCV hif-1
 LPHNVS-SHLDKASVMRLTISYLVVRLLDAG-DLDIEDDMKAQM hHIF1Alp
 IVEEGTVTHLDRIALLHVAATICRLRG--TSENVEENNLENEITN hif-1
 NCEYL---KA LDGEFVVLTDGDMIZISDNV NK MCLTQFEETG hHIF1Alp
 EVWTEDTIAEC LDGPFVIVDSESSILVVTESWAMT LCLTQTDLTG hif-1
 HSVFDETHFC LHEEMREMLTHRNLV-KKGKEQNTQ-----RSFFL hHIF1Alp
 RALRDFLHPSLYD DFD-----KQSKMLHWRPRGEDTDTTGI-NMVL hif-1
 RMECTLTSRCRTMNIKSA TWKVLHCTGHIHVYDT-NSNQPCCCYK hHIF1Alp
 RMR TVISPRGRCLNLKSA LYKSVSEFLVHISKYSTGCHVSFMC GITI hif-1
 -KPPHTCLVLICEPIPHPSNIFILDSKTF LSKHSLUHKFSYCD hHIF1Alp
 PAGQGT-----TNANASAMTKYTESEMC--AGTTRHTCDBRITFVSD hif-1
 RITELMGYEPPEELLGRSIEEYHALLSDHLTKTHHDMETKGQVTT hHIF1Alp
 KFNYILKSELKTLMGTSFYE LVH PADMMIVSK SMKELFAEESHIRT hif-1
 GQYRMLAKRGGYVWVE TQATVLYNT-KNSQPCIVCUNYV VSGII hHIF1Alp
 RYVRLIANDOTLAEIQ T EAT THTGTNGQKG YVICHVYVLCIQG hif-1
 QHDLIFSLQQTECVLKPEVE-SSDMKMTQLFTKVESDOTSSL-EDK hHIF1Alp
 AEESLVVCTDSMPAGHQVDIKKEVODTRDYIGRQPEIVECVDSTP hif-1
 LKKEPDLAL-TLLASAG--DTIISLDFGSNDTETED-----QC hHIF1Alp
 LIEPEEPFDTVIEEVVGGZEPVKQALMGARKNSYDDVLQWLFRD hif-1
 LEEVEL--YNDV-----MLPSPN----- hHIF1Alp
 PSSPBPARYRSADRFRRTTEPSNFGSALASRD FMDSSSRTSRPKTS hif-1
 --EKLQININLAMSPLPTAETPKPLRSAD-PALNQEVA---LKLE hHIF1Alp
 YGRRACSGQSRTTGSSSTSASATLPHSANYSPLEAGISQCGINSP hif-1
 ENPESLELSSTMPQIQDQTPSSSDGSTRQSSPEGNPSEYCFYVD hHIF1Alp
 FSIKSGQVVYGDARSMGRSCDPSDSRRPSALSFSDTLNVSSSTRG hif-1
 SDMVNEFKLELVEKLFAEECTEAKNPFSTQDT----- hHIF1Alp
 INPWIGSNDVFSTMPFA-DSIAIAERIDSSPTLTSGEPILCDDLO hif-1
 --DLDEMLAPYIPMODDFQLRSFDQLSELESSESSSPESASPOST hHIF1Alp
 WEEPDELSCLAPFVDTYEMMO---MDECLDPE-LQALYDLPDFTPA hif-1
 VTVFQQTQIQEPTANATTTTATTDELKTVTKDRMEDIKILIASPS hHIF1Alp
 VPQA-----PAAR hif-1
 FTHLHKETTSATSSPYRDTQSRTASPNRACKGVIEOTEKSHPRS hHIF1Alp
 EVHID-----ESPAPKPM-----GQSG hif-1
 NVLSVALSURTTPPEELNPKILALQNAQRKRKMEHDCSLFCAVG hHIF1Alp
 SLDLFMYTAH-----YOP-- hif-1
 IGTLLQCPDDHAATTSLSKRVKGCKXSSEQNGMEQKTIILIFSDL hHIF1Alp
 ---FOCDETY-----EQ-----GQQQQNQCQPSYSF--- hif-1
 ACRLLGQSMDESGLFQETS YDCEVNAPIQGSRNLLQGEELLRALD hHIF1Alp
 -----FPMLS. hif-1

Figure 2

3/23

HIF-1 α	380	SEDTSFLFDKLLKKEPDALTLLAPAAAGDTIISLDFGSND	417
		— — —	
HIF-2 α	383	SEKSNFLFTKLKEEPEELAQLAPTPGDAIISLDFGNQN	420
		— — —	
HIF-1 α	556	DLGLEMLAPYIPMDDDFQL	574
		— — —	

Figure 3A

4/23




Figure 3B

5/23




Figure 4

6/23




Figure 5

7/23



Hep3B**U2OS**

Figure 6

8/23






Figure 7

9/23



CAS.Fe(II).2-OG



DAOCS.Fe(II).2-OG

Figure 8

10/23

EGL9	IDHDIGGRSRAMLAIYPGNGTRYVKHVDNPFVK
EGLN1	GSYKINGRTRKAMVACYPGNGTGIVVRHVDNPNP
EGLN2	GSYVINGRTRKAMVACYPGNGLGYVRHVDNPNP
EGLN3	KYYVKERSKAMVACYPGNOTGYVRHVDNPNP
Rat SM20	GKYYVKERSKAMVACYPGNGTGIVVRHVDNPNP
P30H	FDGTHLQMARSRNLKNAIVIPHRDFVEL

1

2



EGL9	DGRCITTIYYCNENWDMA
EGLN1	DGRCVTCIYYLNKNWDMA
EGLN2	DGRCITCIYYLNQNWDMA
EGLN3	DGRCITCIYYLNKNWDMA
Rat SM20	DGRCITCIYYLNKNWDMA
P30H	DREVDR YFRTF MV

3

EGL9	TDGGTLRLYPETSMTFMDIDPRADRLVFFWSDRRNPHE
EGLN1	VSQILIRIFPEGKAQFADIEPKFDRLLEFFWSDRRNPHE
EGLN2	VHGGLLQIFPEGRPUVANIEPLEFDRLLEFFWSDRRNPHE
EGLN3	LHGILIRIFPEGKSFADVEPIFDRLLEFFWSDRRNPHE
Rat SM20	LHGGVLRIFPEGKSFVADVEPIFDRLLEFFWSDRRNPHE
P30H	LEDSPA FHSNEDTVIHMRP GEIWFL DAATVHS

4

5

6

7



EGL9	VMPVFRHRFAITIWMDKSERDKALAKGKES
EGLN1	VQPAYATRYAITVWYFDADER ARAKVKY
EGLN2	VKPAYATRYAITVWYFDAKER AAKDKY
EGLN3	VQPSYATRYAMTVWYFDAEER AEAKKRF
Rat SM20	VQPSYATRYAMTVWYFDAEER AEAKKRF
P30H	AVNFSEISRQSLCVDFAF

7

8



Figure 9

11/23





Figure 10

12/23

Fig.11.



SUBSTITUTE SHEET (RULE 26)

13/23




Figure 12

14/23




Figure 13A




Figure 13B

15/23




Figure 13C




Figure 13D

16/23




Figure 13E

N-ODD:

hHIF1 α 390 LKKEPDALTLLAPAAGDTISLDFGSND 417

mHIF1 α 378 LKKEPDALTLLAPAAGDTISLDFGSDD 395

C-ODD:

hHIF1 α 556 _____ DLDLEMLAPYIPMDDDFQL 574

mHIF1 α 543 _____ DLDLEMLAPYIPMDDDFQL 561

motif _____ LxxLAP

Figure 13F

17/23




Figure 14 A




Figure 14 B




Figure 15

19/23




Figure 16

20/23




Figure 17

21/23




Figure 18

22/23




Figure 19




Figure 20

1/35

SEQUENCE LISTING

<110> ISIS INNOVATION LIMITED
 <120> ASSAYS, METHODS AND MEANS
 5 <130> N.84515A PEJ
 <150> GB 0107123.8
 <151> 2001-03-21
 <150> GB 0118952.1
 <151> 2001-08-02
 10 <160> 48
 <170> PatentIn version 3.1
 <210> 1
 15 <211> 2110
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> CDS
 20 <222> (297)..(1517)
 <223>
 <400> 1
 gctttccct gcctgcctgt ctctagtttc tctcacatcc ctttttttt tcctttctct 60
 25 agccaccctg aagggtccct tccaagccc ttagggaccg cagaggactt ggggaccagc 120
 aagcaacccc cagggcacga gaagagctct tgctgtctgc cctgcctcac cctgccccac 180
 accaggcccg gtggcccca gctgcatcaa gtggaggcgg aggaggaggc ggaggagggt 240
 ggcaccatgg gcccgggcgg tgccctccat gcccggggga tgaagacact gctgcc atg 299
 Met
 30 1
 gac agc ccg tgc cag ccg cag ccc cta agt cag gct ctc cct cag tta 347
 Asp Ser Pro Cys Gln Pro Gln Pro Leu Ser Gln Ala Leu Pro Gln Leu
 5 10 15
 cca ggg tct tgc tca gag ccc ttg gag cct gag cct ggc cgg gcc agg 395
 35 Pro Gly Ser Ser Ser Glu Pro Leu Glu Pro Glu Pro Gly Arg Ala Arg
 20 25 30
 atg gga gtg gag agt tac ctg ccc tgt ccc ctg ctc ccc tcc tac cac 443
 Met Gly Val Glu Ser Tyr Leu Pro Cys Pro Leu Leu Pro Ser Tyr His
 40 35 40 45
 tgt cca gga gtg cct agt gag gcc tgc gca ggg agt ggg acc ccc aga 491
 Cys Pro Gly Val Pro Ser Glu Ala Ser Ala Gly Ser Gly Thr Pro Arg
 50 55 60 65
 gcc aca gcc acc tct acc act gcc agc cct ctt cgg gac ggt ttt ggc 539
 45 Ala Thr Ala Thr Ser Thr Thr Ala Ser Pro Leu Arg Asp Gly Phe Gly
 70 75 80
 ggg cag gat ggt ggt gag ctg cgg ccg ctg cag agt gaa ggc gct gca 587
 Gly Gln Asp Gly Gly Glu Leu Arg Pro Leu Gln Ser Glu Gly Ala Ala
 85 90 95
 50 gcg ctg gtc acc aag ggg tgc cag cga ttg gca gcc cag ggc gca cgg 635
 Ala Leu Val Thr Lys Gly Cys Gln Arg Leu Ala Ala Gln Gly Ala Arg

2/35

	100	105	110	
	cct gag gcc ccc aaa cgg aaa tgg gcc gag gat ggt ggg gat gcc cct			683
	Pro Glu Ala Pro Lys Arg Lys Trp Ala Glu Asp Gly Gly Asp Ala Pro			
5	115	120	125	731
	tca ccc agc aaa cgg ccc tgg gcc agg caa gag aac cag gag gca gag			
	Ser Pro Ser Lys Arg Pro Trp Ala Arg Gln Glu Asn Gln Glu Ala Glu			
	130	135	140	145
	cgg gag ggt ggc atg agc tgc agc tgc agc agt ggc agt ggt gag gcc			779
	Arg Glu Gly Gly Met Ser Cys Ser Cys Ser Ser Gly Ser Gly Glu Ala			
10	150	155	160	
	agt gct ggg ctg atg gag gag gcg ctg ccc tct gcg ccc gag cgc ctg			827
	Ser Ala Gly Leu Met Glu Glu Ala Leu Pro Ser Ala Pro Glu Arg Leu			
	165	170	175	
15	gcc ctg gac tat atc gtg ccc tgc atg cgg tac tac ggc atc tgc gtc			875
	Ala Leu Asp Tyr Ile Val Pro Cys Met Arg Tyr Tyr Gly Ile Cys Val			
	180	185	190	
	aag gac agc ttc ctg ggg gca gca ctg ggc ggt cgc gtg ctg gcc gag			923
	Lys Asp Ser Phe Leu Gly Ala Ala Leu Gly Gly Arg Val Leu Ala Glu			
	195	200	205	
20	gtg gag gcc ctc aaa cgg ggt ggg cgc ctg cga gac ggg cag cta gtg			971
	Val Glu Ala Leu Lys Arg Gly Gly Arg Leu Arg Asp Gly Gln Leu Val			
	210	215	220	225
	agc cag agg gcg atc ccg ccg cgc agc atc cgt ggg gac cag att gcc			1019
	Ser Gln Arg Ala Ile Pro Pro Arg Ser Ile Arg Gly Asp Gln Ile Ala			
25	230	235	240	
	tgg gtg gaa ggc cat gaa cca ggc tgt cga agc att ggt gcc ctc atg			1067
	Trp Val Glu Gly His Glu Pro Gly Cys Arg Ser Ile Gly Ala Leu Met			
	245	250	255	
30	gcc cat gtg gac gcc gtc atc cgc cac tgc gca ggg cgg ctg ggc agc			1115
	Ala His Val Asp Ala Val Ile Arg His Cys Ala Gly Arg Leu Gly Ser			
	260	265	270	
	tat gtc atc aac ggg cgc acc aag gcc atg gtg gcg tgt tac cca ggc			1163
	Tyr Val Ile Asn Gly Arg Thr Lys Ala Met Val Ala Cys Tyr Pro Gly			
	275	280	285	
35	aac ggg ctc ggg tac gta agg cac gtt gac aat ccc cac ggc gat ggg			1211
	Asn Gly Leu Gly Tyr Val Arg His Val Asp Asn Pro His Gly Asp Gly			
	290	295	300	305
	cgc tgc atc acc tgt atc tat tac ctg aat cag aac tgg gac gtt aag			1259
	Arg Cys Ile Thr Cys Ile Tyr Tyr Leu Asn Gln Asn Trp Asp Val Lys			
40	310	315	320	
	gtg cat ggc ggc ctg ctg cag atc ttc cct gag ggc cgg ccc gtg gta			1307
	Val His Gly Gly Leu Leu Gln Ile Phe Pro Glu Gly Arg Pro Val Val			
	325	330	335	
45	gcc aac atc gag cca ctc ttt gac cgg ttg ctc att ttc tgg tct gac			1355
	Ala Asn Ile Glu Pro Leu Phe Asp Arg Leu Leu Ile Phe Trp Ser Asp			
	340	345	350	
	cgg cgg aac ccc cac gag gtg aag cca gcc tat gcc acc agg tac gcc			1403
	Arg Arg Asn Pro His Glu Val Lys Pro Ala Tyr Ala Thr Arg Tyr Ala			
	355	360	365	
50	atc act gtc tgg tat ttt gat gcc aag gag cgg gca gcc aaa gac			1451
	Ile Thr Val Trp Tyr Phe Asp Ala Lys Glu Arg Ala Ala Ala Lys Asp			

3/35

	370	375	380	385	
	aag tat cag cta gca tca gga cag aaa ggt gtc caa gta cct gta tca				1499
	Lys Tyr Gln Leu Ala Ser Gly Gln Lys Gly Val Gln Val Pro Val Ser				
		390	395	400	
5	cag ccg cct acg ccc acc tagtgccag tccagagcc gcatggcaga				1547
	Gln Pro Pro Thr Pro Thr				
		405			
	cagcttaaat gacttcagga gagccctggg cctgtgctgg ctgctccttc cctgccaccg				1607
	ctgctgcttc tgactttgcc tctgtcctgc ctgggtgtgga gggctctgtc tgttgcctgag				1667
10	gaccaaggag gagaagagac ctttgcctgcc ccatcatggg ggctgggggtt gtcacctgga				1727
	cagggggcag ccgtggaggc caccgttacc aactgaagct gggggcctgg gtcctaccct				1787
	gtctggtcat gacccatta ggtatggaga gctgggagga ggcattgtca cttcccacca				1847
	ggatgcagga cttgggggtg aggtgagtca tggcctcttg ctggcaatgg ggtgggagga				1907
	gtaccccaaa gtctctcac tcctccagcc tggaaatgtga agtgactccc caacccttt				1967
15	ggccatggca ggcacctttt ggactgggct gccactgctt gggcagagta aaaggtgccca				2027
	ggaggagcat ggggtgtggaa gtcctgtcag ccaagaaata aaagtttacc tcagagctgc				2087
	aaaaaaaaa aaaaaaaaaa aaa				2110
	<210> 2				
20	<211> 407				
	<212> PRT				
	<213> Homo sapiens				
	<400> 2				
25	Met Asp Ser Pro Cys Gln Pro Gln Pro Leu Ser Gln Ala Leu Pro Gln				
	1 5 10 15				
	Leu Pro Gly Ser Ser Ser Glu Pro Leu Glu Pro Glu Pro Gly Arg Ala				
	20 25 30				
30	Arg Met Gly Val Glu Ser Tyr Leu Pro Cys Pro Leu Leu Pro Ser Tyr				
	35 40 45				
	His Cys Pro Gly Val Pro Ser Glu Ala Ser Ala Gly Ser Gly Thr Pro				
	50 55 60				
	Arg Ala Thr Ala Thr Ser Thr Thr Ala Ser Pro Leu Arg Asp Gly Phe				
	65 70 75 80				
35	Gly Gly Gln Asp Gly Gly Glu Leu Arg Pro Leu Gln Ser Glu Gly Ala				
	85 90 95				
	Ala Ala Leu Val Thr Lys Gly Cys Gln Arg Leu Ala Ala Gln Gly Ala				
	100 105 110				
	Arg Pro Glu Ala Pro Lys Arg Lys Trp Ala Glu Asp Gly Gly Asp Ala				
40	115 120 125				
	Pro Ser Pro Ser Lys Arg Pro Trp Ala Arg Gln Glu Asn Gln Glu Ala				
	130 135 140				
	Glu Arg Glu Gly Gly Met Ser Cys Ser Cys Ser Ser Gly Ser Gly Glu				
	145 150 155 160				
45	Ala Ser Ala Gly Leu Met Glu Glu Ala Leu Pro Ser Ala Pro Glu Arg				
	165 170 175				
	Leu Ala Leu Asp Tyr Ile Val Pro Cys Met Arg Tyr Tyr Gly Ile Cys				
	180 185 190				
	Val Lys Asp Ser Phe Leu Gly Ala Ala Leu Gly Gly Arg Val Leu Ala				
50	195 200 205				
	Glu Val Glu Ala Leu Lys Arg Gly Gly Arg Leu Arg Asp Gly Gln Leu				

4/35

210 215 220
 Val Ser Gln Arg Ala Ile Pro Pro Arg Ser Ile Arg Gly Asp Gln Ile
 225 230 235 240
 5 Ala Trp Val Glu Gly His Glu Pro Gly Cys Arg Ser Ile Gly Ala Leu
 245 250 255
 Met Ala His Val Asp Ala Val Ile Arg His Cys Ala Gly Arg Leu Gly
 260 265 270
 Ser Tyr Val Ile Asn Gly Arg Thr Lys Ala Met Val Ala Cys Tyr Pro
 275 280 285
 10 Gly Asn Gly Leu Gly Tyr Val Arg His Val Asp Asn Pro His Gly Asp
 290 295 300
 Gly Arg Cys Ile Thr Cys Ile Tyr Tyr Leu Asn Gln Asn Trp Asp Val
 305 310 315 320
 15 Lys Val His Gly Gly Leu Leu Gln Ile Phe Pro Glu Gly Arg Pro Val
 325 330 335
 Val Ala Asn Ile Glu Pro Leu Phe Asp Arg Leu Leu Ile Phe Trp Ser
 340 345 350
 Asp Arg Arg Asn Pro His Glu Val Lys Pro Ala Tyr Ala Thr Arg Tyr
 355 360 365
 20 Ala Ile Thr Val Trp Tyr Phe Asp Ala Lys Glu Arg Ala Ala Ala Lys
 370 375 380
 Asp Lys Tyr Gln Leu Ala Ser Gly Gln Lys Gly Val Gln Val Pro Val
 385 390 395 400
 Ser Gln Pro Pro Thr Pro Thr
 25 405

<210> 3
 <211> 5163
 <212> DNA
 30 <213> Homo sapiens
 <220>
 <220>
 <221> CDS
 35 <222> (3157)..(4434)
 <223>

<400> 3
 40 ttaggggcag aaaaacattt gtaataatta atggctttga gagacacaag gctttggttg 60
 cccagagta ttagttaacc cacctagtgc tcctaatacat acaatattaa ggattgggag 120
 ggacattcat tgcctcactc tctatttggt tcaccttctg taaaattggt agaataatag 180
 tacccacttc atagcattgt atgatgatta aattggttaa tttttttaa atgcttagaa 240
 cacagattgg gcacataaca gcaagcacca catgtgttta taagataaat tcctttgtgt 300
 tgccttcctg taaagtttaa ataagtaaata aaataaataa atacttgcac gacattttga 360
 45 agtctctcta taacatctga gtaagtggcg gctgcgacaa tgctactgga gttccagaat 420
 cgtgttggtg acaagattgt tcaccagcat atggtgtggt gaaaactcac taatttgga 480
 ttagttcaga ttattaagcc tgaataggtg aaatcctga aatcaaggat ctttggaact 540
 atttgaaatc agtattttat attttcctgt tgtattcatt aaagtgttgc aagtgttcta 600
 tttgatggat taagtatatt taggatatac atgttcaatt tgtgattttg tataactaat 660
 50 tggaacaaga aagctaataa aggttttgat atggacatct attcttttaa gtaaacttca 720
 atgaaaatat atgagtagag catatagaga tgtaaataat ttgtggacac accacagact 780

5/35

	gaaatagcaa	atttaaaaga	aattgttggga	agaatcaagt	gtttgtggaa	tgagtcctcc	840
	tagtaaagtt	cctgctcttg	tgaataatta	agcctcatgt	ataattacta	tagcaaaagg	900
	aagcctaaga	agtattagac	tctacttgta	tttaaattac	attttacata	atztatgtgt	960
	atgaaaaatg	ttttaaatgc	ttattttcgt	aagccatgag	atagctcctt	tatattttta	1020
5	gaatttctga	attaatttgc	ttggatttta	ttagtgcata	tgacagagct	agcaattcct	1080
	ttttctgtgt	tcccatcca	tcctattcat	ccctctttta	ggaaactctg	aactctggat	1140
	tgctctgtgt	tacatacctg	cctcctgcat	tggaactatg	gtctctgagt	gtagtatgac	1200
	taattcattt	gtttgtcaag	gactctcaat	gcatttgttg	aacagcctaa	ttagtaatgt	1260
	ctgcaacaat	gacattttac	tgtatttaat	aaagctctgg	gaaagtagga	tacacataag	1320
10	acaggcttag	gtctaaattc	tttacagaaa	cttggatttt	tagttcgggt	tgaaatttga	1380
	agatgtgagt	atatttatct	cagtttccca	aaggacaagc	taattggaat	tatcatcctc	1440
	tttcaactga	ttggatcccc	agaatgccat	ttacgcagtc	agcaggattt	tataacagtt	1500
	ttaaattctg	tatatttgat	gaagagggtt	tatatttttg	gattcaagcc	tctttttaaa	1560
	cttctacaat	atgggtttaca	ataattcctt	atatctgct	tttgaaatac	atattacaac	1620
15	tttttaagtt	tggaaggcta	tatttcaagg	actgaagtta	cagtatactc	aagtataca	1680
	caagcctagc	accccaactt	ccacatagtg	ttcgataaag	attgataaac	tcgaaatcac	1740
	agacctttta	attcttaaga	caaatagcag	cagaaagaaa	catctttggc	ttatttctgg	1800
	taagggtttt	atgctctgta	aaacaaagaa	ttgtattcat	ccgcgcagca	cagattctat	1860
	taaaaataaa	tgtagagagc	gttaatgtag	tactgctcat	ttaccatcaa	aattcacttt	1920
20	tcaggaataa	tcccatcagt	ttaaattgga	tattggaatg	agcattgatt	acatttaact	1980
	tggtagccca	aaatttcttc	atggggtttt	gaactcggcg	ggatttcaaa	ggttttaaaa	2040
	atgagttttt	gatttttttt	aaaaccctca	aatttcatta	cctttaaact	aggtcgaaac	2100
	ggggcgcaag	agattggatt	aacaccatag	taatacttat	ttgttctta	accatttcag	2160
	ggcttcttga	aatagaggct	gtatggtgta	atggaaaaaa	cagccttgga	atctgggagc	2220
25	ctgattcctg	gattcagtc	cagttttgcg	tgacctggg	caagttactt	tacttctctg	2280
	aatttccgtt	tcctcctctg	caaaatgagg	atcgcaatag	ccaccttgca	accttgactg	2340
	gagcgagcct	cgacaccccc	gcgccggcct	ggagggaagag	cagccatgat	tacgccgcct	2400
	tcgctccgct	accgcttg	ggctggcgcc	ctcctccagc	aggtgtaggc	gctgccgcgc	2460
	tgccccacgc	cttccgccc	ctcgcgccc	tgccctcgg	cgtccccgag	gaggccgctg	2520
30	cgggctgagg	tagcgcaccg	gcctctcggc	gtccagtc	ggtcccgggc	ggagggaag	2580
	cgggcgaccc	acctccgagg	cagaagccga	ggcccgccc	cgccgagtc	ggaggagcgc	2640
	aggcagcccc	cgccctcgg	ccctccccc	ggccctccc	gcccctccc	cgccctccc	2700
	gccctcgcgc	gccgcccgc	cggtcgccc	cgggccgctg	gtgtacgtg	agagcgcgca	2760
	gagcgagtgg	cgcccgatg	ccctgcgctc	ctccacagcc	tgggccgggc	cgccggggac	2820
35	gctgaggcgg	cggcggcggc	cgagggggcc	ggtcttgcc	tcccaggcc	cgcgcgctg	2880
	agcccaggtt	gccattcgcc	gcacaggccc	tattctctca	gccctcggcg	gcgatgaggc	2940
	gctgaggcgg	ctgccggcgc	tgcccgagg	cttaggactc	ggaagcgcc	gggcccagg	3000
	cgtgggggtg	cgccctccct	gaggcgagg	tagcgggtg	atggcgagc	aacggccct	3060
	atctctctcc	ccgctcccca	gcctcgggcg	aggccgtccg	gccgtaccc	ctcctgctcg	3120
40	gccgcccag	tcgccgtgc	cgcccgccc	gccgc atg	gcc aat gac agc ggc		3174
				Met Ala Asn Asp Ser Gly			
				1	5		
	ggg ccc ggc	ggg ccg agc	ccg agc gag	cga gac	cgg cag tac	tgc gag	3222
	Gly Pro Gly	Gly Pro Ser	Pro Ser Glu	Arg Asp Arg	Gln Tyr Cys	Glu	
45		10	15	20			
	ctg tgc ggg	aag atg gag	aac ctg ctg	cgc tgc agc	cgc tgc cgc	agc	3270
	Leu Cys Gly	Lys Met Glu	Asn Leu Leu	Arg Cys Ser	Arg Cys Arg	Ser	
		25	30	35			
	tcc ttc tac	tgc tgc aag	gag cac cag	cgt cag gac	tgg aag aag	cac	3318
50	Ser Phe Tyr	Cys Cys Lys	Glu His Gln	Arg Gln Asp	Trp Lys Lys	His	
	40	45	50				

6/35

	aag ctc gtg tgc cag ggc agc gag ggc gcc ctc ggc cac gga gtg ggc	3366
	Lys Leu Val Cys Gln Gly Ser Glu Gly Ala Leu Gly His Gly Val Gly	
	55 60 65 70	
5	cca cac cag cat tcc ggc ccc gcg ccg cgt gca gtg ccg ccg ccc	3414
	Pro His Gln His Ser Gly Pro Ala Pro Ala Ala Val Pro Pro Pro	
	75 80 85	
	agg gcc ggg gcc cgg gag ccc agg aag gca gcg gcg cgc cgg gac aac	3462
	Arg Ala Gly Ala Arg Glu Pro Arg Lys Ala Ala Ala Arg Arg Asp Asn	
	90 95 100	
10	gcc tcc ggg gac gcg gcc aag gga aaa gta aag gcc aag ccc ccg gcc	3510
	Ala Ser Gly Asp Ala Ala Lys Gly Lys Val Lys Ala Lys Pro Pro Ala	
	105 110 115	
	gac cca gcg gcg gcc gcg tcg ccg tgt cgt gcg gcc gcc ggc ggc cag	3558
	Asp Pro Ala Ala Ala Ala Ser Pro Cys Arg Ala Ala Ala Gly Gly Gln	
15	120 125 130	
	ggc tcg gcg gtg gct gcc gaa gcc gag ccc ggc aag gag gag ccg ccg	3606
	Gly Ser Ala Val Ala Ala Glu Ala Glu Pro Gly Lys Glu Glu Pro Pro	
	135 140 145 150	
20	gcc cgc tca tcg ctg ttc cag gag aag gcg aac ctg tac ccc cca agc	3654
	Ala Arg Ser Ser Leu Phe Gln Glu Lys Ala Asn Leu Tyr Pro Pro Ser	
	155 160 165	
	aac acg ccc ggg gat gcg ctg agc ccc ggc ggc ggc ctg cgg ccc aac	3702
	Asn Thr Pro Gly Asp Ala Leu Ser Pro Gly Gly Gly Leu Arg Pro Asn	
	170 175 180	
25	ggg cag acg aag ccc ctg ccg gcg ctg aag ctg gcg ctc gag tac atc	3750
	Gly Gln Thr Lys Pro Leu Pro Ala Leu Lys Leu Ala Leu Glu Tyr Ile	
	185 190 195	
	gtg ccg tgc atg aac aag cac ggc atc tgt gtg gtg gac gac ttc ctc	3798
	Val Pro Cys Met Asn Lys His Gly Ile Cys Val Val Asp Asp Phe Leu	
30	200 205 210	
	ggc aag gag acc gga cag cag atc ggc gac gag gtg cgc gcc ctg cac	3846
	Gly Lys Glu Thr Gly Gln Gln Ile Gly Asp Glu Val Arg Ala Leu His	
	215 220 225 230	
35	gac acc ggg aag ttc acg gac ggg cag ctg gtc agc cag aag agt gac	3894
	Asp Thr Gly Lys Phe Thr Asp Gly Gln Leu Val Ser Gln Lys Ser Asp	
	235 240 245	
	tcg tcc aag gac atc cga ggc gat aag atc acc tgg atc gag ggc aag	3942
	Ser Ser Lys Asp Ile Arg Gly Asp Lys Ile Thr Trp Ile Glu Gly Lys	
	250 255 260	
40	gag ccc ggc tgc gaa acc att ggg ctg ctc atg agc agc atg gac gac	3990
	Glu Pro Gly Cys Glu Thr Ile Gly Leu Leu Met Ser Ser Met Asp Asp	
	265 270 275	
	ctg ata cgc cac tgt aac ggg aag ctg ggc agc tac aaa atc aat ggc	4038
	Leu Ile Arg His Cys Asn Gly Lys Leu Gly Ser Tyr Lys Ile Asn Gly	
45	280 285 290	
	cgg acg aaa gcc atg gtt gct tgt tat ccg ggc aat gga acg ggt tat	4086
	Arg Thr Lys Ala Met Val Ala Cys Tyr Pro Gly Asn Gly Thr Gly Tyr	
	295 300 305 310	
50	gta cgt cat gtt gat aat cca aat gga gat gga aga tgt gtg aca tgt	4134
	Val Arg His Val Asp Asn Pro Asn Gly Asp Gly Arg Cys Val Thr Cys	
	315 320 325	

7/35

	ata tat tat ctt aat aaa gac tgg gat gcc aag gta agt gga ggt ata	4182
	Ile Tyr Tyr Leu Asn Lys Asp Trp Asp Ala Lys Val Ser Gly Gly Ile	
	330 335 340	
5	ctt cga att ttt tca gaa ggc aaa gcc cag ttt gct gac att gaa ccc	4230
	Leu Arg Ile Phe Pro Glu Gly Lys Ala Gln Phe Ala Asp Ile Glu Pro	
	345 350 355	
	aaa ttt gat aga ctg ctg ttt ttc tgg tct gac cgt cgc aac cct cat	4278
	Lys Phe Asp Arg Leu Leu Phe Trp Ser Asp Arg Arg Asn Pro His	
	360 365 370	
10	gaa gta caa cca gca tat gct aca agg tac gca ata act gtt tgg tat	4326
	Glu Val Gln Pro Ala Tyr Ala Thr Arg Tyr Ala Ile Thr Val Trp Tyr	
	375 380 385 390	
	ttt gat gca gat gag aga gca cga gct aaa gta aaa tat cta aca ggt	4374
	Phe Asp Ala Asp Glu Arg Ala Arg Ala Lys Val Lys Tyr Leu Thr Gly	
15	395 400 405	
	gaa aaa ggt gtg agg gtt gaa ctc aat aaa cct tca gat tcg gtc ggt	4422
	Glu Lys Gly Val Arg Val Glu Leu Asn Lys Pro Ser Asp Ser Val Gly	
	410 415 420	
20	aaa gac gtc ttc tagagccttt gatccagcaa taccacactt cacctacaat	4474
	Lys Asp Val Phe	
	425	
	attgttaact attgtttaac ttgtgaatac gaataaatgg gataaagaaa aatagacaac	4534
	cagttcgcat ttaataagg aaacagaaac aactttttgt gttgcatcaa acagaagatt	4594
	ttgactgctg tgactttgta ctgcatgac aacttcaa atgtgattgc ttacaggagg	4654
25	aagataagct actaattgaa aatgggtttt acatctggat atgaaataag tgcctgtgt	4714
	agaatttttt tcattcttat attttgccag atctgttatc tagctgagtt catttcatct	4774
	ctcccttttt tatatcaagt ttgaatttgg gataattttt ctatattagg tacaatttat	4834
	ctaaactgaa ttgagaaaaa attacagtat tattcctcaa aataacatca atctattttt	4894
	gtaaacctgt tcatactatt aaattttgcc ctaaaagacc tcttaataat gattgttgcc	4954
30	agtgactgat gattaatttt attttactta aaataagaaa aggagcactt taattacaac	5014
	tgaaaaatca gattgttttg cagtccttcc ttactactaat ttgaactctt aaagattgct	5074
	gctttttttt tgacattgtc aataacgaaa cctaattgta aaacagtcac catttactac	5134
	caataacttt tagttaatgt tttaacaagg	5163
35	<210> 4	
	<211> 426	
	<212> PRT	
	<213> Homo sapiens	
	<400> 4	
40	Met Ala Asn Asp Ser Gly Gly Pro Gly Gly Pro Ser Pro Ser Glu Arg	
	1 5 10 15	
	Asp Arg Gln Tyr Cys Glu Leu Cys Gly Lys Met Glu Asn Leu Leu Arg	
	20 25 30	
45	Cys Ser Arg Cys Arg Ser Ser Phe Tyr Cys Cys Lys Glu His Gln Arg	
	35 40 45	
	Gln Asp Trp Lys Lys His Lys Leu Val Cys Gln Gly Ser Glu Gly Ala	
	50 55 60	
50	Leu Gly His Gly Val Gly Pro His Gln His Ser Gly Pro Ala Pro Pro	
	65 70 75 80	
	Ala Ala Val Pro Pro Pro Arg Ala Gly Ala Arg Glu Pro Arg Lys Ala	

8/35

			85			90			95							
	Ala	Ala	Arg	Arg	Asp	Asn	Ala	Ser	Gly	Asp	Ala	Ala	Lys	Gly	Lys	Val
			100						105					110		
5	Lys	Ala	Lys	Pro	Pro	Ala	Asp	Pro	Ala	Ala	Ala	Ala	Ser	Pro	Cys	Arg
			115					120						125		
	Ala	Ala	Ala	Gly	Gly	Gln	Gly	Ser	Ala	Val	Ala	Ala	Glu	Ala	Glu	Pro
			130				135					140				
	Gly	Lys	Glu	Glu	Pro	Pro	Ala	Arg	Ser	Ser	Leu	Phe	Gln	Glu	Lys	Ala
			145			150					155				160	
10	Asn	Leu	Tyr	Pro	Pro	Ser	Asn	Thr	Pro	Gly	Asp	Ala	Leu	Ser	Pro	Gly
				165						170					175	
	Gly	Gly	Leu	Arg	Pro	Asn	Gly	Gln	Thr	Lys	Pro	Leu	Pro	Ala	Leu	Lys
			180						185					190		
	Leu	Ala	Leu	Glu	Tyr	Ile	Val	Pro	Cys	Met	Asn	Lys	His	Gly	Ile	Cys
15			195					200					205			
	Val	Val	Asp	Asp	Phe	Leu	Gly	Lys	Glu	Thr	Gly	Gln	Gln	Ile	Gly	Asp
			210				215					220				
	Glu	Val	Arg	Ala	Leu	His	Asp	Thr	Gly	Lys	Phe	Thr	Asp	Gly	Gln	Leu
			225			230					235				240	
20	Val	Ser	Gln	Lys	Ser	Asp	Ser	Ser	Lys	Asp	Ile	Arg	Gly	Asp	Lys	Ile
				245						250					255	
	Thr	Trp	Ile	Glu	Gly	Lys	Glu	Pro	Gly	Cys	Glu	Thr	Ile	Gly	Leu	Leu
			260						265					270		
	Met	Ser	Ser	Met	Asp	Asp	Leu	Ile	Arg	His	Cys	Asn	Gly	Lys	Leu	Gly
25			275					280					285			
	Ser	Tyr	Lys	Ile	Asn	Gly	Arg	Thr	Lys	Ala	Met	Val	Ala	Cys	Tyr	Pro
			290				295					300				
	Gly	Asn	Gly	Thr	Gly	Tyr	Val	Arg	His	Val	Asp	Asn	Pro	Asn	Gly	Asp
			305			310					315				320	
30	Gly	Arg	Cys	Val	Thr	Cys	Ile	Tyr	Tyr	Leu	Asn	Lys	Asp	Trp	Asp	Ala
				325						330					335	
	Lys	Val	Ser	Gly	Gly	Ile	Leu	Arg	Ile	Phe	Pro	Glu	Gly	Lys	Ala	Gln
			340						345					350		
	Phe	Ala	Asp	Ile	Glu	Pro	Lys	Phe	Asp	Arg	Leu	Leu	Phe	Phe	Trp	Ser
35			355					360					365			
	Asp	Arg	Arg	Asn	Pro	His	Glu	Val	Gln	Pro	Ala	Tyr	Ala	Thr	Arg	Tyr
			370				375					380				
	Ala	Ile	Thr	Val	Trp	Tyr	Phe	Asp	Ala	Asp	Glu	Arg	Ala	Arg	Ala	Lys
			385			390					395				400	
40	Val	Lys	Tyr	Leu	Thr	Gly	Glu	Lys	Gly	Val	Arg	Val	Glu	Leu	Asn	Lys
				405						410					415	
	Pro	Ser	Asp	Ser	Val	Gly	Lys	Asp	Val	Phe						
				420					425							
45	<210>	5														
	<211>	2770														
	<212>	DNA														
	<213>	Homo sapiens														
	<220>															
50	<221>	CDS														
	<222>	(327)..(1043)														

9/35

<223>

<400> 5

	gagtcctggcc	gcagtcgcgg	cagtggtggc	ttcccatccc	caaaaggcgc	cctccgactc	60
5	cttgccgcgc	actgctcgcc	gggcccagtc	ggaaacgggt	cgtggagctc	cgcaccactc	120
	ccgctggttc	ccgaaggcag	atcccttctc	ccgagagttg	cgagaaaactt	tcccttgctc	180
	ccgacgctgc	agcggctcgg	gtaccgtggc	agccgcaggt	ttctgaaccc	cgggccacgc	240
	tcccgcgcgc	tcggttcgc	gctcgtgtag	atcgttccct	ctctggttgc	acgctgggga	300
	tcccggacct	cgattctgcg	ggcgag atg	ccc ctg gga	cac atc atg	agg ctg	353
10			Met Pro Leu Gly His Ile Met Arg Leu				
			1		5		
	gac ctg gag	aaa att gcc	ctg gag tac	atc gtg ccc	tgt ctg cac	gag	401
	Asp Leu Glu	Lys Ile Ala	Leu Glu Tyr	Ile Val Pro	Cys Leu His	Glu	
	10	15	20	25			
15	gtg ggc ttc	tgc tac ctg	gac aac ttc	ctg ggc gag	gtg gtg ggc	gac	449
	Val Gly Phe	Cys Tyr Leu	Asp Asn Phe	Leu Gly Glu	Val Val Gly	Asp	
	30	35	40				
	tgc gtc ctg	gag cgc gtc	aag cag ctg	cac tgc acc	ggg gcc ctg	cgg	497
	Cys Val Leu	Glu Arg Val	Lys Gln Leu	His Cys Thr	Gly Ala Leu	Arg	
20		45	50	55			
	gac ggc cag	ctg gcg ggg	ccg cgc gcc	ggc gtc tcc	aag cga cac	ctg	545
	Asp Gly Gln	Leu Ala Gly	Pro Arg Ala	Gly Val Ser	Lys Arg His	Leu	
	60	65	70				
	cgg ggc gac	cag atc acg	tgg atc ggg	ggc aac gag	gag ggc tgc	gag	593
25	Arg Gly Asp	Gln Ile Thr	Trp Ile Gly	Gly Asn Glu	Glu Gly Cys	Glu	
	75	80	85				
	gcc atc agc	ttc ctc ctg	tcc ctc atc	gac agg ctg	gtc ctc tac	tgc	641
	Ala Ile Ser	Phe Leu Leu	Ser Leu Ile	Asp Arg Leu	Val Leu Tyr	Cys	
	90	95	100	105			
30	ggg agc cgg	ctg ggc aaa	tac tac gtc	aag gag agg	tct aag gca	atg	689
	Gly Ser Arg	Leu Lys Tyr	Tyr Val Lys	Glu Arg Ser	Lys Ala Met		
	110	115	120				
	gtg gct tgc	tat ccg gga	aat gga aca	ggt tat gtt	cgc cac gtg	gac	737
	Val Ala Cys	Tyr Pro Gly	Asn Gly Thr	Gly Tyr Val	Arg His Val	Asp	
35		125	130	135			
	aac ccc aac	ggt gat ggt	cgc tgc atc	acc tgc atc	tac tat ctg	aac	785
	Asn Pro Asn	Gly Asp Gly	Arg Cys Ile	Thr Cys Ile	Tyr Tyr Leu	Asn	
	140	145	150				
	aag aat tgg	gat gcc aag	cta cat ggt	ggg atc ctg	cgg ata ttt	cca	833
40	Lys Asn Trp	Asp Ala Lys	Leu His Gly	Gly Ile Leu	Arg Ile Phe	Pro	
	155	160	165				
	gag ggg aaa	tca ttc ata	gca gat gtg	gag ccc att	ttt gac aga	ctc	881
	Glu Gly Lys	Ser Phe Ile	Ala Asp Val	Glu Pro Ile	Phe Asp Arg	Leu	
	170	175	180	185			
45	ctg ttc ttc	tgg tca gat	cgt agg aac	cca cac gaa	gtg cag ccc	tct	929
	Leu Phe Phe	Trp Ser Asp	Arg Arg Asn	Pro His Glu	Val Gln Pro	Ser	
	190	195	200				
	tac gca acc	aga tat gct	atg act gtc	tgg tac ttt	gat gct gaa	gaa	977
	Tyr Ala Thr	Arg Tyr Ala	Met Thr Val	Trp Tyr Phe	Asp Ala Glu	Glu	
50		205	210	215			
	agg gca gaa	gcc aaa aag	aaa ttc agg	aat tta act	agg aaa act	gaa	1025

10/35

	Arg	Ala	Glu	Ala	Lys	Lys	Lys	Phe	Arg	Asn	Leu	Thr	Arg	Lys	Thr	Glu	
	220							225					230				
	tct	gcc	ctc	act	gaa	gac	tgacc	gtgct	ctgaa	atctg	ctggc	cttgt					1073
	Ser	Ala	Leu	Thr	Glu	Asp											
5	235																
	tcatt	tttagt	aac	ggttc	cct	gaatt	ctctt	aaatt	ctttg	agat	ccaaag	atggc	ctctt				1133
	cagt	gacaac	aat	ctcc	ctg	ctact	ttctt	cat	cttcac	atcc	ctgtct	tgt	gtgtggt				1193
	act	tcatgtt	ttct	tgccaa	gact	gtgtt	g	atct	tcagat	act	ctctt	g	ccagatgaag				1253
	ttatt	tgcta	act	ccagaaa	ttc	ctgcaga	cat	cttactc	ggcc	agcgg	ttac	ctgata					1313
10	gatt	cggtaa	tact	atcaag	aga	agagcct	agg	agcacag	cgag	ggaatg	aac	cttactt					1373
	gcact	ttatg	tata	cttcct	gatt	tgaaag	gagg	aggtt	gaaa	agaaaa	aaat	ggagg	g				1433
	ggtag	atgcc	acag	agagg	atc	acggaag	cct	taacagc	agg	aaacaga	gaa	attt	gtg				1493
	tcatt	ctgaac	aatt	tcaga	tg	ttcttaat	ccag	ggctgt	tggg	gttct	ggag	aattat					1553
	caca	acctaa	tgac	attaat	acct	ctagaa	agg	gctgtg	tcata	gtgaa	caatt	tataa					1613
15	gtgt	cccatg	ggg	cagacac	tc	ctttttc	ccag	tctgc	aac	ctggatt	ttct	gcctca					1673
	gct	ccatttt	gct	gaaaata	atg	actttct	gaata	aaagat	ggca	acacaa	tttt	ttctcc					1733
	at	tttcagtt	ctt	acctggg	aac	ctaattc	ccc	agaagct	aaaa	aactag	acatt	agttg					1793
	tttt	ggttgc	ttt	gttgaa	tgga	attta	at	taaatga	aagg	aaaaat	atat	ccctgg					1853
	tag	ttttgtg	tta	accactg	ata	actgtgg	aa	agagctag	gtc	tactgat	ata	caataaa					1913
20	cat	gtgtgca	tct	gaacaa	ttt	gagagg	gagg	tggagt	tgg	aatgtg	ggt	gttcctg					1973
	tttt	tttttt	tttt	tttttagt	ttt	cttttt	aat	gagctca	cc	tttaaca							2033
	caaaaa	aagc	agg	gtgatgt	at	tttaaaaa	agg	aagtgg	ata	aaaaaaa	tct	caaagct					2093
	att	tgattc	tcg	tctgtcc	ctag	cagtct	tt	cttcagct	cact	tggctc	tct	agatcca					2153
	ctgt	ggttgg	cagt	atgacc	aga	atcatgg	aact	tgtctag	aact	gtggaa	gct	tctactc					2213
25	ctg	cagtaag	cac	agatcgc	act	gcctcaa	ta	acttggta	ttg	agcacgt	at	tttgcaaa					2273
	ag	tactttt	cct	agttttc	agt	attactt	tcat	gtttta	aaa	atccctt	ta	tttcttg					2333
	ctt	gaaaatc	ccat	gaacat	taa	agagcca	gaa	atatttt	ct	ttgttat	gtac	ggatat					2393
	at	atatatat	atag	tcttcc	aag	atagaag	ttt	acttttt	cct	cttctgg	ttt	tgaaaa					2453
	ttt	ccagata	agac	atgtca	ccat	taattc	tca	acgactg	ctc	tattttg	ttg	tacggta					2513
30	atag	ttatca	cct	tctaaat	tact	atgtaa	ttt	actcact	tatt	atgttt	att	gtcctgt					2573
	at	ctttctc	tgg	agtgtaa	gcac	aatgaa	gac	aggaatt	ttg	tatattt	tta	accaatg					2633
	caac	atactc	tcag	cacct	aaat	agtgcc	ggg	aacatag	taag	ggctca	gtaa	atactt					2693
	gtt	gaataaa	ctc	agtctcc	tac	attagca	ttc	taaaaaa	aaaa	aaaaaaa	aaaa	aaaaaaa					2753
	aaaa	aaaaaaa	aaaa	aag													2770
35	<210>	6															
	<211>	239															
	<212>	PRT															
	<213>	Homo sapiens															
40	<400>	6															
	Met	Pro	Leu	Gly	His	Ile	Met	Arg	Leu	Asp	Leu	Glu	Lys	Ile	Ala	Leu	
	1				5					10					15		
45	Glu	Tyr	Ile	Val	Pro	Cys	Leu	His	Glu	Val	Gly	Phe	Cys	Tyr	Leu	Asp	
			20						25					30			
	Asn	Phe	Leu	Gly	Glu	Val	Val	Gly	Asp	Cys	Val	Leu	Glu	Arg	Val	Lys	
			35					40					45				
	Gln	Leu	His	Cys	Thr	Gly	Ala	Leu	Arg	Asp	Gly	Gln	Leu	Ala	Gly	Pro	
		50					55					60					
50	Arg	Ala	Gly	Val	Ser	Lys	Arg	His	Leu	Arg	Gly	Asp	Gln	Ile	Thr	Trp	
	65					70				75					80		

11/35

Ile Gly Gly Asn Glu Glu Gly Cys Glu Ala Ile Ser Phe Leu Leu Ser
 85 90 95
 Leu Ile Asp Arg Leu Val Leu Tyr Cys Gly Ser Arg Leu Gly Lys Tyr
 100 105 110
 5 Tyr Val Lys Glu Arg Ser Lys Ala Met Val Ala Cys Tyr Pro Gly Asn
 115 120 125
 Gly Thr Gly Tyr Val Arg His Val Asp Asn Pro Asn Gly Asp Gly Arg
 130 135 140
 10 Cys Ile Thr Cys Ile Tyr Tyr Leu Asn Lys Asn Trp Asp Ala Lys Leu
 145 150 155 160
 His Gly Gly Ile Leu Arg Ile Phe Pro Glu Gly Lys Ser Phe Ile Ala
 165 170 175
 Asp Val Glu Pro Ile Phe Asp Arg Leu Leu Phe Phe Trp Ser Asp Arg
 180 185 190
 15 Arg Asn Pro His Glu Val Gln Pro Ser Tyr Ala Thr Arg Tyr Ala Met
 195 200 205
 Thr Val Trp Tyr Phe Asp Ala Glu Glu Arg Ala Glu Ala Lys Lys Lys
 210 215 220
 20 Phe Arg Asn Leu Thr Arg Lys Thr Glu Ser Ala Leu Thr Glu Asp
 225 230 235

<210> 7

<211> 2369

25 <212> DNA

<213> Caenorhabditis elegans

<220>

<221> CDS

30 <222> (11)..(2182)

<223>

<400> 7

agcacatgac atg agc agt gcc cca aat gat gac tgt gag atc gac aag 49
 35 Met Ser Ser Ala Pro Asn Asp Asp Cys Glu Ile Asp Lys
 1 5 10
 gga aca cct tct acc gct tca ctt ttt aca acg ctg atg ctc agt caa 97
 Gly Thr Pro Ser Thr Ala Ser Leu Phe Thr Thr Leu Met Leu Ser Gln
 15 20 25
 40 cca tct tct tct aca gct gtt tta cag tgt aca tat tgt gga agc tcg 145
 Pro Ser Ser Ser Thr Ala Val Leu Gln Cys Thr Tyr Cys Gly Ser Ser
 30 35 40 45
 tgc aca tct tcc caa ttg caa aca tgt tta ttc tgt gga aca gtg gct 193
 Cys Thr Ser Ser Gln Leu Gln Thr Cys Leu Phe Cys Gly Thr Val Ala
 45 50 55 60
 tat tgt tcc aag gag cac cag caa ctc gat tgg cta aca cat aaa atg 241
 Tyr Cys Ser Lys Glu His Gln Gln Leu Asp Trp Leu Thr His Lys Met
 65 70 75
 50 ata tgc aag tca ctt caa aca agt ggc atg gtg cca agt aat ttg atg 289
 Ile Cys Lys Ser Leu Gln Thr Ser Gly Met Val Pro Ser Asn Leu Met
 80 85 90

12/35

	cct cag gca gca cct gct gtt atg gct cca att cca cct act gtt tcg	337
	Pro Gln Ala Ala Pro Ala Val Met Ala Pro Ile Pro Pro Thr Val Ser	
	95 100 105	
5	ttt gat gat cct gca ctt acc acg tca ctt ctt cta tct ctt caa aat	385
	Phe Asp Asp Pro Ala Leu Thr Thr Ser Leu Leu Leu Ser Leu Gln Asn	
	110 115 120 125	
	aat cca att ctg aat caa act att tca aat ttt ccg cca aca ttt tcg	433
	Asn Pro Ile Leu Asn Gln Thr Ile Ser Asn Phe Pro Pro Thr Phe Ser	
	130 135 140	
10	atc aca tcg aag acc gaa cca gag cca tcg att cca atc caa att cca	481
	Ile Thr Ser Lys Thr Glu Pro Glu Pro Ser Ile Pro Ile Gln Ile Pro	
	145 150 155	
	caa agg ata tca tca aca agt aca gta ccg ttc agt agt gaa gga agt	529
15	Gln Arg Ile Ser Ser Thr Ser Thr Val Pro Phe Ser Ser Glu Gly Ser	
	160 165 170	
	gca ttc aaa cca tac aga aat acg cat gtg ttt aat tca att tct tct	577
	Ala Phe Lys Pro Tyr Arg Asn Thr His Val Phe Asn Ser Ile Ser Ser	
	175 180 185	
20	gaa tca atg tct tcc atg tgc aca tca cat gaa gca tca ctt gaa cac	625
	Glu Ser Met Ser Ser Met Cys Thr Ser His Glu Ala Ser Leu Glu His	
	190 195 200 205	
	atg tca tca gct tcc ctt gca atg ttc cca aca agt agt act gct caa	673
	Met Ser Ser Ala Ser Leu Ala Met Phe Pro Thr Ser Ser Thr Ala Gln	
	210 215 220	
25	agt gat atc agt aga ctc gct caa gtt ttg agt ctt gct gga gat tca	721
	Ser Asp Ile Ser Arg Leu Ala Gln Val Leu Ser Leu Ala Gly Asp Ser	
	225 230 235	
	cca gct tcg ttg gct ctt gtc aca act tcg gta ccg tca act gct tcc	769
30	Pro Ala Ser Leu Ala Leu Val Thr Thr Ser Val Pro Ser Thr Ala Ser	
	240 245 250	
	aca gca act att cca cct cca gcg aca acg aca agt tca gct aca agt	817
	Thr Ala Thr Ile Pro Pro Pro Ala Thr Thr Ser Ser Ala Thr Ser	
	255 260 265	
35	tca ggc aaa agc gag aca ata act gtt gga aaa gaa aag ata att caa	865
	Ser Gly Lys Ser Glu Thr Ile Thr Val Gly Lys Glu Lys Ile Ile Gln	
	270 275 280 285	
	act gat gat ccg gat att cag atc atc gaa aca gaa ggt gga tca aaa	913
	Thr Asp Asp Pro Asp Ile Gln Ile Ile Glu Thr Glu Gly Gly Ser Lys	
	290 295 300	
40	cca acg gta tcc aga aca cgg aaa cga cca act cct tct aac tcc gct	961
	Pro Thr Val Ser Arg Thr Arg Lys Arg Pro Thr Pro Ser Asn Ser Ala	
	305 310 315	
	gac cca aaa att aat tac aag gat cac aat aag aat gtc gtt tac tcg	1009
45	Asp Pro Lys Ile Asn Tyr Lys Asp His Asn Lys Asn Val Val Tyr Ser	
	320 325 330	
	aca acc ctc caa gaa cat cag aag cat ctt cag aat cgt ggt ctc gca	1057
	Thr Thr Leu Gln Glu His Gln Lys His Leu Gln Asn Arg Gly Leu Ala	
	335 340 345	
50	cta agc att cac caa gca atg gtt cta cgg tta aga tac att gcc gag	1105
	Leu Ser Ile His Gln Ala Met Val Leu Arg Leu Arg Tyr Ile Ala Glu	
	350 355 360 365	

13/35

	cat	gtg	atc	aga	agc	ctg	aat	gag	ttt	gga	tgg	gcc	gtt	gtt	gac	aat	1153
	His	Val	Ile	Arg	Ser	Leu	Asn	Glu	Phe	Gly	Trp	Ala	Val	Val	Asp	Asn	
				370						375					380		
5	ttt	ctg	ggc	tcg	gat	cac	tac	aaa	ttt	acc	gcg	aaa	gaa	att	gag	cga	1201
	Phe	Leu	Gly	Ser	Asp	His	Tyr	Lys	Phe	Thr	Ala	Lys	Glu	Ile	Glu	Arg	
				385					390						395		
	ctc	tat	gaa	cgg	gga	ctc	ttc	agc	cct	ggt	cag	ttg	atg	gaa	gca	aaa	1249
	Leu	Tyr	Glu	Arg	Gly	Leu	Phe	Ser	Pro	Gly	Gln	Leu	Met	Glu	Ala	Lys	
			400					405						410			
10	cac	aaa	gac	gaa	ttt	cac	atc	aaa	gat	att	cga	tct	gac	cac	att	tac	1297
	His	Lys	Asp	Glu	Phe	His	Ile	Lys	Asp	Ile	Arg	Ser	Asp	His	Ile	Tyr	
			415					420					425				
	tgg	tat	gat	ggt	tat	gat	gga	cgt	gcc	aag	gat	gct	gca	act	gtt	cgt	1345
	Trp	Tyr	Asp	Gly	Tyr	Asp	Gly	Arg	Ala	Lys	Asp	Ala	Ala	Thr	Val	Arg	
15	430						435				440					445	
	cta	ttg	att	tca	atg	att	gat	tct	gta	att	caa	cat	ttc	aaa	aaa	cga	1393
	Leu	Leu	Ile	Ser	Met	Ile	Asp	Ser	Val	Ile	Gln	His	Phe	Lys	Lys	Arg	
				450					455						460		
	att	gat	cat	gat	att	gga	gga	cgt	tct	cgt	gca	atg	ctt	gcc	atc	tat	1441
20	Ile	Asp	His	Asp	Ile	Gly	Gly	Arg	Ser	Arg	Ala	Met	Leu	Ala	Ile	Tyr	
				465					470						475		
	cct	gga	aat	gga	act	cgt	tat	gtg	aag	cat	gta	gat	aat	ccg	gta	aaa	1489
	Pro	Gly	Asn	Gly	Thr	Arg	Tyr	Val	Lys	His	Val	Asp	Asn	Pro	Val	Lys	
			480					485						490			
25	gat	gga	aga	tgt	ata	acc	act	att	tat	tac	tgt	aat	gaa	aat	tgg	gat	1537
	Asp	Gly	Arg	Cys	Ile	Thr	Thr	Ile	Tyr	Tyr	Cys	Asn	Glu	Asn	Trp	Asp	
			495					500					505				
	atg	gca	act	gat	ggt	ggt	act	ctc	aga	tta	tat	cca	gag	act	tca	atg	1585
	Met	Ala	Thr	Asp	Gly	Gly	Thr	Leu	Arg	Leu	Tyr	Pro	Glu	Thr	Ser	Met	
30	510					515					520					525	
	act	cca	atg	gat	att	gat	cca	agg	gct	gat	cgt	ctg	gta	ttc	ttc	tgg	1633
	Thr	Pro	Met	Asp	Ile	Asp	Pro	Arg	Ala	Asp	Arg	Leu	Val	Phe	Phe	Trp	
				530					535						540		
	tcc	gat	cgt	cgc	aat	cct	cat	gaa	gtc	atg	cca	gtc	ttc	cgt	cat	cgt	1681
35	Ser	Asp	Arg	Arg	Asn	Pro	His	Glu	Val	Met	Pro	Val	Phe	Arg	His	Arg	
				545					550					555			
	ttc	gca	att	act	att	tgg	tat	atg	gat	aaa	tcc	gaa	aga	gat	aag	gct	1729
	Phe	Ala	Ile	Thr	Ile	Trp	Tyr	Met	Asp	Lys	Ser	Glu	Arg	Asp	Lys	Ala	
			560					565					570				
40	ttg	gca	aaa	gga	aaa	gag	tca	gat	gcg	gca	tgt	gct	tca	aag	aaa	gag	1777
	Leu	Ala	Lys	Gly	Lys	Glu	Ser	Asp	Ala	Ala	Cys	Ala	Ser	Lys	Lys	Glu	
			575					580					585				
	aat	gat	cca	aca	agc	tct	tca	cta	aat	tcc	ctt	att	gga	tca	ctt	ttg	1825
	Asn	Asp	Pro	Thr	Ser	Ser	Ser	Leu	Asn	Ser	Leu	Ile	Gly	Ser	Leu	Leu	
45	590					595					600				605		
	aga	cca	cgg	aaa	aat	cca	agt	act	cac	gat	tta	tca	aaa	ctt	gac	ctt	1873
	Arg	Pro	Arg	Lys	Asn	Pro	Ser	Thr	His	Asp	Leu	Ser	Lys	Leu	Asp	Leu	
				610					615					620			
	cga	ctc	ttc	ccg	tcc	aca	tca	tcc	gat	cca	gct	ctg	gta	tct	gca	gca	1921
50	Arg	Leu	Phe	Pro	Ser	Thr	Ser	Ser	Asp	Pro	Ala	Leu	Val	Ser	Ala	Ala	
				625					630					635			

14/35

gat gaa gat aga gtt gac atc tct gcc gac ttt caa tcc act tca agt 1969
 Asp Glu Asp Arg Val Asp Ile Ser Ala Asp Phe Gln Ser Thr Ser Ser
 640 645 650
 5 ctg gct cat ccg gaa tct act gac tcg gga gta tct ctc tcc acc ttc 2017
 Leu Ala His Pro Glu Ser Thr Asp Ser Gly Val Ser Leu Ser Thr Phe
 655 660 665
 aat gtc gct cat aat cac atg gaa cgt act acc agt ctc cag tcg atc 2065
 Asn Val Ala His Asn His Met Glu Arg Thr Thr Ser Leu Gln Ser Ile
 670 675 680 685
 10 tcc gat cat ttc cgt tcc gaa aga tca cac gaa cgt cgc agc tca aca 2113
 Ser Asp His Phe Arg Ser Glu Arg Ser His Glu Arg Arg Ser Ser Thr
 690 695 700
 agc agc gat caa gat cta gac gaa ggg ctc cca cca cct cct tcc aca 2161
 Ser Ser Asp Gln Asp Leu Asp Glu Gly Leu Pro Pro Pro Pro Ser Thr
 705 710 715
 15 aac cca gag tat tac atc tga agtttttctg gtttttgta ctttctatat 2212
 Asn Pro Glu Tyr Tyr Ile
 720
 atatatatgt caccttcatt caatacgggt aaagtcaatc ttgaaattcc gattcccgag 2272
 20 aaaatcattg tcattcgagt ttttttatgt atggactcta caaattatta tctcgacttt 2332
 tccatgtgag atagagtacc aattcaacat ggttttt 2369

<210> 8

25 <211> 723
 <212> PRT
 <213> Caenorhabditis elegans

<400> 8

30 Met Ser Ser Ala Pro Asn Asp Asp Cys Glu Ile Asp Lys Gly Thr Pro
 1 5 10 15
 Ser Thr Ala Ser Leu Phe Thr Thr Leu Met Leu Ser Gln Pro Ser Ser
 20 25 30
 35 Ser Thr Ala Val Leu Gln Cys Thr Tyr Cys Gly Ser Ser Cys Thr Ser
 35 40 45
 Ser Gln Leu Gln Thr Cys Leu Phe Cys Gly Thr Val Ala Tyr Cys Ser
 50 55 60
 Lys Glu His Gln Gln Leu Asp Trp Leu Thr His Lys Met Ile Cys Lys
 65 70 75 80
 40 Ser Leu Gln Thr Ser Gly Met Val Pro Ser Asn Leu Met Pro Gln Ala
 85 90 95
 Ala Pro Ala Val Met Ala Pro Ile Pro Pro Thr Val Ser Phe Asp Asp
 100 105 110
 Pro Ala Leu Thr Thr Ser Leu Leu Leu Ser Leu Gln Asn Asn Pro Ile
 115 120 125
 45 Leu Asn Gln Thr Ile Ser Asn Phe Pro Pro Thr Phe Ser Ile Thr Ser
 130 135 140
 Lys Thr Glu Pro Glu Pro Ser Ile Pro Ile Gln Ile Pro Gln Arg Ile
 145 150 155 160
 50 Ser Ser Thr Ser Thr Val Pro Phe Ser Ser Glu Gly Ser Ala Phe Lys
 165 170 175

15/35

Pro Tyr Arg Asn Thr His Val Phe Asn Ser Ile Ser Ser Glu Ser Met
 180 185 190
 Ser Ser Met Cys Thr Ser His Glu Ala Ser Leu Glu His Met Ser Ser
 195 200 205
 5 Ala Ser Leu Ala Met Phe Pro Thr Ser Ser Thr Ala Gln Ser Asp Ile
 210 215 220
 Ser Arg Leu Ala Gln Val Leu Ser Leu Ala Gly Asp Ser Pro Ala Ser
 225 230 235 240
 10 Leu Ala Leu Val Thr Thr Ser Val Pro Ser Thr Ala Ser Thr Ala Thr
 245 250 255
 Ile Pro Pro Pro Ala Thr Thr Thr Ser Ser Ala Thr Ser Ser Gly Lys
 260 265 270
 Ser Glu Thr Ile Thr Val Gly Lys Glu Lys Ile Ile Gln Thr Asp Asp
 275 280 285
 15 Pro Asp Ile Gln Ile Ile Glu Thr Glu Gly Gly Ser Lys Pro Thr Val
 290 295 300
 Ser Arg Thr Arg Lys Arg Pro Thr Pro Ser Asn Ser Ala Asp Pro Lys
 305 310 315 320
 20 Ile Asn Tyr Lys Asp His Asn Lys Asn Val Val Tyr Ser Thr Thr Leu
 325 330 335
 Gln Glu His Gln Lys His Leu Gln Asn Arg Gly Leu Ala Leu Ser Ile
 340 345 350
 His Gln Ala Met Val Leu Arg Leu Arg Tyr Ile Ala Glu His Val Ile
 355 360 365
 25 Arg Ser Leu Asn Glu Phe Gly Trp Ala Val Val Asp Asn Phe Leu Gly
 370 375 380
 Ser Asp His Tyr Lys Phe Thr Ala Lys Glu Ile Glu Arg Leu Tyr Glu
 385 390 395 400
 Arg Gly Leu Phe Ser Pro Gly Gln Leu Met Glu Ala Lys His Lys Asp
 405 410 415
 30 Glu Phe His Ile Lys Asp Ile Arg Ser Asp His Ile Tyr Trp Tyr Asp
 420 425 430
 Gly Tyr Asp Gly Arg Ala Lys Asp Ala Ala Thr Val Arg Leu Leu Ile
 435 440 445
 35 Ser Met Ile Asp Ser Val Ile Gln His Phe Lys Lys Arg Ile Asp His
 450 455 460
 Asp Ile Gly Gly Arg Ser Arg Ala Met Leu Ala Ile Tyr Pro Gly Asn
 465 470 475 480
 40 Gly Thr Arg Tyr Val Lys His Val Asp Asn Pro Val Lys Asp Gly Arg
 485 490 495
 Cys Ile Thr Thr Ile Tyr Tyr Cys Asn Glu Asn Trp Asp Met Ala Thr
 500 505 510
 Asp Gly Gly Thr Leu Arg Leu Tyr Pro Glu Thr Ser Met Thr Pro Met
 515 520 525
 45 Asp Ile Asp Pro Arg Ala Asp Arg Leu Val Phe Phe Trp Ser Asp Arg
 530 535 540
 Arg Asn Pro His Glu Val Met Pro Val Phe Arg His Arg Phe Ala Ile
 545 550 555 560
 Thr Ile Trp Tyr Met Asp Lys Ser Glu Arg Asp Lys Ala Leu Ala Lys
 565 570 575
 50 Gly Lys Glu Ser Asp Ala Ala Cys Ala Ser Lys Lys Glu Asn Asp Pro

16/35

580 585 590
 Thr Ser Ser Ser Leu Asn Ser Leu Ile Gly Ser Leu Leu Arg Pro Arg
 595 600 605
 Lys Asn Pro Ser Thr His Asp Leu Ser Lys Leu Asp Leu Arg Leu Phe
 610 615 620
 5 Pro Ser Thr Ser Ser Asp Pro Ala Leu Val Ser Ala Ala Asp Glu Asp
 625 630 635 640
 Arg Val Asp Ile Ser Ala Asp Phe Gln Ser Thr Ser Ser Leu Ala His
 645 650 655
 10 Pro Glu Ser Thr Asp Ser Gly Val Ser Leu Ser Thr Phe Asn Val Ala
 660 665 670
 His Asn His Met Glu Arg Thr Thr Ser Leu Gln Ser Ile Ser Asp His
 675 680 685
 Phe Arg Ser Glu Arg Ser His Glu Arg Arg Ser Ser Thr Ser Ser Asp
 690 695 700
 15 Gln Asp Leu Asp Glu Gly Leu Pro Pro Pro Pro Ser Thr Asn Pro Glu
 705 710 715 720
 Tyr Tyr Ile

20

<210> 9
 <211> 19
 <212> PRT
 <213> artificial sequence

25

<220>
 <221> MISC_FEATURE
 <222> (9)..(9)
 <223> 4Hyp

30

<400> 9

Asp Leu Asp Leu Glu Met Leu Ala Xaa Tyr Ile Pro Met Asp Asp Asp
 1 5 10 15

35 Phe Gln Leu

<210> 10
 <211> 19
 <212> PRT
 <213> artificial sequence
 <220>
 <221> MISC_FEATURE
 <222> (9)..(9)
 <223> 4Hyp

45

<400> 10

Asp Leu Asp Leu Glu Met Leu Ala Xaa Tyr Ile Ser Met Asp Asp Asp
 1 5 10 15

50 Phe Gln Leu

17/35

<210> 11
<211> 19
<212> PRT
<213> artificial sequence
5 <220>
<221> MISC_FEATURE
<222> (9)..(9)
<223> 4Hyp

10 <400> 11

Asp Leu Asp Leu Glu Met Leu Leu Xaa Tyr Ile Pro Met Asp Asp Asp
1 5 10 15
Phe Gln Leu

15 <210> 12
<211> 19
<212> PRT
<213> artificial sequence
20 <220>
<221> MISC_FEATURE
<222> (9)..(9)
<223> 4Hyp

25 <400> 12

Asp Leu Asp Leu Glu Met Leu Val Xaa Tyr Ile Pro Met Asp Asp Asp
1 5 10 15
Phe Gln Leu

30 <210> 13
<211> 19
<212> PRT
<213> artificial sequence
35 <220>
<221> MISC_FEATURE
<222> (9)..(9)
<223> 4Hyp

40 <400> 13

Asp Leu Asp Leu Glu Met Ile Ala Xaa Tyr Ile Pro Met Asp Asp Asp
1 5 10 15
Phe Gln Leu

45 <210> 14
<211> 19
<212> PRT
<213> artificial sequence
50 <220>
<221> MISC_FEATURE

18/35

<222> (9)..(9)

<223> 4Hyp

<400> 14

5

Asp Leu Asp Leu Glu Met Ile Ala Xaa Tyr Ile Pro Met Glu Asp Asp
 1 5 10 15
 Phe Gln Leu

10

<210> 15

<211> 19

<212> PRT

<213> artificial sequence

<220>

15

<221> MISC_FEATURE

<222> (9)..(9)

<223> 4Hyp

<400> 15

20

Asp Leu Asp Leu Glu Met Leu Val Xaa Tyr Ile Ser Met Asp Asp Asp
 1 5 10 15
 Phe Gln Leu

25

<210> 16

<211> 34

<212> PRT

<213> artificial sequence

<400> 16

30

Pro Phe Ser Thr Gln Asp Thr Asp Leu Asp Leu Glu Met Leu Ala Pro
 1 5 10 15
 Tyr Ile Pro Met Asp Asp Asp Phe Gln Leu Arg Ser Phe Asp Gln Leu
 20 25 30

35

Ser Pro

<210> 17

<211> 6

<212> PRT

40

<213> artificial sequence motif

<220>

<221> MISC_FEATURE

<222> (2)..(3)

<223> X=any amino acid

45

<400> 17

Leu Xaa Xaa Leu Ala Pro
 1 5

50

<210> 18

19/35

<211> 5
 <212> PRT
 <213> artificial sequence motif
 <220>
 5 <221> MISC_FEATURE
 <222> (2)..(2)
 <223> X=any amino acid

 <220>
 10 <221> MISC_FEATURE
 <222> (4)..(4)
 <223> X=any amino acid and n=1 to 200

 <400> 18
 15 His Xaa Asp Xaa His
 1 5

 <210> 19
 20 <211> 62
 <212> PRT
 <213> artificial sequence motif
 <220>
 <221> MISC_FEATURE
 25 <222> (2)..(2)
 <223> X=any amino acid

 <220>
 <221> MISC_FEATURE
 30 <222> (4)..(61)
 <223> X=any amino acid

 <400> 19

 35 His Xaa Asp Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 1 5 10 15
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 20 25 30
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 40 35 40 45
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa His
 50 55 60

 <210> 20
 45 <211> 85
 <212> PRT
 <213> artificial sequence motif
 <220>
 <221> MISC_FEATURE
 50 <222> (2)..(14)
 <223> X=any amino acid

20/35

<220>
 <221> MISC_FEATURE
 <222> (16)..(16)
 5 <223> X=any amino acid

 <220>
 <221> MISC_FEATURE
 <222> (18)..(21)
 10 <223> X=any amino acid

 <220>
 <221> MISC_FEATURE
 <222> (23)..(29)
 15 <223> X=any amino acid

 <220>
 <221> MISC_FEATURE
 <222> (31)..(59)
 20 <223> X=any amino acid

 <220>
 <221> MISC_FEATURE
 <222> (61)..(70)
 25 <223> X=any amino acid

 <220>
 <221> MISC_FEATURE
 <222> (72)..(75)
 30 <223> X=any amino acid

 <220>
 <221> MISC_FEATURE
 <222> (77)..(77)
 <223> X=any amino acid
 35 <220>
 <221> MISC_FEATURE
 <222> (79)..(84)
 <223> X=any amino acid

 <400> 20
 40
 Met Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa His Xaa
 1 5 10 15
 Asp Xaa Xaa Xaa Xaa Asp Xaa Xaa Xaa Xaa Xaa Xaa Xaa Tyr Xaa Xaa
 20 25 30
 45 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Leu Xaa Xaa Xaa
 35 40 45
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Pro Xaa Xaa Xaa Xaa
 50 55 60
 Xaa Xaa Xaa Xaa Xaa Xaa Asp Xaa Xaa Xaa Xaa His Xaa Val Xaa Xaa
 50 65 70 75 80
 Xaa Xaa Xaa Xaa Arg

21/35

85

5 <210> 21
 <211> 110
 <212> PRT
 <213> artificial sequence motif
 <220>
 <221> MISC_FEATURE
 <222> (2)..(5)
10 <223> X=any amino acid

 <220>
 <221> MISC_FEATURE
 <222> (6)..(8)
15 <223> X=any amino acid

 <220>
 <221> MISC_FEATURE
 <222> (14)..(15)
20 <223> X=any amino acid

 <220>
 <221> MISC_FEATURE
 <222> (18)..(18)
25 <223> X=any amino acid

 <220>
 <221> MISC_FEATURE
 <222> (24)..(25)
30 <223> X=any amino acid

 <220>
 <221> MISC_FEATURE
 <222> (30)..(30)
35 <223> X=any amino acid

 <220>
 <221> MISC_FEATURE
 <222> (32)..(32)
40 <223> X=any amino acid

 <220>
 <221> MISC_FEATURE
 <222> (36)..(36)
45 <223> X=any amino acid

 <220>
 <221> MISC_FEATURE
 <222> (38)..(39)
50 <223> X=any amino acid

22/35

5 <220>
 <221> MISC_FEATURE
 <222> (42)..(45)
 <223> X=any amino acid

10 <220>
 <221> MISC_FEATURE
 <222> (48)..(48)
 <223> X=any amino acid

15 <220>
 <221> MISC_FEATURE
 <222> (50)..(51)
 <223> X=any amino acid

20 <220>
 <221> MISC_FEATURE
 <222> (55)..(63)
 <223> X=any amino acid

25 <220>
 <221> MISC_FEATURE
 <222> (65)..(66)
 <223> X=any amino acid

30 <220>
 <221> MISC_FEATURE
 <222> (70)..(70)
 <223> X=any amino acid

35 <220>
 <221> MISC_FEATURE
 <222> (72)..(72)
 <223> X=any amino acid

40 <220>
 <221> MISC_FEATURE
 <222> (77)..(83)
 <223> X=any amino acid

45 <220>
 <221> MISC_FEATURE
 <222> (85)..(88)
 <223> X=any amino acid

50 <220>
 <221> MISC_FEATURE
 <222> (90)..(90)
 <223> X=any amino acid

55 <220>
 <221> MISC_FEATURE
 <222> (92)..(92)

23/35

<223> X=any amino acid
 <220>
 <221> MISC_FEATURE
 5 <222> (97)..(97)
 <223> X=any amino acid
 <220>
 <221> MISC_FEATURE
 10 <222> (99)..(100)
 <223> X=any amino acid
 <220>
 <221> MISC_FEATURE
 15 <222> (103)..(104)
 <223> X=any amino acid
 <220>
 <221> MISC_FEATURE
 20 <222> (106)..(109)
 <223> X=any amino acid
 <400> 21
 25 Arg Xaa Xaa Xaa Met Xaa Xaa Xaa Tyr Pro Gly Asn Gly Xaa Xaa Tyr
 1 5 10 15
 Val Xaa His Val Asp Asn Pro Xaa Xaa Asp Gly Arg Cys Xaa Thr Xaa
 20 25 30
 Ile Tyr Tyr Xaa Asn Xaa Xaa Trp Asp Xaa Xaa Xaa Xaa Gly Gly Xaa
 30 35 40 45
 Leu Xaa Xaa Phe Pro Glu Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Pro
 50 55 60
 Xaa Xaa Asp Arg Leu Xaa Phe Xaa Trp Ser Asp Arg Arg Asn Pro His
 65 70 75 80
 35 Glu Val Xaa Pro Xaa Xaa Xaa Arg Xaa Ala Xaa Thr Val Trp Tyr
 85 90 95
 Xaa Asp Xaa Xaa Glu Arg Xaa Xaa Ala Xaa Ala Lys Xaa Lys
 100 105 110
 40 <210> 22
 <211> 6
 <212> PRT
 <213> artificial sequence motif
 <220>
 45 <221> MISC_FEATURE
 <222> (2)..(3)
 <223> X=any amino acid
 <220>
 50 <221> MISC_FEATURE
 <222> (5)..(5)

24/35

<223> X=any amino acid

<220>

<221> MISC_FEATURE

5 <222> (6)..(6)

<223> 4Hyp

<400> 22

10 Leu Xaa Xaa Leu Xaa Xaa
 1 5

<210> 23

<211> 19

15 <212> PRT

<213> Homo sapiens

<400> 23

20 Asp Leu Asp Leu Glu Met Leu Ala Pro Tyr Ile Pro Met Asp Asp Asp
 1 5 10 15
 Phe Gln Leu

<210> 24

<211> 20

25 <212> PRT

<213> Homo sapiens

<400> 24

30 Glu Leu Asp Leu Glu Thr Leu Ala Pro Tyr Ile Pro Met Asp Gly Glu
 1 5 10 15
 Asp Phe Gln Leu
 20

<210> 25

35 <211> 19

<212> PRT

<213> Xenopus laevis

<400> 25

40 Asp Leu Asp Leu Glu Met Leu Ala Pro Tyr Ile Pro Met Asp Asp Asp
 1 5 10 15
 Phe Gln Leu

<210> 26

45 <211> 19

<212> PRT

<213> Drosophila melanogaster

<400> 26

50 Phe Glu Ala Phe Ala Met Arg Ala Pro Tyr Ile Pro Ile Asp Asp Asp
 1 5 10 15

25/35

Met Pro Leu

<210> 27

<211> 19

5 <212> PRT

<213> Caenorhabditis elegans

<400> 27

10 Glu Pro Asp Leu Ser Cys Leu Ala Pro Phe Val Asp Thr Tyr Asp Met
 1 5 10 15
 Met Gln Met

<210> 28

<211> 34

15 <212> PRT

<213> artificial sequence

<220>

<223> Synthetic polypeptide

20 <400> 28

Pro Phe Ser Thr Gln Asp Thr Asp Leu Asp Leu Glu Met Leu Ala Pro
 1 5 10 15
 Tyr Ile Pro Met Asp Asp Asp Phe Gln Leu Arg Ser Phe Asp Gln Leu
 25 20 25 30
 Ser Pro

<210> 29

<211> 19

30 <212> PRT

<213> artificial sequence

<220>

<223> Synthetic polypeptide

35 <400> 29

Asp Leu Asp Leu Glu Met Leu Ala Pro Tyr Ile Pro Met Asp Asp Asp
 1 5 10 15
 Phe Gln Leu

40

<210> 30

<211> 16

<212> PRT

<213> artificial sequence

45 <220>

<223> Synthetic polypeptide

<400> 30

50 Leu Glu Met Leu Ala Pro Tyr Ile Pro Met Asp Asp Asp Phe Gln Leu
 1 5 10 15

26/35

<210> 31
<211> 13
<212> PRT
5 <213> artificial sequence
<220>
<223> Synthetic polypeptide

<400> 31
10
Leu Ala Pro Tyr Ile Pro Met Asp Asp Asp Phe Gln Leu
1 5 10

<210> 32
15 <211> 14
<212> PRT
<213> artificial sequence
<220>
<223> Synthetic polypeptide
20
<400> 32

Asp Leu Asp Leu Glu Met Leu Ala Pro Tyr Ile Pro Met Asp
1 5 10
25
<210> 33
<211> 19
<212> PRT
<213> artificial sequence
30 <220>
<223> Synthetic polypeptide

<400> 33

35 Asp Leu Asp Leu Glu Met Leu Ala Gly Tyr Ile Pro Met Asp Asp Asp
1 5 10 15
Phe Gln Leu

<210> 34
40 <211> 19
<212> PRT
<213> artificial sequence
<220>
<223> Synthetic polypeptide
45
<400> 34

Asp Leu Asp Leu Glu Met Leu Ala Pro Tyr Ile Pro Met Asp Asp Asp
1 5 10 15
50 Phe Gln Leu

27/35

<210> 35

<211> 826

<212> PRT

5 <213> Homo sapiens

<400> 35

10 Met Glu Gly Ala Gly Gly Ala Asn Asp Lys Lys Lys Ile Ser Ser Glu
 1 5 10 15
 Arg Arg Lys Glu Lys Ser Arg Asp Ala Ala Arg Ser Arg Arg Ser Lys
 20 25 30
 Glu Ser Glu Val Phe Tyr Glu Leu Ala His Gln Leu Pro Leu Pro His
 35 40 45
 15 Asn Val Ser Ser His Leu Asp Lys Ala Ser Val Met Arg Leu Thr Ile
 50 55 60
 Ser Tyr Leu Arg Val Arg Lys Leu Leu Asp Ala Gly Asp Leu Asp Ile
 65 70 75 80
 Glu Asp Asp Met Lys Ala Gln Met Asn Cys Phe Tyr Leu Lys Ala Leu
 20 85 90 95
 Asp Gly Phe Val Met Val Leu Thr Asp Asp Gly Asp Met Ile Tyr Ile
 100 105 110
 Ser Asp Asn Val Asn Lys Tyr Met Gly Leu Thr Gln Phe Glu Leu Thr
 115 120 125
 25 Gly His Ser Val Phe Asp Phe Thr His Pro Cys Asp His Glu Glu Met
 130 135 140
 Arg Glu Met Leu Thr His Arg Asn Gly Leu Val Lys Lys Gly Lys Glu
 145 150 155 160
 Gln Asn Thr Gln Arg Ser Phe Phe Leu Arg Met Lys Cys Thr Leu Thr
 30 165 170 175
 Ser Arg Gly Arg Thr Met Asn Ile Lys Ser Ala Thr Trp Lys Val Leu
 180 185 190
 His Cys Thr Gly His Ile His Val Tyr Asp Thr Asn Ser Asn Gln Pro
 195 200 205
 35 Gln Cys Gly Tyr Lys Lys Pro Pro Met Thr Cys Leu Val Leu Ile Cys
 210 215 220
 Glu Pro Ile Pro His Pro Ser Asn Ile Glu Ile Pro Leu Asp Ser Lys
 225 230 235 240
 Thr Phe Leu Ser Arg His Ser Leu Asp Met Lys Phe Ser Tyr Cys Asp
 40 245 250 255
 Glu Arg Ile Thr Glu Leu Met Gly Tyr Glu Pro Glu Glu Leu Leu Gly
 260 265 270
 Arg Ser Ile Tyr Glu Tyr Tyr His Ala Leu Asp Ser Asp His Leu Thr
 275 280 285
 45 Lys Thr His His Asp Met Phe Thr Lys Gly Gln Val Thr Thr Gly Gln
 290 295 300
 Tyr Arg Met Leu Ala Lys Arg Gly Gly Tyr Val Trp Val Glu Thr Gln
 305 310 315 320
 Ala Thr Val Ile Tyr Asn Thr Lys Asn Ser Gln Pro Gln Cys Ile Val
 50 325 330 335
 Cys Val Asn Tyr Val Val Ser Gly Ile Ile Gln His Asp Leu Ile Phe

28/35

			340				345				350					
	Ser	Leu	Gln	Gln	Thr	Glu	Cys	Val	Leu	Lys	Pro	Val	Glu	Ser	Ser	Asp
			355					360					365			
5	Met	Lys	Met	Thr	Gln	Leu	Phe	Thr	Lys	Val	Glu	Ser	Glu	Asp	Thr	Ser
		370					375					380				
	Ser	Leu	Phe	Asp	Lys	Leu	Lys	Lys	Glu	Pro	Asp	Ala	Leu	Thr	Leu	Leu
	385					390					395					400
	Ala	Pro	Ala	Ala	Gly	Asp	Thr	Ile	Ile	Ser	Leu	Asp	Phe	Gly	Ser	Asn
					405					410					415	
10	Asp	Thr	Glu	Thr	Asp	Asp	Gln	Gln	Leu	Glu	Glu	Val	Pro	Leu	Tyr	Asn
					420					425					430	
	Asp	Val	Met	Leu	Pro	Ser	Pro	Asn	Glu	Lys	Leu	Gln	Asn	Ile	Asn	Leu
			435					440					445			
15	Ala	Met	Ser	Pro	Leu	Pro	Thr	Ala	Glu	Thr	Pro	Lys	Pro	Leu	Arg	Ser
		450						455					460			
	Ser	Ala	Asp	Pro	Ala	Leu	Asn	Gln	Glu	Val	Ala	Leu	Lys	Leu	Glu	Pro
	465					470					475					480
	Asn	Pro	Glu	Ser	Leu	Glu	Leu	Ser	Phe	Thr	Met	Pro	Gln	Ile	Gln	Asp
					485					490					495	
20	Gln	Thr	Pro	Ser	Pro	Ser	Asp	Gly	Ser	Thr	Arg	Gln	Ser	Ser	Pro	Glu
					500					505					510	
	Pro	Asn	Ser	Pro	Ser	Glu	Tyr	Cys	Phe	Tyr	Val	Asp	Ser	Asp	Met	Val
			515					520					525			
25	Asn	Glu	Phe	Lys	Leu	Glu	Leu	Val	Glu	Lys	Leu	Phe	Ala	Glu	Asp	Thr
		530					535					540				
	Glu	Ala	Lys	Asn	Pro	Phe	Ser	Thr	Gln	Asp	Thr	Asp	Leu	Asp	Leu	Glu
	545					550					555					560
	Met	Leu	Ala	Pro	Tyr	Ile	Pro	Met	Asp	Asp	Phe	Gln	Leu	Arg	Ser	
					565					570					575	
30	Phe	Asp	Gln	Leu	Ser	Pro	Leu	Glu	Ser	Ser	Ser	Ala	Ser	Pro	Glu	Ser
					580					585					590	
	Ala	Ser	Pro	Gln	Ser	Thr	Val	Thr	Val	Phe	Gln	Gln	Thr	Gln	Ile	Gln
			595					600					605			
35	Glu	Pro	Thr	Ala	Asn	Ala	Thr	Thr	Thr	Thr	Ala	Thr	Thr	Asp	Glu	Leu
		610					615					620				
	Lys	Thr	Val	Thr	Lys	Asp	Arg	Met	Glu	Asp	Ile	Lys	Ile	Leu	Ile	Ala
	625					630					635					640
	Ser	Pro	Ser	Pro	Thr	His	Ile	His	Lys	Glu	Thr	Thr	Ser	Ala	Thr	Ser
					645					650					655	
40	Ser	Pro	Tyr	Arg	Asp	Thr	Gln	Ser	Arg	Thr	Ala	Ser	Pro	Asn	Arg	Ala
					660					665					670	
	Gly	Lys	Gly	Val	Ile	Glu	Gln	Thr	Glu	Lys	Ser	His	Pro	Arg	Ser	Pro
			675					680					685			
45	Asn	Val	Leu	Ser	Val	Ala	Leu	Ser	Gln	Arg	Thr	Thr	Val	Pro	Glu	Glu
		690					695						700			
	Glu	Leu	Asn	Pro	Lys	Ile	Leu	Ala	Leu	Gln	Asn	Ala	Gln	Arg	Lys	Arg
	705					710					715					720
	Lys	Met	Glu	His	Asp	Gly	Ser	Leu	Phe	Gln	Ala	Val	Gly	Ile	Gly	Thr
					725					730					735	
50	Leu	Leu	Gln	Gln	Pro	Asp	Asp	His	Ala	Ala	Thr	Thr	Ser	Leu	Ser	Trp
					740					745					750	

29/35

Lys Arg Val Lys Gly Cys Lys Ser Ser Glu Gln Asn Gly Met Glu Gln
 755 760 765
 Lys Thr Ile Ile Leu Ile Pro Ser Asp Leu Ala Cys Arg Leu Leu Gly
 770 775 780
 5 Gln Ser Met Asp Glu Ser Gly Leu Pro Gln Leu Thr Ser Tyr Asp Cys
 785 790 795 800
 Glu Val Asn Ala Pro Ile Gln Gly Ser Arg Asn Leu Leu Gln Gly Glu
 805 810 815
 10 Glu Leu Leu Arg Ala Leu Asp Gln Val Asn
 820 825

 <210> 36

 <211> 719
 15 <212> PRT
 <213> Caenorhabditis elegans

 <400> 36

 20 Met Glu Asp Asn Arg Lys Arg Asn Met Glu Arg Arg Arg Glu Thr Ser
 1 5 10 15
 Arg His Ala Ala Arg Asp Arg Arg Ser Lys Glu Ser Asp Ile Phe Asp
 20 25 30
 Asp Leu Lys Met Cys Val Pro Ile Val Glu Glu Gly Thr Val Thr His
 25 35 40 45
 Leu Asp Arg Ile Ala Leu Leu Arg Val Ala Ala Thr Ile Cys Arg Leu
 50 55 60
 Arg Lys Thr Ala Gly Asn Val Leu Glu Asn Asn Leu Asp Asn Glu Ile
 65 70 75 80
 30 Thr Asn Glu Val Trp Thr Glu Asp Thr Ile Ala Glu Cys Leu Asp Gly
 85 90 95
 Phe Val Met Ile Val Asp Ser Asp Ser Ser Ile Leu Tyr Val Thr Glu
 100 105 110
 Ser Val Ala Met Tyr Leu Gly Leu Thr Gln Thr Asp Leu Thr Gly Arg
 35 115 120 125
 Ala Leu Arg Asp Phe Leu His Lys Pro Ser Asp Tyr Asp Glu Phe Asp Lys
 130 135 140
 Gln Ser Lys Met Leu His Lys Pro Arg Gly Glu Asp Thr Asp Thr Thr
 145 150 155 160
 40 Gly Ile Asn Met Val Leu Arg Met Lys Thr Val Ile Ser Pro Arg Gly
 165 170 175
 Arg Cys Leu Asn Leu Lys Ser Ala Leu Tyr Lys Ser Val Ser Phe Leu
 180 185 190
 Val His Ser Lys Val Ser Thr Gly Gly His Val Ser Phe Met Gln Gly
 45 195 200 205
 Ile Thr Ile Pro Ala Gly Gln Gly Thr Thr Asn Ala Asn Ala Ser Ala
 210 215 220
 Met Thr Lys Tyr Thr Glu Ser Pro Met Gly Ala Phe Thr Thr Arg His
 225 230 235 240
 50 Thr Cys Asp Met Arg Ile Thr Phe Val Ser Asp Lys Phe Asn Tyr Ile
 245 250 255

30/35

Leu Lys Ser Glu Leu Lys Thr Leu Met Gly Thr Ser Phe Tyr Glu Leu
 260 265 270
 Val His Pro Ala Asp Met Met Ile Val Ser Lys Ser Met Lys Glu Leu
 275 280 285
 5 Phe Ala Lys Gly His Ile Arg Thr Pro Tyr Tyr Arg Leu Ile Ala Ala
 290 295 300
 Asn Asp Thr Leu Ala Trp Ile Gln Thr Glu Ala Thr Thr Ile Thr His
 305 310 315 320
 10 Thr Thr Lys Gly Gln Lys Gly Gln Tyr Val Ile Cys Val His Tyr Val
 325 330 335
 Leu Gly Ile Gln Gly Ala Glu Glu Ser Leu Val Val Cys Thr Asp Ser
 340 345 350
 Met Pro Ala Gly Met Gln Val Asp Ile Lys Lys Glu Val Asp Asp Thr
 355 360 365
 15 Arg Asp Tyr Ile Gly Arg Gln Pro Glu Ile Val Glu Cys Val Asp Phe
 370 375 380
 Thr Pro Leu Ile Glu Pro Glu Asp Pro Phe Asp Thr Val Ile Glu Pro
 385 390 395 400
 Val Val Gly Gly Glu Glu Pro Val Lys Gln Ala Asp Met Gly Ala Arg
 405 410 415
 20 Lys Asn Ser Tyr Asp Asp Val Leu Gln Trp Leu Phe Arg Asp Gln Pro
 420 425 430
 Ser Ser Pro Pro Pro Ala Arg Tyr Arg Ser Ala Asp Arg Phe Arg Thr
 435 440 445
 25 Thr Glu Pro Ser Asn Phe Gly Ser Ala Leu Ala Ser Pro Asp Phe Met
 450 455 460
 Asp Ser Ser Ser Arg Thr Ser Arg Pro Lys Thr Ser Tyr Gly Arg Arg
 465 470 475 480
 Ala Gln Ser Gln Gly Ser Arg Thr Thr Gly Ser Ser Ser Thr Ser Ala
 485 490 495
 30 Ser Ala Thr Leu Pro His Ser Ala Asn Tyr Ser Pro Leu Ala Glu Gly
 500 505 510
 Ile Ser Gln Cys Gly Leu Asn Ser Pro Pro Ser Ile Lys Ser Gly Gln
 515 520 525
 35 Val Val Tyr Gly Asp Ala Arg Ser Met Gly Arg Ser Cys Asp Pro Ser
 530 535 540
 Asp Ser Ser Arg Arg Phe Ser Ala Leu Ser Pro Ser Asp Thr Leu Asn
 545 550 555 560
 Val Ser Ser Thr Arg Gly Ile Asn Pro Val Ile Gly Ser Asn Asp Val
 565 570 575
 40 Phe Ser Thr Met Pro Phe Ala Asp Ser Ile Ala Ile Ala Glu Arg Ile
 580 585 590
 Asp Ser Ser Pro Thr Leu Thr Ser Gly Glu Pro Ile Leu Cys Asp Asp
 595 600 605
 45 Leu Gln Trp Glu Glu Pro Asp Leu Ser Cys Leu Ala Pro Phe Val Asp
 610 615 620
 Thr Tyr Asp Met Met Gln Met Asp Glu Gly Leu Pro Pro Glu Leu Gln
 625 630 635 640
 Ala Leu Tyr Asp Leu Pro Asp Phe Thr Pro Ala Val Pro Gln Ala Pro
 645 650 655
 50 Ala Ala Arg Pro Val His Ile Asp Arg Ser Pro Pro Ala Lys Arg Met

31/35

660 665 670
 His Gln Ser Gly Pro Ser Asp Leu Asp Phe Met Tyr Thr Gln His Tyr
 675 680 685
 Gln Pro Phe Gln Gln Asp Glu Thr Tyr Trp Gln Gly Gln Gln Gln
 5 690 695 700
 Asn Glu Gln Gln Pro Ser Ser Tyr Ser Pro Phe Pro Met Leu Ser
 705 710 715

 <210> 37
 10 <211> 38
 <212> PRT
 <213> Homo sapiens
 <400> 37

 15 Ser Glu Asp Thr Ser Ser Leu Phe Asp Lys Leu Lys Lys Glu Pro Asp
 1 5 10 15
 Ala Leu Thr Leu Leu Ala Pro Ala Ala Gly Asp Thr Ile Ile Ser Leu
 20 25 30
 Asp Phe Gly Ser Asn Asp
 20 35

 <210> 38
 <211> 38
 <212> PRT
 25 <213> Homo sapiens
 <400> 38

 Ser Glu Lys Ser Asn Phe Leu Phe Thr Lys Leu Lys Glu Glu Pro Glu
 1 5 10 15
 30 Glu Leu Ala Gln Leu Ala Pro Thr Pro Gly Asp Ala Ile Ile Ser Leu
 20 25 30
 Asp Phe Gly Asn Gln Asn
 35

 35 <210> 39
 <211> 19
 <212> PRT
 <213> Homo sapiens
 <400> 39

 40 Asp Leu Asp Leu Glu Met Leu Ala Pro Tyr Ile Pro Met Asp Asp Asp
 1 5 10 15
 Phe Gln Leu

 45 <210> 40
 <211> 119
 <212> PRT
 <213> Caenorhabditis elegans
 <400> 40

 50 Ile Asp His Asp Ile Gly Gly Arg Ser Arg Ala Met Leu Ala Ile Tyr

32/35

1 5 10 15
 Pro Gly Asn Gly Thr Arg Tyr Val Lys His Val Asp Asn Pro Val Lys
 20 25 30
 5 Asp Gly Arg Cys Ile Thr Thr Ile Tyr Tyr Cys Asn Glu Asn Trp Asp
 35 40 45
 Met Ala Thr Asp Gly Gly Thr Leu Arg Leu Tyr Pro Glu Thr Ser Met
 50 55 60
 Thr Pro Met Asp Ile Asp Pro Arg Ala Asp Arg Leu Val Phe Phe Trp
 65 70 75 80
 10 Ser Asp Arg Arg Asn Pro His Glu Val Met Pro Val Phe Arg His Arg
 85 90 95
 Phe Ala Ile Thr Ile Trp Tyr Met Asp Lys Ser Glu Arg Asp Lys Ala
 100 105 110
 Leu Ala Lys Gly Lys Glu Ser
 15 115

<210> 41
 <211> 116
 <212> PRT
 20 <213> Homo sapiens
 <400> 41

Gly Ser Tyr Lys Ile Asn Gly Arg Thr Lys Ala Met Val Ala Cys Tyr
 1 5 10 15
 25 Pro Gly Asn Gly Thr Gly Tyr Val Arg His Val Asp Asn Pro Asn Gly
 20 25 30
 Asp Gly Arg Cys Val Thr Cys Ile Tyr Tyr Leu Asn Lys Asp Trp Asp
 35 40 45
 Ala Lys Val Ser Gly Gly Ile Leu Arg Ile Phe Pro Glu Gly Lys Ala
 30 50 55 60
 Gln Phe Ala Asp Ile Glu Pro Lys Phe Asp Arg Leu Leu Phe Phe Trp
 65 70 75 80
 Ser Asp Arg Arg Asn Pro His Glu Val Gln Pro Ala Tyr Ala Thr Arg
 85 90 95
 35 Tyr Ala Ile Thr Val Trp Tyr Phe Asp Ala Asp Glu Arg Ala Arg Ala
 100 105 110
 Lys Val Lys Tyr
 115

40 <210> 42
 <211> 116
 <212> PRT
 <213> Homo sapiens
 <400> 42

45 Gly Ser Tyr Val Ile Asn Gly Arg Thr Lys Ala Met Val Ala Cys Tyr
 1 5 10 15
 Pro Gly Asn Gly Leu Gly Tyr Val Arg His Val Asp Asn Pro His Gly
 20 25 30
 50 Asp Gly Arg Cys Ile Thr Cys Ile Tyr Tyr Leu Asn Gln Asn Trp Asp
 35 40 45

33/35

Val Lys Val His Gly Gly Leu Leu Gln Ile Phe Pro Glu Gly Arg Pro
 50 55 60
 Val Val Ala Asn Ile Glu Pro Leu Phe Asp Arg Leu Leu Ile Phe Trp
 65 70 75 80
 5 Ser Asp Arg Arg Asn Pro His Glu Val Lys Pro Ala Tyr Ala Thr Arg
 85 90 95
 Tyr Ala Ile Thr Val Trp Tyr Phe Asp Ala Lys Glu Arg Ala Ala Ala
 100 105 110
 Lys Asp Lys Tyr
 10 115

 <210> 43
 <211> 115
 <212> PRT
 15 <213> Homo sapiens
 <400> 43

Lys Tyr Tyr Val Lys Glu Arg Ser Lys Ala Met Val Ala Cys Tyr Pro
 1 5 10 15
 20 Gly Asn Gly Thr Gly Tyr Val Arg His Val Asp Asn Pro Asn Gly Asp
 20 25 30
 Gly Arg Cys Ile Thr Cys Ile Tyr Tyr Leu Asn Lys Asn Trp Asp Ala
 35 40 45
 25 Lys Leu His Gly Gly Ile Leu Arg Ile Phe Pro Glu Gly Lys Ser Phe
 50 55 60
 Ile Ala Asp Val Glu Pro Ile Phe Asp Arg Leu Leu Phe Phe Trp Ser
 65 70 75 80
 Asp Arg Arg Asn Pro His Glu Val Gln Pro Ser Tyr Ala Thr Arg Tyr
 85 90 95
 30 Ala Met Thr Val Trp Tyr Phe Asp Ala Glu Glu Arg Ala Glu Ala Lys
 100 105 110
 Lys Lys Phe
 115

 35 <210> 44
 <211> 116
 <212> PRT
 <213> Rattus sp.
 <400> 44

40 Gly Lys Tyr Tyr Val Lys Glu Arg Ser Lys Ala Met Val Ala Cys Tyr
 1 5 10 15
 Pro Gly Asn Gly Thr Gly Tyr Val Arg His Val Asp Asn Pro Asn Gly
 20 25 30
 45 Asp Gly Arg Cys Ile Thr Cys Ile Tyr Tyr Leu Asn Lys Asn Trp Asp
 35 40 45
 Ala Lys Leu His Gly Gly Val Leu Arg Ile Phe Pro Glu Gly Lys Ser
 50 55 60
 Phe Val Ala Asp Val Glu Pro Ile Phe Asp Arg Leu Leu Phe Ser Trp
 50 65 70 75 80
 Ser Asp Arg Arg Asn Pro His Glu Val Gln Pro Ser Tyr Ala Thr Arg

34/35

85 90 95
 Tyr Ala Met Thr Val Trp Tyr Phe Asp Ala Glu Glu Arg Ala Glu Ala
 100 105 110
 Lys Lys Lys Phe
 115

5
 <210> 45
 <211> 92
 <212> PRT
 10 <213> Streptomyces sp.
 <400> 45

Phe Asp Gly Thr His Leu Gln Met Ala Arg Ser Arg Asn Leu Lys Asn
 1 5 10 15
 15 Ala Ile Val Ile Pro His Arg Asp Phe Val Glu Leu Asp Arg Glu Val
 20 25 30
 Asp Arg Tyr Phe Arg Thr Phe Met Val Leu Glu Asp Ser Pro Leu Ala
 35 40 45
 Phe His Ser Asn Glu Asp Thr Val Ile His Met Arg Pro Gly Glu Ile
 20 50 55 60
 Trp Phe Leu Asp Ala Ala Thr Val His Ser Ala Val Asn Phe Ser Glu
 65 70 75 80
 Ile Ser Arg Gln Ser Leu Cys Val Asp Phe Ala Phe
 85 90

25
 <210> 46
 <211> 37
 <212> DNA
 <213> artificial sequence
 30 <220>
 <223> forward primer

<400> 46
 gatttatcgt gcttggcagg attcgttgac acttatg

37

35
 <210> 47
 <211> 36
 <212> DNA
 <213> artificial sequence
 40 <220>
 <223> reverse primer

<400> 47
 gtgtcaacga atcctgccaac gcacgataaa tcaggc

36

45
 <210> 48
 <211> 6
 <212> PRT
 <213> artificial sequence motif
 50 <220>
 <221> MISC_FEATURE

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 02/01381

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N9/02 C07K7/08 G01N33/535

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 G01N C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, EPO-Internal, BIOSIS, PAJ, CHEM ABS Data, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 00 69908 A (MAXWELL PATRICK HENRY ;ISIS INNOVATION (GB); PUGH CHRISTOPHER WILL) 23 November 2000 (2000-11-23) the whole document ---	1-55
A	SHIBASAKI T ET AL: "Substrate Selectivities of Proline Hydroxylases" TETRAHEDRON LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 40, no. 28, 9 July 1999 (1999-07-09), pages 5227-5230, XP004170060 ISSN: 0040-4039 the whole document --- -/--	1-55

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

27 August 2002

Date of mailing of the international search report

19/09/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel: (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

G. Willièrè

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 02/01381

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MILLHORN D E ET AL: "REGULATION OF GENE EXPRESSION FOR TYROSINE HYDROXYLASE IN OXYGEN SENSITIVE CELLS BY HYPOXIA" KIDNEY INTERNATIONAL, NEW YORK, NY, US, vol. 51, no. 2, 1997, pages 527-535, XP000907384 ISSN: 0085-2538 the whole document	1-55
T	JAAKOLA ET AL.: "Targeting of HIF-alpha to the Hippel-Lindau ubiquitylation complex by O2-regulated prolyl hydroxylation" SCIENCE, vol. 292, no. 5516, 5 April 2001 (2001-04-05), pages 468-472, XP002211146 the whole document	1-55

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/GB 02/01381

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 0069908	A	23-11-2000	EP	1179018 A1	13-02-2002
			WO	0069908 A1	23-11-2000